



Biomarkers >

Volume 14, 2009 - [Issue 2](#)


419 | 156

Views | CrossRef citations to date | 3

Altmetric

Research Article

Biological half-life of cadmium in the urine of inhabitants after cessation of cadmium exposure

Yasushi Suwazono , Teruhiko Kido, Hideaki Nakagawa, Muneko Nishijo, Ryumon Honda, Etsuko Kobayashi, ... [show all](#)

Pages 77-81 | Received 01 Sep 2008, Accepted 07 Jan 2009, Published online: 01 Mar 2009

 Cite this article  <https://doi.org/10.1080/13547500902730698>

Sample our
Bioscience
Journals
>> [Sign in here](#) to start your access
to the latest two volumes for 14 days



 Full Article

 Figures & data

 References

 Citations

 Metrics

 Reprints & Permissions

[Read this article](#)

 Share

Abstract

We investigated the biological half-life of the urinary cadmium concentration (U-Cd) based on a 24-year follow-up study after cessation of cadmium exposure in a cadmium-polluted area. Spot urine samples were obtained from all inhabitants in this area in 1979, 1986, 1991, 1999 and 2003. Biological half-life was calculated in the inhabitants whose U-Cd was more than $5 \mu\text{g l}^{-1}$ (9 men and 12 women) or $5 \mu\text{g g}^{-1}$ creatinine (9 men and 19 women) using a one-compartment model. The estimated half-life and 95% confidence intervals were 13.6 years (9.0–28.2 years) and 13.9 years (9.6–25.6 years) for unadjusted U-Cd in men and women, respectively. For creatinine-adjusted U-Cd, they were 14.2 years (11.2–19.4 years) and 23.5 years (17.7–35.0 years) in men and

women, respectively. The biological half-lives of U-Cd obtained in this study were identical with the values of total body burden determined by a different method.

Keywords::

- Urinary cadmium
- biological half-life
- long-term follow-up study
- risk assessment
- human

Acknowledgment

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Related Research Data

- URINARY CADMIUM ELIMINATION AS A BIOMARKER OF EXPOSURE FOR EVALUATING A CADMIUM DIETARY EXPOSURE - BIOKINETICS MODEL

Source: Journal of Toxicology and Environmental Health Part A

The relation of individual cadmium concentration in urine with total cadmium intake in Kakehashi River basin, Japan

Source: Toxicology Letters
- In vivo measurement of liver and kidney cadmium in workers exposed to this metal: Its significance with respect to cadmium in blood and urine

Source: Environmental Research

Cadmium Metabolism in Man

Source: Human Toxicology
- Significance of Urinary Cadmium Concentration in a Japanese Population Environmentally Exposed to Cadmium

Source: Archives of Environmental Health An International Journal

A comparison between fecal cadmium and urinary β 2-microglobulin, total protein, and

Related research

Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2025 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)

 Taylor and Francis Group

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG