



Biomarkers >

Volume 14, 2009 - Issue 2

427 | 158 | 3
Views | CrossRef citations to date | Altmetric

Research Article

Biological half-life of cadmium in the urine of inhabitants after cessation of cadmium exposure

Yasushi Suwazono , Teruhiko Kido, Hideaki Nakagawa, Muneko Nishijo, Ryumon Honda, Etsuko Kobayashi, ... [show all](#)

Pages 77-81 | Received 01 Sep 2008, Accepted 07 Jan 2009, Published online: 01 Mar 2009

[Cite this article](#) <https://doi.org/10.1080/13547500902730698>

Sample our Medicine, Dentistry, Nursing & Allied Health Journals
>> [Sign in here](#) to start your access to the latest two volumes for 14 days

[Full Article](#)

[Figures & data](#)

[References](#)

[Citations](#)

[Metrics](#)

[Reprints & Permissions](#)

[Read this article](#)

[Share](#)

Abstract

We investigated the biological half-life of the urinary cadmium concentration (U-Cd) based on a 24-year follow-up study after cessation of cadmium exposure in a cadmium-polluted area. Spot urine samples were obtained from all inhabitants in this area in 1979, 1986, 1991, 1999 and 2003. Biological half-life was calculated in the inhabitants whose U-Cd was more than $5 \mu\text{g l}^{-1}$ (9 men and 12 women) or $5 \mu\text{g g}^{-1}$ creatinine (9 men and 19 women) using a one-compartment model. The estimated half-life and 95% confidence intervals were 13.6 years (9.0–28.2 years) and 13.9 years (9.6–25.6 years) for unadjusted U-Cd in men and women, respectively. For creatinine-adjusted U-Cd, they were 14.2 years (11.2–19.4 years) and 23.5 years (17.7–35.0 years) in men and

women, respectively. The biological half-lives of U-Cd obtained in this study were identical with the values of total body burden determined by a different method.

Keywords::

Urinary cadmium

biological half-life

long-term follow-up study

risk assessment

human

Acknowledgment

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Related Research Data

URINARY CADMIUM ELIMINATION AS A BIOMARKER OF EXPOSURE FOR EVALUATING A CADMIUM DIETARY EXPOSURE - BIOKINETICS MODEL

Source: Journal of Toxicology and Environmental Health Part A

The relation of individual cadmium concentration in urine with total cadmium intake in Kakehashi River basin, Japan

Source: Toxicology Letters

In vivo measurement of liver and kidney cadmium in workers exposed to this metal: Its significance with respect to cadmium in blood and urine

Source: Environmental Research

Cadmium Metabolism in Man

Source: Human Toxicology

Significance of Urinary Cadmium Concentration in a Japanese Population

Environmentally Exposed to Cadmium

Source: Archives of Environmental Health An International Journal

A comparison between fecal cadmium and urinary β 2-microglobulin, total protein, and

Related research

People also read

Recommended articles

Cited by
158

Information for	Open access
Authors	Overview
R&D professionals	Open journals
Editors	Open Select
Librarians	Dove Medical Press
Societies	F1000Research
Opportunities	Help and information
Reprints and e-prints	Help and contact
Advertising solutions	Newsroom
Accelerated publication	All journals
Corporate access solutions	Books

Keep up to date

Register to receive personalised research and resources
by email

 Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)
[Accessibility](#)



Registered in England & Wales No. 01072954
 5 Howick Place | London | SW1P 1WG