

Pharmaceutical Biology >

Volume 45, 2007 - [Issue 4](#)

 Free access

50,255 | 29 | 0
Views | CrossRef citations to date | Altmetric

Listen

Research Article

Effect of Leaves and Fruits of *Moringa oleifera*. on Gastric and Duodenal Ulcers

V.C. Devaraj, Mohammed Asad & Satya Prasad

Pages 332-338 | Accepted 13 Nov 2006, Published online: 07 Oct 2008

 Cite this article  <https://doi.org/10.1080/13880200701212924>

 Full Article

 Figures & data

 References

 Citations

 Metrics

 Reprints & Permissions

 View PDF

 Share

Abstract

The effect of different extracts of leaves and fruits of *Moringa oleifera*. Lam. (Moringaceae) on gastric and duodenal ulcers was evaluated by using different gastric ulcer models and cysteamine-induced duodenal ulcer method. The leaf extracts (500 mg/kg, p.o.) of *Moringa oleifera*. showed gastric ulcer healing effect in acetic acid-induced chronic gastric ulcers. The acetone extract and methanol extract of the leaves produced gastric antisecretory effect in pylorus-ligated rats and showed gastric cytoprotective effect in ethanol-induced and indomethacin-induced gastric ulcers. The leaf extracts also produced a significant reduction of stress-induced gastric ulcers and cysteamine-induced duodenal ulcers. None of the extracts of the fruits showed any significant antiulcer effect. It was concluded that leaves of *Moringa oleifera*. increase healing of gastric ulcers and also prevent the development of experimentally induced gastric ulcers and duodenal ulcers in rats.

Introduction

Peptic ulcer is a breach in the gastric and duodenal epithelium associated with acute or chronic inflammation and is the most common gastrointestinal disorder in clinical practice. In spite of established antiulcer drugs, a rational therapy for peptic ulcer remains elusive, and a search for safer potential drugs is being carried out. The use of natural drugs in gastric ulcer has been reported (Sairam et al., [2001](#)).

The development and progression of gastric ulcer depends to some extent on the type of the food consumed by the patient. It has been shown that spicy food, fatty food, or food containing caffeine stimulates acid secretion in the stomach (Crawford, [2003](#)), and high-fiber diets such as potatoes, bananas, peas, beans, and so forth, reduce the development of duodenal ulcers (Maryland Medical Center Programs, [2004](#)).

Previous reports on the incidence of gastric ulcers in the South Asian population reveals that the occurrence is lower due to the type of food consumed by the people of this region; one of the foods that is speculated to protect against ulcers is *Moringa oleifera* Lam. (Moringaceae) leaves (Jayaraj et al., [1998](#)). Furthermore, flower bud of *Moringa pterygosperma*, a synonym of *Moringa oleifera*. that is widely consumed in Pakistan, has been reported to possess antiulcer activity against aspirin-induced gastric ulcers in rats (Akhtar & Ahmad, [1995](#)). However, the effect of different extracts of leaves or fruits of *Moringa oleifera*. on gastric and duodenal ulcers is not known. The current study was undertaken to evaluate the effect of fruits and leaves of *Moringa oleifera*. on experimentally induced gastric and duodenal ulcers.

Materials and Methods

Collection of plant material and extraction procedure

The leaves and fruits of *Moringa oleifera*. were collected from the Madiwala area of Bangalore, India, in the month of May and were authenticated by the Regional Research Institute (Bangalore, India). A voucher specimen (RRI/BNG/SMP-Prog/800) has been preserved in the Regional Research Institute for future reference.

The shade-dried leaves and fruits of the plant were subjected to Soxhlet extraction, and the extracts thus obtained were subjected to preliminary phytochemical analysis (Mukaherjee, [2002](#)).

Experimental animals

Albino Wistar rats weighing between 200 and 250 g were used. The institutional animal ethics committee approved the experimental protocol. Animals were maintained under standard conditions in an animal house approved by the Committee for the Purpose of Control, and Supervision on Experiments on Animals (CPCSEA), Government of India.

Acute toxicity study

The acute oral toxicity study was performed according to the OPPTS (Office of Prevention, Pesticide and Toxic Substance) Up and Down procedure (Health Effect Test Guideline, [2004](#)). The different extracts were suspended using 0.5% sodium carboxy methylcellulose and were administered orally. All the extracts of leaves and fruits were safe at a dose of 5000 mg/kg, p.o., and one-tenth of this dose was selected for evaluation of antiulcer activity.

Effect on gastric ulcers

Acetic acid-induced chronic gastric ulcer

Male Wistar rats were fasted for 24 h prior to the experiment. Under light ether anesthesia, a midline epigastric incision was made and the stomach was exposed. Glacial acetic acid (0.05 mL) was applied topically onto the serosal surface using a cylindrical mold (6.0 mm internal diameter), which was allowed to remain there for 60 s. The acid solution was then removed by rinsing the mold with normal saline to prevent possible damage to the surrounding tissues close to the point of application. The animals were treated with different extracts of leaves and fruits of *Moringa oleifera*. (500 mg/kg, p.o.) once daily for 10 days after induction of ulcer while the control group received only vehicle. Rats were sacrificed on the 10th day by overdose of ether

anesthesia; stomach was removed and cut open along the greater curvature (Okabe et al., [1970](#)).

The stomach samples were scanned using a computer scanner, and the total mucosal area and total ulcerated area were measured using public domain image processing and analysis program developed at National Institutes of Health (Bethesda, MD, USA). The PC version of the program was downloaded free from Scion (<http://www.scioncorp.com>; Scion Image for Windows, release beta 4.0.2). The scale was set at 6.1 pixels per millimeter.

The ulcer index was determined using the formula (Ganguly, [1969](#))

where $X = \text{total mucosal area} / \text{total ulcerated area}$.

The ulcers were given scores based on their intensity as follows: 0 = no ulcer, 1 = superficial mucosal erosion, 2 = deep ulcer or transmural necrosis, 3 = perforated or penetrated ulcer.

The stomach samples from groups that showed reduction in ulcer index were subsequently processed for histologic examination. Three indices, namely regenerated glandular epithelial width, capillary density in the scar tissue, and volume of collagen content were selected to reflect the rate and quality of ulcer healing (Wang et al., [1999](#)).

Pylorus ligation-induced ulcers

The animals were fasted for 36 h with water ad libitum. before pylorus ligation (Shay et al., [1945](#); Kulkarni, [1999](#)). Normal saline (1 mL/rat, p.o.) was administered twice daily to all the animals during the fasting period (Asad et al., [2003](#)). Under light ether anesthesia, the pyloric portion of the stomach was ligated. The leaf extracts (500 mg/kg, p.o.) were administered intraduodenally immediately after pylorus ligation. Six hours after pylorus ligation, animals were sacrificed. The stomach was isolated and the content of the stomach was used for estimation of free and total acidity (Hawk et al., [1947](#)), pepsin content (Debnath et al., [1974](#)), total hexoses (Goel et al., [1985](#)), and total proteins (Lowry et al., [1951](#)). The ulcer index was determined using the same method as mentioned earlier.

Healing of indomethacin-induced gastric ulcers

The gastric ulcers were induced by administering indomethacin (5 mg/kg, p.o.) for 5 days to the normally fed animals (Majumdar et al., [2003](#)). The animals were then treated with different extracts of leaves of *Moringa oleifera*. (500 mg/kg, p.o.) once daily for 5 days after induction of ulcer while the control group received only vehicle. The last dose of indomethacin was considered day 0. Rats were sacrificed on the day 0 and day 5. The stomach was removed; ulcer score and ulcer index were determined. The glandular portion of the stomach was then used for estimation of mucin content (Corne et al., [1974](#)), total proteins (Lowry et al., [1951](#)), antioxidant factors like superoxide dismutase activity (Eltner & Heupel, [1976](#)), total tissue sulfhydryl groups (Ellman, [1959](#)), and catalase activity (Link, [1988](#)).

Ethanol-induced ulcers

Male albino rats were fasted for 36 h before administration of 90% ethanol (1 mL/200 g). The leaf extracts (500 mg/kg, p.o.) were administered 1 h before ethanol administration. One hour after ethanol administration, the animals were sacrificed, stomach was isolated, and ulcer index was determined (Brzozowski et al., [1998](#)).

Cold restraint stress-induced ulcers

The animals were placed in a restraint cage, and the cage was placed at a temperature of 2°C for 3 h. The leaf extracts (500 mg/kg, p.o.) were administered 30 min prior to subjection of stress. The animals were sacrificed, stomach was isolated, and ulcer index was determined (Vincent et al., [1977](#)).

Effect on duodenal ulcers

Cysteamine-induced duodenal ulcers

Duodenal ulcers were induced by administering cysteamine hydrochloride (400 mg/kg, p.o.) twice at an interval of 4 h. Leaf extracts (500 mg/kg, p.o.) were administered 30 min prior to each dose of cysteamine hydrochloride. After 24 h of the first dose of cysteamine, animals were sacrificed and the duodenum was excised carefully and cut opened along the antimesenteric side. The duodenal ulcer area, ulcer score, and ulcer index were determined (Szabo, [1978](#)).

The ulcers were given scores based on their intensity as follows: 0 = no ulcer, 1 = superficial mucosal erosion, 2 = deep ulcer or transmural necrosis, 3 = perforated or

The ulcer index (UI) was calculated using the following equation (Desai et al., [1999](#)):

Statistical analysis

The statistical significance was assessed using one-way analysis of variance (ANOVA) followed by Dunnett's comparison test. For comparing nonparametric ulcer scores, ANOVA followed by non-parametric Dunn post test was used. The values are expressed as mean \pm SEM, and $p < 0.05$ was considered significant.

Results

Acetic acid-induced chronic gastric ulcers

The petroleum ether, acetone, and methanol leaf extracts of *Moringa oleifera*. showed a significant reduction in ulcer index when compared with control ($p < 0.001$). The acetone extract of the leaves was most potent; it produced a 79% decrease in the ulcer index. The fruit extracts of *Moringa oleifera*. and the chloroform extract of the leaves did not show any significant effect on ulcer index ([Fig. 1](#)). None of the treatments produced any significant effect on ulcer score ([Fig. 2](#)).

Figure 1 Effect of *Moringa oleifera*. on ulcer index in acetic acid-induced chronic gastric ulcer. All values are mean \pm SEM, $n = 5$ to 6 . * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ when compared with control group. Cont, control; pet leaf, petroleum ether leaf extract (500 mg/kg p.o.); chcl leaf, chloroform leaf extract (500 mg/kg p.o.); act leaf, acetone leaf extract (500 mg/kg p.o.); meth leaf, methanol leaf extract (500 mg/kg); pet frt, petroleum ether fruit extract (500 mg/kg p.o.); chcl frt, chloroform fruit extract (500 mg/kg p.o.); act frt, acetone fruit extract (500 mg/kg); meth frt, methanol fruit extract (500 mg/kg p.o.).





Figure 2 Effect of *Moringa oleifera*. on ulcer score in acetic acid-induced chronic gastric ulcer. All values are mean \pm SEM, n = 5 to 6. *p < 0.05, **p < 0.01, ***p < 0.001 when compared with control group. Cont, control; pet leaf, petroleum ether leaf extract (500 mg/kg p.o.); chcl leaf, chloroform leaf extract (500 mg/kg p.o.); act leaf, acetone leaf extract (500 mg/kg p.o.); meth leaf, methanol leaf extract (500 mg/kg); pet frt, petroleum ether fruit extract (500 mg/kg p.o.); chcl frt, chloroform fruit extract (500 mg/kg p.o.); act frt, acetone fruit extract (500 mg/kg), meth frt, methanol fruit extract (500 mg/kg p.o.).



[Display full size](#)

Histologic examination of the ulcerated area revealed that there was a significant increase in regenerated glandular epithelium width after treatment with acetone extract of the leaves ($p < 0.05$) when compared with control. The collagen content in the ulcerated tissue was significantly increased by all the three extracts of the leaves, with acetone extract showing the maximum effect. No significant difference on capillary density in scar tissue was observed after treatment ([Table 1](#)).

Table 1. Effect of *Moringa oleifera*. Lam. on regenerated glandular epithelium width, capillary density, and volume of collagen content in acetic acid-induced chronic gastric ulcers



[Download CSV](#)

[Display Table](#)

Pylorus ligation-induced gastric ulcers

The acetone leaf extracts of *Moringa oleifera*. showed a significant reduction in free acidity and total acidity when compared with control ($p < 0.05$) whereas the methanol

of the treatments produced any significant effect on ulcer index, total hexose content, mucin content, pepsin activity, and total protein ([Table 2](#)).

[Table 2..](#) Effect of *Moringa oleifera*. leaf extracts on free acidity, total acidity, ulcer index, and total hexoses in pylorus-ligated rats



[Download CSV](#)

[Display Table](#)

Healing of indomethacin-induced gastric ulcers

The acetone and methanol leaf extracts of *Moringa oleifera*. showed a significant reduction in ulcer index ($p < 0.01$) and a significant increase in mucus content when compared with control. None of the treatments produced any significant effect on total protein and antioxidant factors like SOD activity, total tissue sulfhydryl group (glutathione), and catalase activity ([Table 3](#)).

[Table 3..](#) Effect of *Moringa oleifera*. leaf extracts on indomethacin-induced ulcers



[Download CSV](#)

[Display Table](#)

Ethanol-induced gastric ulcers

The acetone and methanol leaf extracts of *Moringa oleifera*. showed a significant reduction in ulcer index when compared with control ($p < 0.05$). The petroleum ether extract did not show significant reduction in ulcer index when compared with control ([Table 4](#)).

[Table 4..](#) Effect of *Moringa oleifera*. leaf extracts in ethanol-induced gastric ulcers, stress-induced gastric ulcers, and cysteamine-induced duodenal ulcers



[Download CSV](#)

[Display Table](#)

Stress-induced gastric ulcers

All the three leaf extracts of *Moringa oleifera*. showed a significant reduction in ulcer index when compared with control. Again, the acetone extract of the leaves was most potent in reducing ulcer index ([Table 4](#)).

Cysteamine-induced duodenal ulcers

The acetone leaf extract of *Moringa oleifera*. showed a highly significant reduction in ulcer area when compared with control ($p < 0.01$). The methanol leaf extracts of *Moringa oleifera*. showed a significant reduction in ulcer area when compared with that of control ($p < 0.05$). The petroleum ether leaf extract did not show any significant reduction in ulcer area. None of the treatments produced any significant effect on ulcer score and ulcer index ([Table 4](#)).

Discussion

The current study dealt with the effect of different extracts of *Moringa oleifera*. leaves and fruits on the gastric and duodenal ulcers in rats. The extracts that showed ulcer-healing effect in acetic acid-induced gastric ulcers were screened further to determine their effect on gastric cytoprotection and gastric secretion.

Application of glacial acetic acid (0.05 mL) on to the serosal surface of the rat stomach produces deep penetrating gastric ulcers (Okabe et al., [1971](#)). The fruit extracts of the plant did not show any effect on the healing of gastric ulcers induced by acetic acid, whereas the petroleum ether, acetone, and methanol leaf extracts of *Moringa oleifera*. increased healing of the acetic acid-induced gastric ulcers. The histopathologic examination was carried out to determine the effect of extracts on regeneration of glandular epithelium, formation of collagen, and capillary density, all of which are essential processes for the healing of ulcers. The acetone extract of the leaves increased the collagen content and regeneration of glandular epithelium, whereas the methanol and petroleum ether extract of the leaves produced only an increase in collagen content.

Pylorus ligation-induced ulcer was used to study the effect of extracts on gastric secretion. The ligation of the pyloric end of the stomach causes accumulation of gastric acid in the stomach that produces ulcers in the stomach. Agents that decrease gastric

induced by this method. The acetone and methanol extracts of *Moringa oleifera*. produced reduction in gastric acid secretion proving their antisecretory effect.

Ethanol-induced and indomethacin-induced gastric ulcer was employed to study the cytoprotective effect of the extracts. The acetone and methanol extracts of *Moringa oleifera*. leaves were effective in reducing ulcer index in both these models and significantly increased the mucus content in indomethacin-induced gastric ulcers. However, the leaf extracts of *Moringa oleifera*. were not effective in altering the antioxidant factors like SOD activity, total tissue sulfhydryl group (glutathione), and catalase activity suggesting that the healing of ulcers or prevention of development of gastric ulcers may not be due to antioxidant effect.

Stress plays an important role in ulcerogenesis. The pathophysiology of stress-induced gastric ulcers is complex. Stress-induced ulcers are probably mediated by histamine release with enhancement in acid secretion and a reduction in mucus production (Brodie & Hanson, [1960](#); Peters & Richardson, [1983](#)). The petroleum ether, acetone, and methanol extracts of *Moringa oleifera*. were effective in reducing the ulcers induced by stress. Cysteamine-induced duodenal ulcer in rat is a widely used model of peptic ulcer disease. Cysteamine hydrochloride inhibits the alkaline mucus secretion from the Brunner glands in the proximal duodenum and stimulates gastric acid secretion rate. Gastric emptying is also delayed, and serum gastrin concentration is increased (Parmar & Desai, [1993](#)). The acetone and methanol leaf extracts of *Moringa oleifera*. were effective in reducing the ulcer area in cysteamine-induced duodenal ulcers.

The acetone and methanol extracts of the leaves were effective in all the tested models of peptic ulcer disease, whereas the petroleum ether was effective only in healing chronic gastric ulcers and stress-induced gastric ulcers. Hence, the healing of gastric ulcers in acetic acid-induced gastric ulcers may be due to decreased acid secretion, increased mucus secretion, or decreased GI motility in case of acetone and methanol extracts, and the ulcer healing in case of petroleum ether may be mainly due to reduction in gastric motility.

Moringa oleifera. contains a number of flavonoids, triterpenes, steroids, alkaloids, and many other chemical constituents (Ross, [1999](#)). The flavonoid quercetin present in the leaves is a well-known antiulcer agent (Suzuki et al., [1998](#)). Further, the leaves contain

[2000](#)). This explains more potent ulcer healing effect of acetone and methanol extracts of the leaves compared with petroleum extract. Apart from flavonoids, the leaves of the plant contain steroids such as β -sitosterol and β -carotene, and both of these are known to reduce the development of gastric ulcers (Xiao et al., [1992](#)). This probably explains the activity of petroleum ether extract.

The results of the current study suggest that consumption of the leaves of *Moringa oleifera*. may be beneficial in healing of ulcers in patients suffering from peptic ulcer disease.

Acknowledgment

The authors are thankful to Prof. Suresh Nagpal, chairman, Krupanidhi Educational Trust, and Prof. Sunil Dhaminigi, secretary, Krupanidhi Educational Trust, for providing facilities to carry out the work. The authors are also thankful to Prof. Dr. Amit Kumar Das, principal, Krupanidhi College of Pharmacy, for his valuable suggestions during the work.

Related Research Data

[A Study of the Factors Involved in the Production of Gastric Ulcers by the Restraint Technique](#)

Source: Gastroenterology

[PROTEIN MEASUREMENT WITH THE FOLIN PHENOL REAGENT](#)

Source: Journal of Biological Chemistry

[Evidence for protective and antioxidant properties of rutin, a natural flavone, against ethanol induced gastric lesions](#)

Source: Journal of Ethnopharmacology

[The mechanism of pH-dependent hydrogen peroxide cytotoxicity in vitro](#)

Source: Archives of Biochemistry and Biophysics

References

1. Akhtar AH, Ahmad KU (1995): Anti-ulcerogenic evaluation of the methanolic extracts of some indigenous medicinal plants of Pakistan in aspirin-ulcerated rats. *J Ethnopharmacol* 46: 1-6.

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

2. Asad M, Rajesh KS, Ahmad M (2003): Effect of administration of saline or glucose during fasting on the development of gastric ulcers in pylorus ligated rats. *Ethiop Pharm J* 16: 1-6.

[Google Scholar](#)

3. Brodie DAQ, Hanson HM (1960): A study of the factors involved in the production of gastric ulcers by the restraint technique. *Gastroenterology* 38: 353-360.

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

4. Brzozowski T, Konturek SJ, Kwiecien S, Pajdo R, Brzozowski I, Hahn EG (1998): Involvement of endogenous cholecystikinin and somatostatin in gastro protection induced by intra duodenal fat. *J Clin Gastroenterol* 27: 125-137.

[Web of Science ®](#)

[Google Scholar](#)

5. Casa CL, Villegas I, Lastra CA, Motilva V, Calero MJM (2000): Evidence for protective and antioxidant properties of rutin, a natural flavone, against ethanol induced gastric lesions. *J Ethnopharmacol* 71: 45-53.

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

6. Corne SJ, Morrissey SM, Woods RJ (1974): A method for the quantitative estimation of gastric barrier mucus. *J Physiol* 242: 116-117.

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

7. Crawford JM (2003): Robbin's Pathologic Basis of Disease. New Delhi, India, Saunders, pp. 787-802.

[Google Scholar](#)

8. Debnath PK, Gode KO, Govinda DD, Sanyal AK (1974): Effect of propranolol on gastric secretion in rats. Br J Pharmacol 51: 213-216.

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

9. Desai JK, Goyal RK, Parmar NS (1999): Characterization of dopamine receptor subtypes involved in experimentally induced gastric and duodenal ulcers in rats. J Pharm Pharmacol 51: 187-192.

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

10. Ellman GL (1959): Tissue sulfhydryl groups. Arch Biochem Biophys 82: 70-77.

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

11. Elstner EF, Heupel A (1976): Inhibition of nitrite formation from hydroxyl ammonium chloride. A simple assay for super oxide dismutase. Anal Biochem 70: 616-620.

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

12. Ganguly AK (1969): A method for quantitative assessment of experimentally produced ulcers in stomach of rats. Experientia 25: 1224.

[PubMed](#) | [Google Scholar](#)

13. Goel RK, Das DG, Sanyal AK (1985): Effect of vegetable banana powder on changes induced by ulcerogenic agents in dissolved mucosubstances of gastric juice. Indian J Gastroenterol 4: 249-251.

[PubMed](#) | [Google Scholar](#)

14. Hawk PB, Oser BL, Summerson HW (1947): Practical Physiological Chemistry, 12th ed. London, Churchill, p. 347.

5. Health Effect Test Guidelines (2004): Acute Oral Toxicity [computer program], OPPTS 870, 1100 United States Office of Prevention, Pesticides and Toxic Substances, Environmental Protection Agency (7101). Available at <http://www.epa.gov/opptsfrs/home/guidelin.htm>.

[Google Scholar](#)

6. Jayaraj AP, Tovey FI, Clark CG (1998): Possible dietary protective factors in relation to the distribution of duodenal ulcer in India and Bangladesh. *Gut* 21: 1068–1076.

[Google Scholar](#)

7. Kulkarni SK (1999): *Handbook of Experimental Pharmacology*, 3rd ed. New Delhi, India, Vallabh Prakashan, pp. 148–150.

[Google Scholar](#)

8. Link EM (1988): Mechanism of pH dependent hydrogen peroxide cytotoxicity in vitro.. *Arch Biochem Biophys* 265: 362–372.

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

9. Lowry CH, Roseborough NI, Farr AL, Randall RJ (1951): Protein measurement with Folin phenol reagent. *J Biol Chem* 193: 265–275.

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

10. Majumdar B, Chaudhri SGR, Ray A, Bandyopadhyay SK (2003): Effect of ethanol extract of Piper betle. Linn leaf on healing of NSAID-induced experimental ulcer—A novel role of free radical scavenging action. *Indian J Exp Biol* 41: 311–315.

[PubMed](#) | [Google Scholar](#)

11. Maryland Medical Center Programs (2004): Peptic ulcers. Available at http://www.umm.edu/patiented/articles/what_lifestyle_changes_recommended_peptic_ulcers_dyspepsia_000019_11.htm

[Google Scholar](#)

22. Mukaherjee PK (2002): Quality Control of Herbal Drugs (An Approach to Evaluation of Botanicals). New Delhi, Business Horizons, pp. 380-421.

[Google Scholar](#)

23. Okabe S, Pfeiffer CJ, Roth JL (1970): Experimental production of duodenal and antral ulcers in rats. Federal Proc 29: 255.

[Google Scholar](#)

24. Okabe S, Roth JLA, Pfeiffer CJ (1971): Differential-healing periods of the acetic acid ulcer model in rats and cats. Experientia 27: 146-148.

[PubMed](#) | [Google Scholar](#)

25. Parmar NS, Desai JK (1993): A review of the current methodology for the evaluation of gastric and duodenal anti-ulcer agents. Indian J Pharmacol 25: 120-135.

[Google Scholar](#)

26. Peters MN, Richardson CT (1983): Stressful life events, acid hyper secretion, and ulcer disease. Gastroenterology 84: 114-119.

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

27. Ross IA (1999): Medicinal plants of the World. New Jersey, Humana Press, 234-235.

[Google Scholar](#)

28. Sairam K, Rao ChV, Goel R K (2001): Effect of Centella asiatica. Linn on physical and chemical factors induced gastric ulceration and secretion in rats. Indian J Exp Biol 39: 137-142.

[PubMed](#) | [Google Scholar](#)

29. Shay H, Komarov SA, Fele SS, Meranze D, Gruenstein H, Siple H (1945): A simple method for uniform production of gastric ulceration in rat. Gastroenterology 5: 43-61.

[Web of Science ®](#) | [Google Scholar](#)

30. Suzuki Y, Ishihara M, Segami T, Ito M (1998): Anti-ulcer effects of antioxidants, quercetin, α -tocopherol, nifedipine and tetracycline in rats. *Jpn J Pharmacol* 78: 435-441.

[PubMed](#) | [Google Scholar](#)

31. Szabo S (1978): Animal model for human disease: Duodenal ulcer disease. *Am J Pathol* 73: 273-276.

[Google Scholar](#)

32. Vincent GP, Galvin GB, Rutkowski JL, Pare WP (1977): Body orientation, food deprivation and potentiation of restraint induced gastric lesions. *Gastroenterol Clin Biol* 1: 539-543.

[PubMed](#) | [Google Scholar](#)

33. Wang JZ, Wuy J, Rao CM, Gao MT, Li WG (1999): Effect of recombinant human basic fibroblast growth factor on stomach ulcers in rats and mice. *Acta Pharmacol Sin* 209: 763-768.

[Google Scholar](#)

34. Xiao M, Yang Z, Jiu M, You J, Xiao R (1992): The anti-gastroulcerative activity of β -sitosterol- β -D-glucoside and its aglycone in rats. *Hua Xi Yi Ke Da Xue Xue Bao* 23(1): 98-101.

[PubMed](#) | [Google Scholar](#)

[Download PDF](#)

Related research

People also read

Recommended articles

Cited by
29

Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email




Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG

 Taylor and Francis
Group

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)