Taylor & Francis Online

Home ▶ All Journals ▶ Environment & Agriculture ▶ International Journal of Agricultural Sustainability ▶ List of Issues ▶ Volume 17, Issue 2 ▶ Estimating loss of ecosystem services du ....

International Journal of Agricultural Sustainability ›

Volume 17, 2019 - Issue 2

969 Views | 70 CrossRef citations to date | 3 Altmetric

Articles

Estimating loss of ecosystem services due to paddy straw burning in North-west India

Suresh Kumar, D. K. Sharma, D. R. Singh, H. Biswas, K. V. Praveen & Vikas Sharma

Pages 146-157 | Published online: 18 Feb 2019

Cite this article https://doi.org/10.1080/14735903.2019.1581474 Check for updates

Sample our Environment & Agriculture Journals
>> Sign in here to start your access to the latest two volumes for 14 days

Full Article Figures & data References Citations Metrics

Reprints & Permissions Read this article Share

ABSTRACT

Crop residue burning is severe in rice–wheat cropping system of North-western states (Punjab, Haryana, Uttarakhand, and western Uttar Pradesh) of India, where mechanized harvesting of rice using combine harvesters is a common practice, and management of leftover residue in the short window of 10–15 days for timely sowing of wheat is a formidable task. Moreover, there is a lack of user-friendly, cost-effective, and economically viable options and, around 23 million tonnes of rice residue is burnt annually in the region. Burning biomass not only pollutes environment but also results in loss of appreciable amount of plant's essential nutrients. Straw burning releases soot particles, nitrogen oxides, sulphur dioxide, carbon dioxide, carbon monoxide, and polycyclic aromatic hydrocarbons, thus causing serious deterioration in atmospheric quality and human health hazards. We attempted to identify and quantify the

environmental cost of paddy straw burning in North-west India. Using extant coefficients, it is estimated that cost of paddy residue burning is INR (Indian National Rupee) 8953 per ha, and the social cost of burning is INR 3199 crores per annum in the region.

KEYWORDS:

Crop residue burning | ecosystem services | North-west India

Disclosure statement

No potential conflict of interest was reported by the authors.

Related research

People also read | Recommended articles | Cited by 70

Crop residue management options in rice–rice system: a review

Subhendu Bikash Goswami et al.
Archives of Agronomy and Soil Science
Published online: 12 Sep 2019

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email

Sign me up

Copyright © 2026 Informa UK Limited Privacy policy

Cookies Terms & conditions Accessibility

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG

Taylor & Francis by informa