



International Journal of Agricultural Sustainability >

Volume 17, 2019 - Issue 2

931 | 61 | 0  
Views | CrossRef citations to date | Altmetric

Articles

# Estimating loss of ecosystem services due to paddy straw burning in North-west India

Suresh Kumar , D. K. Sharma, D. R. Singh, H. Biswas, K. V. Praveen & Vikas Sharma

Pages 146-157 | Published online: 18 Feb 2019

Cite this article <https://doi.org/10.1080/14735903.2019.1581474>

Check for updates

Sample our   
Environment and Sustainability  
Journals  
>> [Sign in here](#) to start your access  
to the latest two volumes for 14 days

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

 Read this article

Share

## ABSTRACT

Crop residue burning is severe in rice-wheat cropping system of North-western states (Punjab, Haryana, Uttarakhand, and western Uttar Pradesh) of India, where mechanized harvesting of rice using combine harvesters is a common practice, and management of leftover residue in the short window of 10–15 days for timely sowing of wheat is a formidable task. Moreover, there is a lack of user-friendly, cost-effective, and economically viable options and, around 23 million tonnes of rice residue is burnt annually in the region. Burning biomass not only pollutes environment but also results in loss of appreciable amount of plant's essential nutrients. Straw burning releases soot particles, nitrogen oxides, sulphur dioxide, carbon dioxide, carbon monoxide, and polycyclic aromatic hydrocarbons, thus causing serious deterioration in atmospheric quality and human health hazards. We attempted to identify and quantify the

environmental cost of paddy straw burning in North-west India. Using extant coefficients, it is estimated that cost of paddy residue burning is INR (Indian National Rupee) 8953 per ha, and the social cost of burning is INR 3199 crores per annum in the region.

KEYWORDS:

Crop residue burning    ecosystem services    North-west India

---

## Disclosure statement

No potential conflict of interest was reported by the authors.

---

## Related research

People also read

Recommended articles

Cited by  
61

## Information for

Authors

R&D professionals

Editors

Librarians

Societies

## Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

## Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

## Help and information

Help and contact

Newsroom

All journals

Books

## Keep up to date

Register to receive personalised research and resources  
by email

 Sign me up

  

  

Copyright © 2026 Informa UK Limited Privacy policy Cookies Terms & conditions

Accessibility



Registered in England & Wales No. 01072954  
5 Howick Place | London | SW1P 1WG