



The Journal of Maternal-Fetal & Neonatal Medicine >

Volume 34, 2021 - Issue 11

817 | 5 | 2  
Views | CrossRef citations to date | Altmetric

Original Articles

# Clinical and economic outcomes of cesarean deliveries with skin closure through skin staples plus waterproof wound dressings versus 2-octyl cyanoacrylate plus polymer mesh tape

Stephen S. Johnston , Brian Po-Han Chen, Akhil Nayak, Stephanie Hsiao Yu. Lee, Michelle Costa & Giovanni A. Tommaselli

Pages 1711-1720 | Received 19 Dec 2018, Accepted 16 Jul 2019, Accepted author version posted online: 17 Jul 2019, Published online: 01 Aug 2019

Cite this article <https://doi.org/10.1080/14767058.2019.1645830>

Check for updates

Sample our Medicine, Dentistry, Nursing & Allied Health Journals  
>> [Sign in here](#) to start your access to the latest two volumes for 14 days

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

Read this article

Share

## Abstract

## Purpose

To compare clinical and economic outcomes of cesarean deliveries with skin closure through skin staples plus waterproof wound dressings (SSWWD) versus 2-octyl cyanoacrylate plus polymer mesh tape (2OPMT). We hypothesized that cesarean deliveries with skin closure through 2OPMT may be associated with a lower rate of

wound complications and infections as compared with skin closure through SSWWD; we also hypothesized that, accordingly, 2OPMT may be associated with lower hospital length of stay (LOS), hospital costs, and all-cause readmissions as compared with SSWWD.

## Methods

Retrospective, observational study using a research database derived from administrative records routinely contributed by hundreds of hospitals in the USA. We queried the database for patients aged 18-49 years who had an in-hospital low transverse cesarean delivery between 1 January, 2012 and 31 March, 2017. Using records of medical supplies used during deliveries, we identified deliveries for which skin closure was performed by either SSWWD (SSWWD group) or 2OPMT (2OPMT group). Our primary study outcome was a composite endpoint of infection/wound complication diagnosis during the hospital stays in which the deliveries were performed. Our secondary outcomes included: length of stay (LOS) and total hospital costs for the hospital stays in which the deliveries were performed, and all-cause readmissions (30/60/90 days post discharge) to the same hospital in which the delivery was performed. We compared outcomes between propensity-score matched groups using regressions accounting for hospital-level clustering and non-Gaussian empirical outcome distributions.

## Results

Each group comprised 2133 patients (4266 total patients; mean age = 30.3 years [SD = 4.6]). Compared with the SSWWD group, the 2OPMT group had statistically significant lower rates of complications (infection, 0.7 versus 1.6%,  $p = .011$ ; wound complication, 0.6 versus 1.3%,  $p = .036$ ; composite, 0.9 versus 2.0%,  $p = .002$ ), shorter LOS (mean = 3.5 days [SD = 1.6] versus 3.7 days [SD = 1.8],  $p = .007$ ), and lower total hospital costs (mean = \$8879 [SD = \$3157] versus \$9313 [SD = \$3311],  $p = .025$ ). Between-group differences for 30/60/90-day all-cause readmissions were statistically insignificant.

## Conclusions

This large observational study is the first of its kind and provides evidence that cesarean delivery skin closure with 2OPMT is associated with lower rates of in-hospital

infection and wound complications, lower LOS, lower total hospital costs as compared with SSWWD.

Keywords:

2-octyl cyanoacrylate    cesarean delivery    skin closure    skin staples    infection, wound complications

---

## Acknowledgments

We acknowledge the contributions of Niels Derrek-Schmitz, Julia Ting, Geethu Roshan, and Sushama Ramisetti, employees of Johnson & Johnson at the time this study was conducted.

---

## Disclosure statement

Stephen S. Johnston, Brian Po-Han Chen, Stephanie Hsiao Yu Lee, Michelle Costa, and Giovanni A. Tommaselli are employees of Johnson & Johnson. Akhil Nayak is an employee of Mu Sigma, which is a paid consultant to Johnson & Johnson.

---

## Data availability statement

Although we are contractually unable to make the specific dataset on which this study was based publicly available, the overall Premier Healthcare Database is a commercially-available research database.

---

## Additional information

### Funding

This study was funded and conducted by Johnson & Johnson.

---

## Related research

People also read

Recommended articles

Cited by  
5

### Information for

Authors

R&D professionals

Editors

Librarians

Societies

### Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

### Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

### Help and information

Help and contact

Newsroom

All journals

Books

### Keep up to date

Register to receive personalised research and resources  
by email

 Sign me up



Copyright © 2026 Informa UK Limited Privacy policy Cookies Terms & conditions

Accessibility

 Taylor and Francis Group

Registered in England & Wales No. 01072954  
5 Howick Place | London | SW1P 1WG