

641 | 2 | 2
Views | CrossRef citations to date | Altmetric

Original Articles

Clinical and economic outcomes of cesarean deliveries with skin closure through skin staples plus waterproof wound dressings versus 2-octyl cyanoacrylate plus polymer mesh tape

Stephen S. Johnston , Brian Po-Han Chen, Akhil Nayak, Stephanie Hsiao Yu. Lee, Michelle Costa & Giovanni A. Tommaselli

Pages 1711-1720 | Received 19 Dec 2018, Accepted 16 Jul 2019, Published online: 01 Aug 2019

Cite this article <https://doi.org/10.1080/14767058.2019.1645830>

Check for updates

Sample our
Medicine, Dentistry, Nursing
& Allied Health Journals
>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

Full Article

Figures & data

References

Citations

Metrics

Reprints

We Care About Your Privacy

We and our 845 partners store and/or access information on a device, such as unique IDs in cookies to process personal data. You may accept or manage your choices by clicking below, including your right to object where legitimate interest is used, or at any time in the privacy policy page. These choices will be signaled to our partners and will not affect browsing data. [Privacy Policy](#)

We and our partners process data to provide:

Use precise geolocation data. Actively scan device characteristics for identification. Store and/or access information on a device. Personalised advertising and content, advertising and content measurement, audience research and services development.

[List of Partners \(vendors\)](#)

I Accept

Essential Only

Show Purpose closure



Methods

Retrospective, observational study using a research database derived from administrative records routinely contributed by hundreds of hospitals in the USA. We queried the database for patients aged 18–49 years who had an in-hospital low transverse cesarean delivery between 1 January, 2012 and 31 March, 2017. Using records of medical supplies used during deliveries, we identified deliveries for which skin closure was performed by either SSWWD (SSWWD group) or 2OPMT (2OPMT group). Our primary study outcome was a composite endpoint of infection/wound complication diagnosis during the hospital stays in which the deliveries were performed. Our secondary outcomes included: length of stay (LOS) and total hospital costs for the hospital stays in which the deliveries were performed, and all-cause readmissions (30/60/90 days post discharge) to the same hospital in which the delivery was performed. We compared outcomes between propensity-score matched groups using regressions accounting for hospital-level clustering and non-Gaussian empirical outcome distributions.

Results

Each group comprised 2133 patients (4266 total patients; mean age = 30.3 years [SD = 4.6]). Compared with the SSWWD group, the 2OPMT group had statistically significant lower rates of complications (infection, 0.7 versus 1.6%, $p = .011$; wound complication, 0.6 versus 1.3%, $p = .036$; composite, 0.9 versus 2.0%, $p = .002$), shorter LOS (mean = 3.5 days [SD = 1.6] versus 3.7 days [SD = 1.8], $p = .007$), and lower total hospital costs (mean = \$8879 [SD = \$3157] versus \$9313 [SD = \$3311], $p = .025$).

Between-group differences in readmissions were not statistically significant.

Conclusion

This large study found that 2OPMT was associated with statistically significant lower rates of infection, wound complication, and LOS compared with SSWWD.



Q Keyword

infection,

Acknowledgments

We acknowledge the contributions of Niels Derrek-Schmitz, Julia Ting, Geethu Roshan, and Sushama Ramiseti, employees of Johnson & Johnson at the time this study was conducted.

Disclosure statement

Stephen S. Johnston, Brian Po-Han Chen, Stephanie Hsiao Yu Lee, Michelle Costa, and Giovanni A. Tommaselli are employees of Johnson & Johnson. Akhil Nayak is an employee of Mu Sigma, which is a paid consultant to Johnson & Johnson.

Data availability statement

Although we are contractually unable to make the specific dataset on which this study was based publicly available, the overall Premier Healthcare Database is a commercially-available research database.

Additional information

Fundin

This stud



Rela

Effect of delivery

cesarean

Jill M. Westcott et al.

Perineal skin tear repair following vaginal birth; skin adhesive versus conventional suture – a randomised controlled trial >

Rima Anggrenas Dasrilsyah et al.
Journal of Obstetrics and Gynaecology
Published online: 12 Jun 2020

Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive updates by email



Sign up



Copyright

Accessibility

Registered
5 Howick Place



for & Francis Group
orma business