

HIV Clinical Trials >
Volume 19, 2018 - Issue 4

Free access

427 | 2 | 0
Views | CrossRef citations to date | Altmetric[Listen](#)

Brief Report

Short-term folic acid supplementation and aerobic exercise improve vascular reactivity in HIV-infected individuals

Shana S. Grigoletti, Jorge P. Ribeiro, Eduardo Sprinz & Paula A. B. Ribeiro

Pages 148-151 | Published online: 05 Feb 2018

 Cite this article <https://doi.org/10.1080/15284336.2018.1433769>

Check for updates

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

View PDF

Abstract

The aim of this study was to determine the effect of supervised exercise and folic acid supplementation on endothelial function in HIV-infected individuals. A randomized clinical trial, double blinded, was conducted with 16 HIV-infected individuals, antiretroviral therapy (at least 6 months) with undetectable viral load (<50 copies/mL), and CD4 count > 200 cells/mm³. The subjects were randomized to aerobic exercise (n = 5) and control (n = 11).

About Cookies On This Site

We and our partners use cookies to enhance your website experience, learn how our site is used, offer personalised features, measure the effectiveness of our services, and tailor content and ads to your interests while you navigate on the web or interact with us across devices. You can choose to accept all of these cookies or only essential cookies. To learn more or manage your preferences, click "Settings". For further information about the data we collect from you, please see our [Privacy Policy](#).

Accept All

Essential Only

Settings



endothelial function in HIV-infected individuals. As exercise and folate supplementation are safe and relatively inexpensive, this finding deserves more attention in large randomized clinical trials in an attempt to reduce cardiovascular risk in HIV-infected population.

Keywords: Exercise training | Endothelial function | HIV

Introduction

The antiretroviral therapy (ART) has dramatically reduced the morbidity and mortality associated to HIV.¹ On the other hand, with longer life expectancy, metabolic chronic non-communicable diseases are more prevalent in patients under ART therapy when compared to naïve patients.^{2,3} In this scenario, cardiovascular disease (CVD) is one of the leading causes of morbidity and mortality in such population,⁴⁻⁶ and the possible mechanisms of raised incidence could be related to endothelial dysfunction.⁷⁻¹² Therefore, we believe that it should be important to identify and manage individuals with endothelial dysfunction early on, which compromises up to 35% of these patients.^{13,14}

Our previous report has shown that supplementation with folates improves endothelium-dependent vasodilatation in HIV patients.¹⁵ However, non-farmacologic safe strategies¹⁶ have been considered to reduce cardiovascular complications, such as exercise as an additional therapy, since the ARV treatment involves increased cardiovascular risks. Previously, acute responses to exercise demonstrated potential benefits.¹⁷ Also, chronic effects (exercise training) have been used to improve endothelial function in patients such as obese^{18,19} and peripheral arterial disease patients.

About Cookies On This Site

We and our partners use cookies to enhance your website experience, learn how our site is used, offer personalised features, measure the effectiveness of our services, and tailor content and ads to your interests while you navigate on the web or interact with us across devices. You can choose to accept all of these cookies or only essential cookies. To learn more or manage your preferences, click "Settings". For further information about the data we collect from you, please see our [Privacy Policy](#).

Accept All

Essential Only

Settings

and aerobic exercise training would improve vascular reactivity, as assessed by brachial artery reactive hyperemia in HIV-infected people. For that, we conducted a pilot randomized, double-blind, placebo-controlled trial, with HIV-infected individuals, aged 18y or over, on ART for at least 6 months, with undetectable viral load (less than 50 copies/ml), and CD4 counts more than 200 cells/mm³. Exclusion criteria were diabetes mellitus, active infection, liver, kidney or cardiovascular diseases, pregnancy, and illicit drug use. All regular medications were maintained. The protocol was approved by the committee for ethics in research of the Hospital de Clínicas de Porto Alegre, according to Declaration of Helsinki, and all participants gave written informed consent.

Population and methods

Participants were randomly assigned to a 4-week treatment with either folinic acid ($n = 6$) or placebo ($n = 5$) both blinded to the researcher and participants as described before,¹⁵ and a not blinded arm of exercise group ($n = 5$). The primary outcome measure was the assessment of brachial artery vascular responses during reactive hyperemia. Secondary outcomes were the changes on the biochemical and hemodynamic variables. All outcomes were blinded for the evaluators and statistician.

In short, reactive hyperemia (RH) was induced by placing a cuff in the upper-arm at 250 mm Hg, and releasing after 5 min, in dominant forearm. All flow recordings were manually traced by an operator who was blinded to the groups, and time. After 15 min of rest, 2.5 mg of sublingual isosorbide dinitrate (Isordil®/Sigma/Brazil) was administered as an endothelium-independent vasodilator. Five minutes later, endothelium-independent vasodilatation of the brachial artery was measured.²¹ Folinic acid and placebo capsules were indistinguishable in shape, size, as well as color. They



About Cookies On This Site

We and our partners use cookies to enhance your website experience, learn how our site is used, offer personalised features, measure the effectiveness of our services, and tailor content and ads to your interests while you navigate on the web or interact with us across devices. You can choose to accept all of these cookies or only essential cookies. To learn more or manage your preferences, click “Settings”. For further information about the data we collect from you, please see our [Privacy Policy](#).

Accent All


Essential Oils

Settings

Our results showed no significant change in hemodynamic and biochemical variables. Likewise, blood glucose, lipid profile, creatinine, and vitamin B₁₂ did not change significantly in the three groups after intervention (data not showed).

Figure 1 shows the delta of reactive hyperemia response and endothelium-dependent vasodilation responses (blue bars) after the 4 weeks. Exercise improves reactive hyperemia (9.4–15.9 mL/min/100 mL) and folic acid (11.9–19.2 mL/min/100 mL). The same was not observed in the placebo group (from 18.7 to 18.4 ml/min/100 ml, time p < 0.001, interaction p = 0.02). The endothelium-independent responses (green bars) remained unchanged in both groups (groups p = 0.67; interaction p = 0.73). No adverse effects were observed.

Figure 1 Deltas and SE of reactive hyperemia (FMD), endothelium-dependent and independent, after 4-weeks intervention in HIV patients.



[Display full size](#)

About Cookies On This Site

We and our partners use cookies to enhance your website experience, learn how our site is used, offer personalised features, measure the effectiveness of our services, and tailor content and ads to your interests while you navigate on the web or interact with us across devices. You can choose to accept all of these cookies or only essential cookies. To learn more or manage your preferences, click "Settings". For further information about the data we collect from you, please see our [Privacy Policy](#).

[Accept All](#)

[Essential Only](#)

[Settings](#)

The main intervention was similar to knowledge of aerobic exercise in this paper.



could be easily implemented in the clinical site routine. However, non-supervised exercise effect (counseling and home-based strategies) still needs to be tested in these patients.

In conclusion, our preliminary results demonstrate that short-term administration methylated folate derivative from folic acid, and aerobic exercise training improve endothelium-dependent vasodilatation in HIV-infected individuals. This finding deserves more attention in larger randomized clinical trials, and future studies should also evaluate whether prolonged interventions would be effective to improve cardiovascular risks and major cardiovascular events in these population.

Contributors

SSG, ES, JPR, PABR conceived and designed the experiments. SSG was responsible for data collection. SSG, PABR analyzed data. SSG, PABR wrote the paper. The Final review was done by ES, PABR.

Funding

This work was supported by CAPES; CNPq, Brasília, Brazil; FIPE-HCPA, Porto Alegre, Brazil through scholarships.

Acknowledgments

In memoriam of Dr Jorge Pinto Ribeiro that was the first mentor of this study. This study was supported by CAPES, CNPq, FIPE-HCPA, and FAPERGS. We thank the patients and their families for their participation in this study. We also thank the nursing staff of the Hospital de Clínicas de Porto Alegre (HCPA) for their support and help during the data collection process.



About Cookies On This Site

We and our partners use cookies to enhance your website experience, learn how our site is used, offer personalised features, measure the effectiveness of our services, and tailor content and ads to your interests while you navigate on the web or interact with us across devices. You can choose to accept all of these cookies or only essential cookies. To learn more or manage your preferences, click "Settings". For further information about the data we collect from you, please see our [Privacy Policy](#).

Accept All

Essential Only

Settings

References

1. Sterne JA, Hernán MA, Ledergerber B, et al. Long-term effectiveness of potent antiretroviral therapy in preventing AIDS and death: a prospective cohort study. Lancet. 2005; 366(9483): 378–84.10.1016/S0140-6736(05)67022-5
- | PubMed | Web of Science ® | Google Scholar
2. Trevisol F, Alencastro P, Ribeiro P, et al. Association of physical activity with lipodystrophy syndrome in HIV-infected patients. J AIDS Clin Res. 2012; 3(8): 1.
- | Google Scholar
3. Islam FM, Wu J, Jansson J, Wilson DP. Relative risk of cardiovascular disease among people living with HIV: a systematic review and meta-analysis. HIV Med. 2012; 13(8): 453–68.
- | PubMed | Web of Science ® | Google Scholar
4. Grinspoon S, Carr A. Cardiovascular risk and body-fat abnormalities in HIV-infected adults. N Engl J Med. 2005; 352(1): 48–62.10.1056/NEJMra041811
- | PubMed | Web of Science ® | Google Scholar
5. Friis-Møller N, Sabin CA, Weber R, et al. Combination antiretroviral therapy and the risk of myocardial infarction. N Engl J Med. 2003; 349(21): 1993–2003.
- | PubMed | Web of Science ® | Google Scholar
6. Benjamin EJ, Blaha MJ, Chiuve SE, et al. Heart disease and stroke statistics-2017 update: a report from the American Heart Association. Circulation. 2017; 135(10): e146–603.10.1161/CIR.0000000000000485
- | PubMed | Web of Science ® | Google Scholar
7. Blum J, ...
immun...
dysf...
 About Cookies On This Site
We and our partners use cookies to enhance your website experience, learn how our site is used, offer personalised features, measure the effectiveness of our services, and tailor content and ads to your interests while you navigate on the web or interact with us across devices. You can choose to accept all of these cookies or only essential cookies. To learn more or manage your preferences, click "Settings". For further information about the data we collect from you, please see our [Privacy Policy](#).
- Accept All
Essential Only
Settings
8. Solage C, ...
Clin In...
- In this article

9. Stein JH, Klein MA, Bellehumeur JL, et al. Use of human immunodeficiency virus-1 protease inhibitors is associated with atherogenic lipoprotein changes and endothelial dysfunction. *Circulation*. 2001; **104**(3): 257-62.10.1161/01.CIR.104.3.257

| PubMed | Web of Science ® | Google Scholar

10. Blanco JJ, Garcia IS, Cerezo JG, et al. Endothelial function in HIV-infected patients with low or mild cardiovascular risk. *J Antimicrob Chemother*. 2006; **58**(1): 133-9.10.1093/jac/dkl190

| PubMed | Web of Science ® | Google Scholar

11. de Larrañaga GF, Petroni A, Deluchi G, Alonso BS, Benetucci JA. Viral load and disease progression as responsible for endothelial activation and/or injury in human immunodeficiency virus-1-infected patients. *Blood Coagul Fibrinolysis*. 2003; **14**(1): 15-18.10.1097/00001721-200301000-00004

| PubMed | Web of Science ® | Google Scholar

12. Monsuez JJ, Dufaux J, Vittecoq D, Vicaut E. Reduced reactive hyperemia in HIV-infected patients. *J Acquir Immune Defic Syndr*. 2000; **25**(5): 434-42.10.1097/00126334-200012150-00009

| PubMed | Web of Science ® | Google Scholar

13. Balsam P, Mikuła T, Peller M, et al. Evaluation of endothelial function and arterial stiffness in HIV-infected patients: a pilot study. *Kardiol Pol*. 2015; **73**(5): 344-51.10.5603/KP.a2014.0231

| PubMed | Web of Science ® | Google Scholar

14. Currier JS, Lundgren JD, Corr A, et al. Epidemiological evidence for cardiovascular

About Cookies On This Site

We and our partners use cookies to enhance your website experience, learn how our site is used, offer personalised features, measure the effectiveness of our services, and tailor content and ads to your interests while you navigate on the web or interact with us across devices. You can choose to accept all of these cookies or only essential cookies. To learn more or manage your preferences, click "Settings". For further information about the data we collect from you, please see our [Privacy Policy](#).

Accept All

Essential Only

Settings

5. Grigoriadis E, et al. Cardiovascular effects of lipid-lowering therapy in patients with cardiovascular disease: a systematic review and meta-analysis. *Arch Intern Med*. 2005; **165**(11): 1289-1299.10.1001/archinte.155.11.1289

In this article

6. Terry L, Sprinz E, Ribeiro JP. Moderate and high intensity exercise training in HIV-1 seropositive individuals: a randomized trial. *Int J Sports Med.* 1999; **20**(2): 142-6.10.1055/s-2007-971108
| PubMed | Web of Science ® | Google Scholar
7. Deresz LF, Sprinz E, Kramer AS, et al. Regulation of oxidative stress in response to acute aerobic and resistance exercise in HIV-infected subjects: a case-control study. *AIDS Care.* 2010; **22**(11): 1410-7.10.1080/09540121003758549
| PubMed | Web of Science ® | Google Scholar
8. Thomson RL, Brinkworth GD, Noakes M, Clifton PM, Norman RJ, Buckley JD. The effect of diet and exercise on markers of endothelial function in overweight and obese women with polycystic ovary syndrome. *Human Reprod.* 2012; **27**(7):2169-76.10.1093/humrep/des138
| PubMed | Web of Science ® | Google Scholar
9. Cotie LM, Josse AR, Phillips SM, MacDonald MJ. Endothelial function increases after a 16-week diet and exercise intervention in overweight and obese young women. *Biomed Res Int.* 2014; 2014: 327-95.
| Web of Science ® | Google Scholar
10. Januszek R, Mika P, Konik A, Petriczek T, Nowobilski R, Niżankowski R. The effect of treadmill training on endothelial function and walking abilities in patients with peripheral arterial disease. *J Cardiol.* 2014; **64**(2): 145-51.10.1016/j.jcc.2013.12.002
| PubMed | Web of Science ® | Google Scholar
11. Celermajer DS, Sorensen KE, Gooch VM, et al. Non-invasive detection of endothelial dysfunction in children and adolescents. *Lancet.* 1992; **340**(8828): 1111-1115.
| PubMed | Web of Science ® | Google Scholar

About Cookies On This Site

We and our partners use cookies to enhance your website experience, learn how our site is used, offer personalised features, measure the effectiveness of our services, and tailor content and ads to your interests while you navigate on the web or interact with us across devices. You can choose to accept all of these cookies or only essential cookies. To learn more or manage your preferences, click "Settings". For further information about the data we collect from you, please see our [Privacy Policy](#).

X ; 340(8828):
Accept All

Essential Only

Settings

Moderate intensity exercise training reverses functional aerobic impairment in HIV-infected individuals >

Gregory A. Hand et al.

AIDS Care

Published online: 29 Sep 2008

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright

Accepted

Registered
5 Howick Pl

In this article

About Cookies On This Site

We and our partners use cookies to enhance your website experience, learn how our site is used, offer personalised features, measure the effectiveness of our services, and tailor content and ads to your interests while you navigate on the web or interact with us across devices. You can choose to accept all of these cookies or only essential cookies. To learn more or manage your preferences, click "Settings". For further information about the data we collect from you, please see our [Privacy Policy](#).

Accept All

Essential Only

Settings

or & Francis Group
orma business