


[HIV Clinical Trials](#) >Volume 19, 2018 - [Issue 4](#) Free access679 | 3 | 0
Views | CrossRef citations to date | Altmetric[Listen](#)

Brief Report

Short-term folic acid supplementation and aerobic exercise improve vascular reactivity in HIV-infected individuals

Shana S. Grigoletti, Jorge P. Ribeiro, Eduardo Sprinz & Paula A. B. Ribeiro 

Pages 148-151 | Published online: 05 Feb 2018

 Cite this article <https://doi.org/10.1080/15284336.2018.1433769> Full Article Figures & data References Citations Metrics Reprints & Permissions View PDF Share

Abstract

The aim of this study was to determine the effect of supervised exercise and folic acid supplementation on endothelial function in HIV-infected individuals. A randomized clinical trial, double blinded, was conducted with 16 HIV-infected individuals, antiretroviral therapy (at least 6 months) with undetectable viral load (<50 copies/mL), and CD4 count > 200 cells/mm³. The subjects were randomized to aerobic exercise (n = 5) and daily intake for 4 weeks of 5 mg of folic acid (n = 6) or placebo (n = 5) groups. To assess endothelial function, venous occlusion plethysmography in the brachial artery by the protocol of reactive hyperemia was performed. The aerobic protocol consisted in cycling exercise, 3 times/week at 60–80% VO₂max, for 4 weeks. Exercise group (Δ6.5 mL/min/100 mL) and folic acid group (Δ7.3 mL/min/100 mL)

-0.3 ml/min/100 ml, time $p < 0.001$, interaction $p = 0.02$). Results demonstrate that supervised exercise and folic acid supplementation in very short term improve endothelial function in HIV-infected individuals. As exercise and folate supplementation are safe and relatively inexpensive, this finding deserves more attention in large randomized clinical trials in an attempt to reduce cardiovascular risk in HIV-infected population.

Keywords:

Exercise training

Endothelial function

HIV

Introduction

The antiretroviral therapy (ART) has dramatically reduced the morbidity and mortality associated to HIV.^{[1](#)} On the other hand, with longer life expectancy, metabolic chronic non-communicable diseases are more prevalent in patients under ART therapy when compared to naïve patients.^{[2,3](#)} In this scenario, cardiovascular disease (CVD) is one of the leading causes of morbidity and mortality in such population,^{[4-6](#)} and the possible mechanisms of raised incidence could be related to endothelial dysfunction.^{[7-12](#)} Therefore, we believe that it should be important to identify and manage individuals with endothelial dysfunction early on, which compromises up to 35% of these patients.^{[13,14](#)}

Our previous report has shown that supplementation with folates improves endothelium-dependent vasodilatation in HIV patients.^{[15](#)} However, non-pharmacologic safe strategies^{[16](#)} have been considered to reduce cardiovascular complications, such as exercise as an additional therapy, since the ARV treatment involves increased cardiovascular risks. Previously, acute responses to exercise demonstrated potential benefits.^{[17](#)} Also, chronic effects (exercise training) have been used to improve endothelial function in patients such as obese^{[18,19](#)} and peripheral arterial disease patients^{[20](#)} (Table 1).

Table 1 HIV patients' clinical characteristics



Nevertheless, there is still no study evaluating the effect of folic acid supplementation when compared to exercise intervention on endothelial function in HIV patients. This study aimed to determine whether 4-weeks of folic acid supplementation (5 mg/day) and aerobic exercise training would improve vascular reactivity, as assessed by brachial artery reactive hyperemia in HIV-infected people. For that, we conducted a pilot randomized, double-blind, placebo-controlled trial, with HIV-infected individuals, aged 18y or over, on ART for at least 6 months, with undetectable viral load (less than 50 copies/ml), and CD4 counts more than 200 cells/mm³. Exclusion criteria were diabetes mellitus, active infection, liver, kidney or cardiovascular diseases, pregnancy, and illicit drug use. All regular medications were maintained. The protocol was approved by the committee for ethics in research of the Hospital de Clínicas de Porto Alegre, according to Declaration of Helsinki, and all participants gave written informed consent.

Population and methods

Participants were randomly assigned to a 4-week treatment with either folic acid (n = 6) or placebo (n = 5) both blinded to the researcher and participants as described before,^{[15](#)} and a not blinded arm of exercise group (n = 5). The primary outcome measure was the assessment of brachial artery vascular responses during reactive hyperemia. Secondary outcomes were the changes on the biochemical and hemodynamic variables. All outcomes were blinded for the evaluators and statistician.

In short, reactive hyperemia (RH) was induced by placing a cuff in the upper-arm at 250 mm Hg, and releasing after 5 min, in dominant forearm. All flow recordings were manually traced by an operator who was blinded to the groups, and time. After 15 min of rest, 2.5 mg of sublingual isosorbide dinitrate (Isordil®/Sigma/Brazil) was administered as an endothelium-independent vasodilator. Five minutes later, endothelium-independent vasodilatation of the brachial artery was measured.^{[21](#)} Folic acid and placebo capsules were indistinguishable in shape, size, as well as color. They were pre-packed in identical bottles and consecutively numbered for each patient, according to the randomization schedule. Supplements were coded so that neither the investigators nor the participants were aware of the contents. Reported compliance was


100% of prescription in all groups. Missed supervised exercise sessions were replaced in the same week.

Results

Our results showed no significant change in hemodynamic and biochemical variables. Likewise, blood glucose, lipid profile, creatinine, and vitamin B₁₂ did not change significantly in the three groups after intervention (data not showed).

Figure 1 shows the delta of reactive hyperemia response and endothelium-dependent vasodilation responses (blue bars) after the 4 weeks. Exercise improves reactive hyperemia (9.4–15.9 mL/min/100 mL) and folic acid (11.9–19.2 mL/min/100 mL). The same was not observed in the placebo group (from 18.7 to 18.4 ml/min/100 ml, time $p < 0.001$, interaction $p = 0.02$). The endothelium-independent responses (green bars) remained unchanged in both groups (groups $p = 0.67$; interaction $p = 0.73$). No adverse effects were observed.

Figure 1 Deltas and SE of reactive hyperemia (FMD), endothelium-dependent and independent, after 4-weeks intervention in HIV patients.



Display full size

The main finding of this small randomized clinical trial was that short-term interventions of oral folic acid supplementation and aerobic exercise improved similarly endothelial-dependent vasodilation in HIV-infected people. To the best of our knowledge, this is the first study to investigate the effect of short-term intervention of aerobic exercise when compared to folic acid supplementation on endothelial function in this population. Our results extend these vascular benefits to HIV-infected people on stable antiretroviral therapy. More importantly, both interventions are low cost and could be easily implemented in the clinical site routine. However, non-supervised exercise effect (counseling and home-based strategies) still needs to be tested in these patients.

In conclusion, our preliminary results demonstrate that short-term administration methylated folate derivative from folic acid, and aerobic exercise training improve endothelium-dependent vasodilatation in HIV-infected individuals. This finding deserves more attention in larger randomized clinical trials, and future studies should also evaluate whether prolonged interventions would be effective to improve cardiovascular risks and major cardiovascular events in these population.

Contributors

SSG, ES, JPR, PABR conceived and designed the experiments. SSG was responsible for data collection. SSG, PABR analyzed data. SSG, PABR wrote the paper. The Final review was done by ES, PABR.

Funding

This work was supported by CAPES; CNPq, Brasília, Brazil; FIPE-HCPA, Porto Alegre, Brazil through scholarships.

Acknowledgments

In memoriam of Dr Jorge Pinto Ribeiro that was the first mentor of this study. This study

References

1. Sterne JA, Hernán MA, Ledergerber B, et al. Long-term effectiveness of potent antiretroviral therapy in preventing AIDS and death: a prospective cohort study. *Lancet*. 2005; 366(9483): 378–84.10.1016/S0140-6736(05)67022-5
[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
2. Trevisol F, Alencastro P, Ribeiro P, et al. Association of physical activity with lipodystrophy syndrome in HIV-infected patients. *J AIDS Clin Res*. 2012; 3(8): 1.
[Google Scholar](#)
3. Islam FM, Wu J, Jansson J, Wilson DP. Relative risk of cardiovascular disease among people living with HIV: a systematic review and meta-analysis. *HIV Med*. 2012; 13(8): 453–68.
[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
4. Grinspoon S, Carr A. Cardiovascular risk and body-fat abnormalities in HIV-infected adults. *N Engl J Med*. 2005; 352(1): 48–62.10.1056/NEJMra041811
[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
5. Friis-Moller N, Sabin CA, Weber R, et al. Combination antiretroviral therapy and the risk of myocardial infarction. *N Engl J Med*. 2003; 349(21): 1993–2003.
[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
6. Benjamin EJ, Blaha MJ, Chiuve SE, et al. Heart disease and stroke statistics-2017 update: a report from the American Heart Association. *Circulation*. 2017; 135(10): e146–603.10.1161/CIR.0000000000000485
[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

7. Blum A, Hadas V, Burke M, Yust I, Kessler A. Viral load of the human immunodeficiency virus could be an independent risk factor for endothelial dysfunction. *Clin Cardiol*. 2005; 28(3): 149-53.10.1002/clc.v28:3
[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
8. Solages A, Vita JA, Thornton DJ, et al. Endothelial function in HIV-infected persons. *Clin Infect Dis*. 2006; 42(9): 1325-32.10.1086/503261
[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
9. Stein JH, Klein MA, Bellehumeur JL, et al. Use of human immunodeficiency virus-1 protease inhibitors is associated with atherogenic lipoprotein changes and endothelial dysfunction. *Circulation*. 2001; 104(3): 257-62.10.1161/01.CIR.104.3.257
[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
10. Blanco JJ, Garcia IS, Cerezo JG, et al. Endothelial function in HIV-infected patients with low or mild cardiovascular risk. *J Antimicrob Chemother*. 2006; 58(1): 133-9.10.1093/jac/dkl190
[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
11. de Larrañaga GF, Petroni A, Deluchi G, Alonso BS, Benetucci JA. Viral load and disease progression as responsible for endothelial activation and/or injury in human immunodeficiency virus-1-infected patients. *Blood Coagul Fibrinolysis*. 2003; 14(1): 15-18.10.1097/00001721-200301000-00004
[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
12. Monsuez JJ, Dufaux J, Vittecoq D, Vicaut E. Reduced reactive hyperemia in HIV-infected patients. *J Acquir Immune Defic Syndr*. 2000; 25(5): 434-42.10.1097/00126334-200012150-00009
[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
13. Balsam P, Mikuła T, Peller M, et al. Evaluation of endothelial function and arterial stiffness in HIV-infected patients: a pilot study. *Kardiol Pol*. 2015; 73(5): 344-

4. Currier JS, Lundgren JD, Carr A, et al. Epidemiological evidence for cardiovascular disease in HIV-infected patients and relationship to highly active antiretroviral therapy. *Circulation*. 2008; 118(2): e29–35.10.1161/CIRCULATIONAHA.107.189624

5. Grigoletti SS, Guindani G, Moraes RS, Ribeiro JP, Sprinz E. Short-term folinic acid supplementation improves vascular reactivity in HIV-infected individuals: a randomized trial. *Nutrition*. 2013;29(6):886–891. doi:10.1016/j.nut.2013.01.015.

6. Terry L, Sprinz E, Ribeiro JP. Moderate and high intensity exercise training in HIV-1 seropositive individuals: a randomized trial. *Int J Sports Med*. 1999; 20(2): 142–6.10.1055/s-2007-971108

7. Deresz LF, Sprinz E, Kramer AS, et al. Regulation of oxidative stress in response to acute aerobic and resistance exercise in HIV-infected subjects: a case-control study. *AIDS Care*. 2010; 22(11): 1410–7.10.1080/09540121003758549

8. Thomson RL, Brinkworth GD, Noakes M, Clifton PM, Norman RJ, Buckley JD. The effect of diet and exercise on markers of endothelial function in overweight and obese women with polycystic ovary syndrome. *Human Reprod*. 2012; 27(7):2169–76.10.1093/humrep/des138

9. Cotie LM, Josse AR, Phillips SM, MacDonald MJ. Endothelial function increases after a 16-week diet and exercise intervention in overweight and obese young women. *Biomed Res Int*. 2014; 2014: 327–95.

20. Januszek R, Mika P, Konik A, Petriczek T, Nowobilski R, Niżankowski R. The effect of treadmill training on endothelial function and walking abilities in patients with peripheral arterial disease. *J Cardiol*. 2014; 64(2): 145–51.10.1016/j.jjcc.2013.12.002

[PubMed](#)

[Web of Science®](#)

[Google Scholar](#)

21. Celermajer DS, Sorensen KE, Gooch VM, et al. Non-invasive detection of endothelial dysfunction in children and adults at risk of atherosclerosis. *Lancet*. 1992; 340(8828): 1111–5.10.1016/0140-6736(92)93147-F

[PubMed](#)

[Web of Science®](#)

[Google Scholar](#)

[Download PDF](#)

Related research

People also read

Recommended articles

Cited by
3

[Moderate intensity exercise training reverses functional aerobic impairment in HIV-infected individuals >](#)

Gregory A. Hand et al.

AIDS Care

Published online: 29 Sep 2008

[Acute high-intensity interval exercise versus moderate-intensity continuous exercise in heated water-based on hemodynamic, cardiac autonomic, and vascular respo... >](#)

Isabela Roque Marçal et al.

Clinical and Experimental Hypertension

Published online: 19 Apr 2022

Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG

 Taylor and Francis
Group

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)