



International Journal of Fruit Science >

Volume 20, 2020 - [Issue sup2](#)

✓ Free access

4,622 Views | 8 CrossRef citations to date | 0 Altmetric

[Listen](#)

Research Article

Gibberellic Acid and Potassium Nitrate Promote Seed Germination and Growth of Grey-leaved Saucer-berry (*Cordia Sinensis* Lam.) Seedlings

Rahul Dev , Devi Dayal & M. Sureshkumar

Pages S937-S954 | Published online: 11 Jun 2020

Cite this article <https://doi.org/10.1080/15538362.2020.1774465>



Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

View PDF

View EPUB

Share

Formulae display: **MathJax**

ABSTRACT

Cordia sinensis

Lam is a multipurpose underutilized species of arid and semiarid regions. It is commonly propagated through seeds, but germination percentage is low (50–60%) and erratic. Therefore, an investigation was undertaken to standardize the propagation technique during year 2016 at CAZRI RRS Bhuj, India. The seeds of *Cordia gharaf* were treated with gibberellic acid (four levels: 0, 250, 500, and 1000 ppm) and potassium nitrate (four levels: 0%, 0.25%, 0.5%, and 1%) singly or in combination and it was found that combined application of GA₃ and KNO₃ enhanced seed germination

resulted in earliest seed germination (4.33 ± 0.33 days), maximum seed germination percentage (79.47%), and subsequent seedling growth e.g. highest leaf area (25.52 ± 1.41 cm²), and maximum leaf fresh (1.59 g) and dry weight (0.72 g). However, maximum plant height was observed under GA₃ 500 ppm and KNO₃ (1%) treatment. In conclusion, the combination treatment of GA₃ 250 ppm and KNO₃ 1% is an effective and practical method for improving *C. sinensis* seed germination.

KEYWORDS:

Cordia sinensis

gibberellic acid

potassium nitrate

seed germination

Introduction

Cordia sinensis Lam. (*Cordia gharaf* (Forssk.) Ehrenb. ex Asch.) is a multi-stemmed tree, growing up to 8 to 12 m height. *Cordia sinensis* is a potential underutilized plant which is also known as Gondi, narrow-leaved sebastian or Greyleaved saucer-berry and belongs to the family Boraginaceae. The genus *Cordia* consists of about 250 species and the majority of species are tree or shrub (Barroso and Oliveira, [2009](#)). It is well-adapted plant species in arid to semiarid regions of West Africa to East to the Middle East, India, Sri Lanka, and Pakistan due to its various adaptive features e.g. strong tap root system, waxy leaves, hairiness, sunken and covered stomata in leaves, water binding mechanism, moderate tolerance to salinity and alkalinity. These traits make it hardier than *Cordia myxa*, a popular fruit plant of the Boraginaceae family (Meghwal et al., [2014](#)). The fruits of the *C. sinensis* Lam. are consumed by local inhabitants. The leaves provide the protein (13.12%) rich fodder to cattle (Kuria et al., [2005](#)). Various parts of the plant have ethnomedicinal value and are traditionally used to cure different human and livestock ailments (Gumgumjee and Hajar, [2015](#)). Farmers of arid regions grow *C. sinensis* around the farm boundary to protect the main crop from hot and dry wind (Meghwal et al., [2014](#)). Therefore, *C. sinensis* may be a potential species for domestication and utilization in the arid region. The total plant population of *C. sinensis* was about 864 thousand which share only 0.32% of the total plant population of the Gujarat state (Pradeep and Singh, [2010](#)). In Kachchh region of Gujarat, *C. sinensis* was distributed among the 29.09% of surveyed sites (Dey et al., [2016](#)). While it is reported

(Singh, [2018](#)). Generally, *Cordia sinensis* is propagated through the seeds but germination percentage is poor (50–60%) and erratic (Meghwal, [2007](#)). Various germination studies conducted in the other *Cordia* species viz. *Cordia boissieri* reveals conflicting information with regard to breaking seed dormancy through stratification (Schuch et al., [2001](#)). The study conducted in *Cordia myxa* and *C. gharaf* indicated seed treatment with gibberellic acid (GA_3) (250, 500 mg L⁻¹) promotes germination significantly (Meghwal, [2006](#)). Similarly, about 50–60% seed germination could be obtained with fresh seeds of *C. sinensis*, but the germination percentage is low and not uniform (Maundu et al., [2005](#); Meghwal et al., [2014](#)). However, another study in *Cordia biosseri* showed high percentage of seed dormancy in the newly produced seeds (Fulbright, [1992](#); Schuch et al., [2001](#)). Both GA_3 and KNO_3 stimulate the seed germination of dormant seeds of various plant species (Koyuncu [2005](#); Cárdenas et al., [2013](#); Mello et al., [2009](#)). However, the studies carried out by Ding et al. ([2007](#)), Gao et al. ([2011](#)), Cárdenas et al. ([2013](#)) observed the effect of GA_3 and KNO_3 on seed germination of other species but very few on *C. sinensis* seed germination.

The proper seedling growth of germinated seeds is the most important factor for successful sapling production under nursery conditions. The earlier studies showed the variable effect of GA_3 and KNO_3 on growth traits of various plants. Application of GA_3 and KNO_3 promotes overall plant growth in *Brassica oleracea Capitata* (Majumdar, [2013](#)), *Cucumis sativus* (Pal et al., [2016](#)), and *Solanum lycopersicum* (Balaguera-Lopez et al., [2009](#)). GA_3 (250 mg L⁻¹) treatment resulted in maximum seed germination (98.75%), subsequent shoot length, root length, leaf area, shoot, and root dry weight (Shabaq, [2013](#)). Similar growth promotion of seedlings was also reported under the GA_3 treated seeds of *Citrus limon* (Dzayi and Rahman, [2010](#)), *Nothapodytes nimmoniana* (Patil et al., [2015](#)), and *Citrus aurantifolia* (Jaiswal et al., [2018](#)). Lay et al. ([2015](#)), found GA_3 and KNO_3 presowing treatment to seeds of papaya improves the seedling growth, fresh and dry weight of seedlings. However, very few studies are available with quantitative data on the subsequent growth of treated plants. The *C. sinensis* is very important species of arid and semiarid regions due to its multiple uses for fodder, fruits, and medicines, yet the multiplication of species through seeds provides poor and erratic germination. Therefore, present investigation attempted to improve the seed germination rate and quantify the subsequent growth for faster multiplication for the *Cordia sinensis*, an important species of the arid region.

Material and Methods

The study was conducted at Horticulture Nursery, ICAR-Central Arid Zone Research Institute, Regional Research Station, Bhuj Gujarat, India. The seed germination experiment was performed in the laboratory, while, subsequent plant growth study was carried out under the shade net. *Cordia sinensis* fruits were collected from the native population in the Kachchh region of Gujarat (22°84'624 – 23°56'288 N° latitude and 68°95'730 – 70°73'100 E° longitude). Fresh seeds were extracted from the ripened fruits. The mucilaginous pulp was removed from the seeds by rubbing the seeds with sandy soil followed by washing with tap water. Air dried, cleaned seeds were stored in the paper bag for 2 weekss at the room temperature before use.

Experiment 1. Chemical Treatment of Seeds

Before the seed treatment, seeds were disinfected with fungicide (Carbendazim 2%) for 20–30 minutes and rinsed three times with running tap water. For the treatments, seeds were divided into four groups, the total no. of treatment combinations were 16, four levels of each of GA₃ concentration (0, 250, 500, and 1000 ppm) and four KNO₃ concentrations (0%, 0.25%, 0.5%, and 1% KNO₃) purchased from Sigma Aldrich (90% gibberellin A₃ basis) Loba Chemie (99% purity), respectively and were evaluated with 3 replications 1) seeds of first group were transferred into the beaker filled with distilled water and left for 24 h; 2) seeds of second group were soaked in 250, 500, and 1000 mg L⁻¹ (w/v) GA₃ solution for 24 h; 3) similarly, seeds of the third group were soaked in 0.25%, 0.5%, and 1% KNO₃ solution for 24 h; 4) the seeds of the fourth group were treated with the aqueous solution supplemented different combinations of GA₃ and KNO₃ ([Table 1](#)).

Table 1. Test design of chemicals (KNO₃, GA₃) and combinations of KNO₃ and GA₃ treatments



[Download CSV](#) [Display Table](#)

The previously treated seeds of *Cordia sinensis* were taken for germination test. The 16 treatments were arranged in completely randomized design with three replications. The average temperature of laboratory during experiment was about 35°C. A layer of cotton

size of blotting paper. The 20 seeds were wrapped in blotting paper and cotton wool placed in the Petri dishes followed by soaking with distilled water (each treatment with three Petri dishes, e.g., 60 seeds per treatment). The seeds were kept moist uniformly by frequent spraying of water. The initiation of germination was recorded at daily basis and number of germinated seeds on particular day was also noted and percentage of germination was recorded at the weekly interval by unfolding the blotting paper up to 30 days of sowing after which there was no further germination noticed. The germination percentage was calculated as follows (Gashi et al., [2012](#))

$$G\% = \frac{a}{N} \times 100$$

Where G = germination percentage, a = the total no. of germinated seeds till 30 days of sowing and N = the total no. of seeds sown.

While, mean germination time was estimated using the following method (Moradi et al., [2008](#)).

$$MGT = \frac{\sum Dn}{\sum n}$$

Where n = number of seeds which were germinated on day D,

D = number of days taken for seed germination.

Experiment II. Seedling Growth at Net House

After the 30-day, the germinated plantlets of *C. sinensis* were shifted from laboratory to green shade net (50% shade and 1.5 mm thickness) conditions. The plantlets were established in the black-colored nursery poly bags (20 × 10 cm) and filled with sterilized soil (sand: soil: farmyardd manner – 2:2:1) potting mixture. The average temperature and relative humidity under shade net house during experiment were about 30°C and 74%. Different plant characteristics like plant height (cm), stem thickness (mm), no. of leaves per plant were measured at 30, 90, 180, 270, and 360 days of seed germination. Leaf area (cm²), leaf fresh weight (g), leaf dry weight (g), leaf nitrogen (%), phosphorous (%), potassium (%), sodium (%) were estimated at the end of the experiment. The stem thickness and leaf area (cm²) were measured using a digital vernier calliper and the leaf area meter (Licor-3100). For recording leaf dry weight, leaf samples were dried in the oven at 72°C for 48 h or until the constant weight was achieved. The leaf nutrients (e.g. (available N, P, K, and Na) were

solution, followed by single and double distilled water. The cleaned leaf samples were dried in a hot air oven at 72°C for 48 h or until they obtained a constant weight and ground and sieved. The concentration of K and Na was determined using the flame photometer (Model: ESICO1382). While, measurement of leaf P and N content was done through colorimetric and Kjeldahl method, respectively.

Statistical Analysis

The randomized complete block design (RCBD) was used for conducting the experiment in laboratory and shade net. The critical differences (CD) at $P < .05$ and $P < .01$ level of probability were determined after analysis of variance (ANOVA) for all treatments. The experiment was laid out with 16 treatments including control with three replications and 60 seeds per replicate were used. The mean and analysis of variance were performed to determine significant differences among treatments ($P < .05$) using the Statistical Software Package for Agricultural Research Workers (Sheoran et al., [1998](#)).

Result and Discussion

Experiment 1. Chemical Treatment of Seeds

Mean Germination Time (MGT) and Germination Percentage (GP)

The different concentrations of GA₃ and KNO₃ significantly fasten the seed germination as compared to the germination percentage of control seeds. The KNO₃ (0.5%) treatment significantly shortens the germination initiation duration (4 days), compared to other single KNO₃ applications. Whereas the synergistic effect of GA₃ × KNO₃ treatments was observed on the shortening of germination time. The seeds treated with aqueous solution supplemented with GA₃ 250 mg L⁻¹ and 1% KNO₃, took minimum time for germination initiation (4.33 ± 0.33 days), which was about 1 week less than germination initiation in control treatment. Among the treatment, maximum time for germination initiation was taken by the seeds soaked in the KNO₃ 0.25% and 1.0% concentration ([Table 2](#)).

[Table 2](#). Analysis of variance of gibberellic acid and potassium nitrate effects on seed germination properties for Cordia sinensis.



The result of the $\text{GA}_3 \times \text{KNO}_3$ experiment revealed significant enhancement of final germination percentage ([Table 3](#)). Based on the results for germination percentage, the highly significant difference was noted between seeds treated with a different gibberellic acid concentration (250, 500, and 1000 mg L^{-1} GA_3) and with a combination of the different concentration of potassium nitrate (0.25%, 0.5%, and 1%) and untreated seeds (control). The maximum final germination percentage was noted in the seeds treated with 250 mg L^{-1} $\text{GA}_3 \times 1\%$ KNO_3 (79.47%) and 500 mg L^{-1} $\text{GA}_3 \times 0.25\%$ KNO_3 (67.17%), as compared to the germination percentage of control (21.97%). In contrast, when seeds treated with a higher GA_3 concentration along with higher KNO_3 treatment, the final germination percentage reduced drastically. Analysis of data revealed that both single factors (GA_3 and KNO_3 concentration) affected seed germination percentages ($p \leq 0.05$). However, higher germination percentage was reported under the 0.5% KNO_3 treatment, which was higher than all other single treatments. However, GA_3 treatment singly did not significantly increase the final seed germination percentage. Similar study conducted in *Cordia biosseri* reveals a high percentage of dormant seeds and GA_3 treatment, breaks the seed dormancy (Fulbright, [1992](#); Schuch et al., [2001](#)). Similarly, Meghwal (2007) noted 20–30% seed germination in *Cordia myxa* and up to 60% seed germination in *Cordia gharaf* with freshly extracted seeds and seed treatment with GA_3 and mechanical scarification significantly improve (66.66%) germination of species. On the contrary, Maundu et al. ([2005](#)) observed that, seed treatment of fresh seeds of *Cordia sinensis* generally not necessary but, stored seeds need to be soaked with warm water (40°C) until water is cool. However, Acharya and Purohit ([1999](#)) found that GA_3 promoted germination of seeds in *Cordia gharaf* to the maximum extent of 81%. Waman et al. ([2017](#)) recommended that presowing treatments with KNO_3 and GA_3 were effective to produce the healthy seedlings of *Semecarpus kurzii*. In general, seeds treated with KNO_3 (0.5%) had a higher overall percentage seedling emergence than GA_3 treatments. The gibberellic acid promotes the germination of dormant seeds of many species such as, *Citrullus lanatus* (Ding et al., [2007](#)), *Penstemon digitalis* (Mello et al., [2009](#)), *Tamarindus indica* (Patel et al., [2018](#)) and *Elaeocarpus prunifolius* (Iralu and Upadhaya, [2018](#)). The KNO_3 solution increases the germination percentage of *Lilium orientalis* seeds (Gao et al., [2011](#)), *Eremurus spectabilis* seeds (Keskiner and Tuncer, [2019](#)), and *Passiflora edulis* f. *Flavicarpa*

Cárdenas et al. (2013) who reported that increased seed germination was directly proportional to the increase in GA₃ concentrations (500 and 1000 mg L⁻¹). However, at higher GA₃ concentration (1500 mg L⁻¹), germination of *Passiflora* species decreased (Cárdenas et al., 2013). On the contrary, lower concentration (80–250 mg L⁻¹) promotes the seed germination of *Eriobotrya japonica* (Al-Hawezy, 2013), *Nothapodytes nimmoniana* (Patil et al., 2015) and *Citrus aurantifolia* (Jaiswal et al., 2018). Seed germination was improved due to gibberellic acid (GA) treatment possibly through increased synthesis of hydrolytic enzymes, which were further transported to endosperm. The enzymes break down the stored food to supply the energy required for germination (Chen et al., 2008; Varner, 1965). GA₃ may promote cytokinin action and counteracts the germination inhibitor (ABA) which ultimately stimulate the biochemical processes of seed germination (Chen et al., 2008; Cetinbas and Koyuncu, 2006; Tuan et al., 2018). Potassium nitrate improves the seed germination percentage in the various plants species (Duermeyer et al., 2018; Keskiner and Tuncer, 2019). A study on the germination of mountain ash seeds showed that germination of the species was affected by both the concentration of KNO₃ and the time of KNO₃ exposure (Bian et al., 2013). Similarly, Gupta et al. (2011) reported that treatment of *Hippophae salicifolia* seeds with 0.1% KNO₃ pretreatment resulted in 96% seed germination. However, a higher concentration of KNO₃ negatively affects seed germination. A similar trend of reduction of seed germination on the application of higher KNO₃ concentration was reported in *Salvia cyanescens* (Yücel and Yilmaz, 2009) and *Gladiolus alatus* (Ramzan et al., 2010).

Table 3. Mean of germination, growth (MGT, GP, LA, LFW, LDW) and seedling nutrients (NPK and Na) of *Cordia sinensis* under various levels of GA₃ (in ppm) and KNO₃ (in %)



Download CSV

Display Table

According to Gniazdowska et al. (2007), KNO₃ possibly stimulates seed germination by the production of nitrous oxide. Nitrous oxide may increase seed germination and break seed dormancy by regulation of endogenous balance of abscisic acid and gibberellic acid. It may influence the expression of enzymes that triggers a nitrate-induced ABA decrease and the biosynthesis of germination promoters (gibberellins) (Duermeyer et al., 2018; Finch et al., 2007; Sirova et al., 2011). The present study shows a synergistic


an aqueous solution of gibberellins supplemented with potassium nitrate solution. Similar study conducted by Meghwal (2007) indicates that merely water imbibition (20–30% germination) was not sufficient to trigger the germination in *Cordia myxa* but seed treatment with gibberellic acid (GA_3) (500 mg L^{-1}) was essential to promote seed germination (66.66%). However, KNO_3 did not influence the seed germination, irrespective of treatment duration and chemical concentration. Gashi et al. (2012) found the higher germination percentage (92.26%) in the *Ramonda* species' seeds treated with 1000 ppm GA_3 + 0.3% KNO_3 . Treatment of *Sorghum bicolor* seeds with GA_3 (100 ppm) and KNO_3 (1%) resulted in increase of germination percentage to a higher extent (94%) (Shanmugavalli et al., 2007). Similar synergistic results of KNO_3 (1%) and GA_3 (500 ppm) seed treatments were reported by Dewir et al. (2011) in *Sabal palmetto*. Whereas, soaking cracked seeds of *Elaeocarpus prunifolius* in the solution of GA_3 and KNO_3 accelerate the germination rate (Iralu and Upadhaya, 2018).

Experiment 2. Subsequent Quantification of Growth

Plant Height and Collar Diameter

Application of $\text{GA}_3 \times \text{KNO}_3$ singly or in combination, significantly ($p = .05$) affected the plant height (Figure 1). However, plant height under different single KNO_3 applications differed significantly. Lower KNO_3 application (0.25% conc.) tends to increase plant height at par to GA_3 500 mg L^{-1} , but at higher KNO_3 application (1%), there was a reduction in the plant height. The longest plants were observed from the GA_3 (500 mg L^{-1}) and KNO_3 (1%) applications, followed by combined application of 500 mg L^{-1} GA_3 and 0.5% KNO_3 . Whereas, the minimum increase in the plant height was seen when plants seeds soaked in the GA_3 1000 mg L^{-1} and KNO_3 1% solutions for 24 h.


Figure 1. Plant height of *Cordia sinensis*, under various levels of GA_3 (in ppm) and KNO_3 (in %) seed treatment, recorded at three month interval



Display full size

Data presented in [Figure 2](#) illustrated the effect of GA₃ and KNO₃ on the collar diameter of *C. sinensis*. The significant difference was ascertained for *C. sinensis* between seeds treated with GA₃, KNO₃ and H₂O (control). All GA₃ and KNO₃ applications had a positive effect on collar diameter. However, single GA₃ applications were more effective than different single KNO₃ application. Moreover, integrated application of GA₃ and KNO₃ resulted pronounced improvement in collar diameter of plantlets. The maximum increase in the collar diameter in one-year-old plants was observed with the GA₃ 500 mg L⁻¹ and KNO₃ 0.5% treatment followed by GA₃ 1000 mg L⁻¹ and KNO₃ 0.25% treatment. On the contrary, the minimum collar diameter was recorded in the plants of control group. The presowing treatment with lower concentration of GA₃ and KNO₃ improved the seedling height significantly however, reduction in the plant height was observed under the higher concentration of GA₃ and KNO₃. Treatment of *Eriobotrya japonica* seeds with lower concentration of GA₃ (250 mg L⁻¹) resulted maximum collar diameter and plant height comparing to control (Shabaq, [2013](#)). Similarly, maximum plant height and stem diameter of *Citrus aurantifolia* plantlets were observed with GA₃ (200 mg L⁻¹) seed treatment (Meshram et al., [2015](#)).

Figure 2. Collar diameter of *Cordia sinensis*, under various levels of GA₃ (in ppm) and KNO₃ (in %) seed treatment, recorded at three month interval



Display full size

The GA mediated increment in plant height and collar diameter chiefly due to enhancement of cell division in cambium tissues and its prompt cell progeny as supported by the Parab et al. (2017) in *Carica papaya* and Harsha et al. (2017) in *Citrus maxima*.


Leaf Area and Leaves Number per Plant

The single factors (GA₃ and KNO₃ concentration) significantly influenced the leaf area ($p \leq 0.05$) (Table 3). Seeds treated with the different concentrations of GA₃ are found to have more leaf area as compare to KNO₃ treated seed and untreated seeds (control). A lower concentration of GA₃ (250 mg L⁻¹) had a favorable effect on leaf expansion. While at a higher concentration of GA₃ treatment leaf area found to be non-significantly decreased. Similar to GA₃, leaf area of treated KNO₃ seeds was declined with increases of KNO₃ concentration. The GA₃ and KNO₃ treatments significantly affected the leaf area. The maximum leaf area was recorded in the plants treated with GA₃ 250 × KNO₃ 1%, followed by GA₃ 500 × KNO₃ 0.5%. While, treatment combination, GA₃ 1000 mg L⁻¹ × KNO₃ 0.25% had some adverse effect on leaf expansion, and it resulted in minimum leaf area. Similar, leaf expansion at lower GA₃ concentration (250 mg L⁻¹) was observed in *Eriobotrya japonica* (Shabaq, 2013), *Citrus limon* (Dzayi and Rahman, 2010) and *Citrus aurantifolia* (Meshram et al., 2015).

In general, numbers of leaves per plant were not improved by the pretreatment of GA₃ and KNO₃ (Figure 3). However, numbers of leaves per plant were significantly improved with combined application of GA₃ and KNO₃. The presowing treatment of seeds with

followed by seeds treated with GA_3 250 $\text{mg L}^{-1} \times \text{KNO}_3$ 0.5%. In contrast, GA_3 seeds soaked in higher gibberellin concentration (1000 mg L^{-1}) supplemented with medium or high KNO_3 (0.5% or 1%) concentration for 24 h, gave the minimum increase in the leaf numbers per plant. The various previous study reported the opposite relations between leaf growth and plant height increment due to GA_3 treatment (Arney and Mancinelli, 1966). These trends are concurrent with the present study. GA_3 pregermination treatments increase the leaf area and leaf numbers in *Carica papaya* (Parab et al., 2017) and *Citrus maxima* (Harsha et al., 2017). This is possibly due to the stimulation of cell division and enlargement by GA application (Parab et al., 2017).

Figure 3. Leaf number per plant of *Cordia sinensis*, under various levels of GA_3 (in ppm) and KNO_3 (in %) seed treatment, recorded at three month interval



Display full size

Leaf Fresh and Dry Weight

The result indicated that the highest leaf fresh and dry weight was achieved when the *C. sinensis* seeds were treated with the aqueous solution of GA_3 250 mg L^{-1} , supplemented with the KNO_3 1%, followed by the solution containing GA_3 500 mg L^{-1} and KNO_3 0.25%. The minimum leaf fresh weight was noted under the highest KNO_3 (1% conc.) treatment. In general, fresh weight of leaf was higher when seeds were treated with GA_3 than a KNO_3 solution; further fresh weight was decreased when GA_3 concentration increased from 0.25% to 1%. While the lowest dry weight was found

concentration of both the chemicals (GA_3 and KNO_3) resulted in the decline in the dry weight of leaves. Cárdenas et al. (2013) found the higher dry weight, total plant weight, specific leaf area and leaf area of KNO_3 treated passiflora seedling. Similarly, higher fresh and dry shoot weight of GA_3 treated seeds were found in the seedlings of *Carica papaya* (Meena and Jain, 2012 & Parab et al., 2017) and *Citrus maxima* (Harsha et al., 2017). Maximum vegetative dry weight and root weight were observed with GA_3 (250 mg L^{-1}) treatment in *Eriobotrya japonica* (Shabaq, 2013) and similarly, higher shoot and root dry weight were found due to GA_3 treatment in *Citrus limon* (Dzayi and Rahman, 2010) and *Citrus aurantifolia* (Jaiswal et al., 2018). The higher fresh and dry shoot weight in the GA_3 presowing treatment chiefly because of increased mobilization of water, membrane permeability, nutrient uptake, and transportation which ultimately leads to higher production of photosynthetic products (Parab et al., 2017; Harsha et al., 2017). While, higher mobilization of photosynthetic products was also taken place in the KNO_3 treated seeds in the early stage of seed germination (Lay et al., 2015).

Leaf Nutrient Traits

The different concentration of $\text{GA}_3 \times \text{KNO}_3$ affects the leaf N, P, K, and Na content significantly (Table 3). The leaf N content was lower under the different single GA_3 and KNO_3 treatments as compared to control. However, combination treatment of GA_3 1000 mg L^{-1} and KNO_3 0.25% resulted in highest leaf N content. In general leaf N content was higher in the plants treated with GA_3 as compare to KNO_3 treatments. The leaf P content was higher under the GA_3 treated plants it was highest under GA_3 250 mg L^{-1} treatment closely followed by GA_3 $500 \text{ mg L}^{-1} \times \text{KNO}_3$ 1% treatment. The combined application of GA_3 $1000 \text{ mg L}^{-1} \times 0.5\% \text{ KNO}_3$ resulted minimum leaf P content. In general higher leaf P content was achieved when seeds were treated with GA_3 for 24 h than KNO_3 treatment for the same duration. Percentage leaf K content was lower for GA_3 and KNO_3 treated seeds as compared to seeds treated with distilled water (control). However, among the treatments, the highest K content was noted when seeds were soaked in GA_3 500 mg L^{-1} followed by KNO_3 (0.25%). The lowest leaf K content was reported when the seeds were soaked in the aqueous solution of GA_3 1000 mg L^{-1} supplemented with KNO_3 1% concentration. The treatment of GA_3 resulted significantly high leaf Na content than control. However, there was no significant difference between seeds soaked in H_2O (control) and KNO_3 solution for leaf Na content. The seeds soaked in GA_3 (500 mg L^{-1}) and KNO_3 0.25% had the highest leaf

with an aqueous solution of GA₃ 1000 mg L⁻¹ supplemented with the KNO₃ 0.25% for 24 h.

Increase in leaf nitrogen and phosphorous content with foliar application of GA₃ and KNO₃ was reported in Cucumis sativus plants (Pal et al., [2016](#)). While the higher leaf nitrogen and potassium contents were significantly influenced by the different GA₃ application. Maximum leaf nitrogen and potassium were reported under combination of GA₃ (100 ppm) and urea (1%) in the Citrus jambhiri (Kant et al., [2017](#)). Kazemi ([2014](#)) noted that GA₃ application did not significantly affect leaf NPK content, but the application of potassium nitrate influenced leaf NPK content in Solanum lycopersicum leaves. The highest increase was reported with 6 mM KNO₃ application. A similar effect was observed by Zhang et al. ([2002](#)) and Lin and Danfeng ([2003](#)). Application of potassium nitrate might increase the plant metabolism and related functions thereby were responsible for increase in leaf NPK contents (Marschner, [2012](#)). Leaf sodium content was found to be decreased with combined application of gibberellins and potassium nitrate in the study. The similar antagonistic correlation was also observed by Song and Fujiyama ([1996](#)).

Thus, the external application of KNO₃ and GA₃ showed a synergistic effect in promoting seed germination and seedling vigor of Cordia sinensis seed in the present investigation. In conclusion, it can be stated that the application of GA₃ (250 ppm) and KNO₃ (1%) is a simple, effective, and practical method for improving seed germination and seedling growth of Cordia sinensis. Although, further work might be useful for in-depth understanding on effects of chemicals in Cordia sinensis.

Acknowledgments

Author is thankful to Director, ICAR-CAZRI for proving necessary facility for conducting research.

Disclosure Statement

The authors declare that they have no conflict of interest.

Literature Cited

1. Acharya, J., and S.S. Purohit. 1999. Effect of quality of light and growth regulators on seed germination of *Cordia gharaf* and *Cordia dichotoma*. In: Paper presented in national seminar on plant physiol. At interface of Agri Horticulture and Industary, (Vol. 66, 150.), Udaipur, India.

[Google Scholar](#)

2. Al-Hawezy, S.M.N. 2013. The role of the different concentrations of GA3 on Seed Germination and Seedling Growth of Loquat (*Eriobotrya japonica* L.). IOSR Journal of Agriculture and Veterinary Science 4 (5): 3-6.

[Google Scholar](#)

3. Arney, S.E., and P. Mancinelli. 1966. The basic action of gibberellic acid in elongation of 'Meteor' pea stems. *New Phytol.* 65:161-175.

[Web of Science ®](#) | [Google Scholar](#)

4. Balaguera-Lopez, H.E., J.F. Cardenas-Hernandez, and J.G. Alvarez-Herrera. 2009. Effect of gibberellic acid (GA3) on seed germination and growth of tomato (*Solanum lycopersicum* L.). *Acta Hort. (ISHS)*. 821:141-148. doi:10.17660/ActaHortic.2009.821.15.

[Google Scholar](#)

5. Barroso, I.C.E., and F.D. Oliveira. 2009. Pharmacognostic diagnosis of fruits of *Cordia sellowiana* Cham. and *Cordia myxa* L. (Boraginaceae). *Rev. Bras. Farmacogn.* 19:458-470. doi:10.1590/S0102-695X2009000300021.

[Google Scholar](#)

6. Bian, L., L. Yang, J. Wang, and H. Shen. 2013. Effects of KNO3 pretreatment and temperature on seed germination of *Sorbus pohuashanensis*. *J. Forest Res.* 24:309-316. doi:10.1007/s11676-013-0354-9.

7. Çetinbaş, M. and Koyuncu, F. 2006. Improving germination of *Prunus avium* L. seeds by gibberellic acid, potassium nitrate and thiourea. *Horticultural Science - Prague* 33(3): 119–123.

[Google Scholar](#)

8. Cárdenas, J., C. Carranza, D. Miranda, and S. Magnitskiy. 2013. Effect of GA₃, KNO₃, and removing of basal Point of seeds on germination of sweet granadilla (*Passiflora ligularis* Juss) and yellow Passion fruit (*Passiflora edulis* f. *Flavicarpa*). *Rev. Bras. Frutic.* 35(3):853–859. doi: 10.1590/S0100-29452013000300023.

[Web of Science ®](#) | [Google Scholar](#)

9. Chen, S.Y., S.R. Kuo, and C.T. Chien. 2008. Roles of gibberellins and abscisic acid in dormancy and germination of red bayberry (*Myrica rubra*) seeds. *Tree Physiol.* 28(9):1431–1439. doi: 10.1093/treephys/28.9.1431.

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

10. Dewir, Y.H. Mahrouk, M.E. and Naido, Y. 2011. Effect of some mechanical and chemical treatments on seed germination of *Sabal palmetto* and *Thrinax morrisii* palms. *Australian Journal of Crop Science* 5(3): 248-253.

[Google Scholar](#)

11. Dev, R., M.S. Kumar, S. Kumar, and D. Dayal. 2016. Survey, collection and utilization of *Grewia* and *Cordia* species for human and animal nutrition in Arid Kachchh, Gujarat, p.344. In IAC abstracts book. 1st International agrobiodiversity congress. November 6–9, 2016. New Delhi, India.

[Google Scholar](#)

12. Ding, Q.L., X.M. Dang, and Y.F. Zhan. 2007. Effect of potassium nitrate and gibberellin solution soaking on mini-watermelon seed germination. *J. South China Uni. Trop. Agric.* 13:14–16. in Chinese with English abstract.

[Google Scholar](#)

3. Duermeyer, L., E. Khodapanahi, D. Yan, A. Krapp, S. Rothstein, and E. Nambara. 2018. Regulation of seed dormancy and germination by nitrate. *Seed Sci. Res.* 28(3):150–157. doi: 10.1017/S096025851800020X.

[Web of Science ®](#) | [Google Scholar](#)

4. Dzayi, F.H., and Rahman. 2010. Effect of GA3 and Soaking time on Seed Germination and Seedling Growth Lemon (*Citrus limon* L.) High Diploma.Thesis coll.Agric, univ. of Salahaddin – Erbil.

[Google Scholar](#)

5. Finch, W.E., C.S. Cadman, P.E. Toorop, J. Lynn, and H. Hilhorst. 2007. Seed dormancy release in *Arabidopsis Cvi* by dry after-ripening, low temperature, nitrate and light shows common quantitative patterns of gene expression directed by environmentally specific sensing. *Plant Jour. Lond.* 51:60–78. doi:10.1111/j.1365-313X.2007.03118.x.

 Updates

[Google Scholar](#)

6. Fulbright, T. 1992. Temperature Effects on Seed Germination of *Cordia boissieri* A. DC. (*Boraginaceae*). *Southwest. Nat.* 37(2):197–199. doi: 10.2307/3671669.

[Web of Science ®](#) | [Google Scholar](#)

7. Gao, N., G.F. Cui, Y.Q. Lai, S.X. Zhang, J. Li, J.H. Wang, and F.H. Liu. 2011. Effects of Different Treatments on the Germination of Oriental lily Seeds. *Acta Agric. Univ. Jiangxi* 33:660–664. (in Chinese with English abstract).

[Google Scholar](#)

8. Gashi, B. Abdullai, K. Mata, V and Kongjika, E. 2012. Effect of gibberellic acid and potassium nitrate on seed germination of the resurrection plants *Ramonda serbica* and *Ramonda nathaliae*. *African Journal of Biotechnology* 11(20): 4537-4542.

[Google Scholar](#)

9. Gniazdowska, A., U. Dobrzyjska, T. Babajczyk, and R. Bogatek. 2007. Breaking the apple embryo dormancy by nitric oxide involves the stimulation of ethylene

20. Gumgumjee, N.M., and A.S. Hajar. 2015. Antimicrobial efficacy of *Acacia saligna* (Labill.) H.L.Wendl. and *Cordia sinensis* Lam. Leaves extracts against some pathogenic microorganisms. *Int. J. Microbiol. Immunol. Res* 3(4):51–57.

[Google Scholar](#)

21. Gupta, S.M., P. Pandey, A. Grover, and Z. Ahmed. 2011. Breaking seed dormancy in *Hippophae salicifolia*, a high-value medicinal plant. *Physiol. Mol. Biol. Plants*. 17:403–406. doi:10.1007/s12298-011-0082-6.

22. Harsha, H.R., R. Venkat, K.J. Dayamani, and M. Shivanna. 2017. Effect of growth regulators and macronutrients on seedling growth of pummelo (*Citrus maxima* Merrill). *Int. J Recent Sci. Res.* 8(10):20531–20533. doi: 10.24327/IJRSR.

[Google Scholar](#)

23. Iralu, V., and K. Upadhaya. 2018. Seed dormancy, germination and seedling characteristics of *Elaeocarpus prunifolius* Wall. ex Müll. Berol.: A threatened tree species of north-eastern India. *N.Z. J. Of For. Sci.* 48:16. doi:10.1186/s40490-018-0121-y.

24. Jaiswal, S.B., R.V. Nainwad, S.J. Supekar, and S.B. Mane. 2018. Effect of growth regulators and chemicals on growth of Kagzi Lime (*Citrus aurantifolia* Swingle.) seedlings. *Int.J.Curr.Microbiol.App.Sci.* 6:940–944.

[Google Scholar](#)

25. Kant, G., R.P.S. Dalal, and B.S. Beniwal. 2017. Effect of PGR on leaf nutrients content and root growth in rough lemon (*Citrus jambhiri*) seedling. *Int. J. Pure App. Biosci.* 5(4):346–351. doi: 10.18782/2320-7051.5386.

[Google Scholar](#)

6. Kazemi, M. 2014. Effect of gibberellic acid and potassium nitrate spray on vegetative growth and reproductive characteristics of tomato. J. Biol. Environ. Sci. 8(22):1-9.

[Google Scholar](#)

7. Keskiner, K., and B. Tuncer. 2019. Dormancy breaking treatments for wild *Eremurus spectabilis* M.Bieb seeds. Fresen. Environ. Bull. 28(2A):1167-1173.

[Web of Science ®](#) | [Google Scholar](#)

8. Koyuncu, F. (2005). Breaking seed dormancy in black mulberry (*Morus nigra* L.) by cold stratification and exogenous application of gibberellic acid. Acta Biol. Cracoviensia Ser. Bot. 47(2):23-26.

[Web of Science ®](#) | [Google Scholar](#)

9. Kuria, S.G., M.M. Wanyoike, C.K. Gachuiri, and R.G. Wahome. 2005. Nutritive value of important range forage species for camels in marsabit district, Kenya. Trop. Subtrop. Agroecosyst. 5:15-24.

[Google Scholar](#)

10. Lay, P., G.V. Basvaraju, V. Pashte, and M. Gowri. 2015. Studies on effect of gibberellic acid (GA3) and potassium nitrate (KNO3) on breaking of seed dormancy of papaya (*Carica papaya* L.) cv. Surya. The Ecoscan 9:109-113.

[Google Scholar](#)

11. Lin, D., and H. Danfeng. 2003. Effects of potassium levels on photosynthesis and fruit quality of muskmelon in culture medium. Acta Horticulturae Sinica 30(2):221-223.

[Google Scholar](#)

12. Majumdar, F. 2013. Response of gibberellic acid and potash nutrient on growth and yield of late planting cabbage (*Brassica oleracea Capitata*). MS thesis, Department of Horticulture, Sher-e-Bangla Agricultural University, Dhaka, Bangladesh, pp. 62-84.

[Google Scholar](#)

33. Marschner, P. 2012. Marschner's mineral nutrition of higher plants. 3rd. Academic Press, London. p. 672 p.

[Google Scholar](#)

34. Maundu, P.M., B. Tengnäs, N. Muema, and A. Birnie. (2005). *Cordia sinensis* (C. gharaf, C. rothii) in Useful trees and shrubs for Kenya World Agroforestry Centre, Eastern and Central Africa Regional Programme: Nairobi 173–174 Retrieved December 5, 2018, from http://www.worldagroforestry.org/usefultrees/pdflib/Cordia_sinensis

[Google Scholar](#)

35. Meena, R.R., and M.C. Jain. 2012. Effect of seed treatment with gibberellic acid on growth parameters of papaya seedling (*Carica papaya* L.). *Prog. Hort* 44(2):248–250.

[Google Scholar](#)

36. Meghwal, P.R. 2006. Propagation study in Lehsua (*Cordia myxa*), p. 117–119. In: S.N. Ghosh, S.K. Mitra, B.C. Banik, M.A. Hasan, S.K. Sarkar, R.S. Dhua, J. Kabir, and J.K. Hore eds. Proceedings of the national symposium on production, utilization and export of underutilized fruits with commercial potentialities. 22–24 November 2006. Kalyani, Nadia, West Bengal, India.

[Google Scholar](#)

37. Meghwal, P.R. 2007. Propagation studies in lehsua (*Cordia myxa*). *Indian Journal of Agricultural sciences* 77 (11) : 765–767

[Web of Science ®](#) | [Google Scholar](#)

38. Meghwal, P.R., S. Akath, P. Kumar, and B.R. Morwal. 2014. Diversity, distribution and horticultural potential of *Cordia myxa* L.: A promising underutilized fruit species of arid and semi-arid regions of India. *Genet. Resour. Crop Evol.* 61:1633–1643. doi:10.1007/s10722-014-0161-y.

[Web of Science ®](#) | [Google Scholar](#)

39. Mello, A.M., N.A. Streek, E.E. Blankenship, and E.T. Paparozzi. 2009. Gibberellic acid promotes seed germination in *Penstemon digitalis* cv. Husker Red. *HortSci* 44(3):870–873. doi: 10.21273/HORTSCI.44.3.870.

[Web of Science ®](#) | [Google Scholar](#)

40. Meshram, P.C., P.S. Joshi, R.K. Bhoyar, and A.K. Sahoo. 2015. Effect of different plant growth regulators on seedling growth of acid lime (*Citrus aurantifolia* Swingle). *Res. Envrn. Life Sci.* 8(4):725–728.

[Google Scholar](#)

41. Moradi, D.P., F. Sharif-zadeh, and M. Janmohammadi. 2008. Influence of priming techniques on seed germination behavior of maize inbred lines (*Zea mays* L.). *J. Agr. Biol. Sci.* 3(3):22–25.

[Google Scholar](#)

42. Pal, P., K. Yadav, K. Kumar, and N. Singhm. 2016. Effect of gibberellic acid and potassium foliar sprays on productivity and physiological and biochemical parameters of parthenocarpic cucumber (*Cucumis sativus*) cv. 'seven star F1. *J. Horticultural Res.* 24(1):93–100. doi: 10.1515/johr-2016-0011.

[Google Scholar](#)

43. Parab, A.M., J.C. Mathad, and K.V. Malshe. 2017. Effect of pre-soaking chemicals on germination and subsequent seedling growth of papaya (*Carica papaya* L.) cv. Solo. *Int. J. Chem. Stud.* 5(4):1812–1816.

[Google Scholar](#)

44. Patel, M., R.V. Tank, D.R. Bhanderi, H.M. Patil, V. Patel, and M. Desai. 2018. Response of soaking time and chemicals on germination and growth of tamarind (*Tamarindus indica* L.). *Plant Archives* 18:51–56.

[Google Scholar](#)

45. Patil, A.S., S.R. Patil, and A.S. Kale. 2015. Influence of growth Regulators on

j.graham under shade net conditions. Global J. Res. Med. Plants Indigen. Med. 4(1):01-09.

[Google Scholar](#)

46. Pradip, K. and Singh H.S. 2010. Tree wealth of the non-forest areas of Gujarat. A report tree census in non-forest areas-2009. Forest department, Gujarat state. p 47

[Google Scholar](#)

47. Ramzan, A., I.A. Hafiz, T. Ahmad, and N.A. Abbasi. 2010. Effect of priming with potassium nitrate and dehusking on seed germination of gladiolus (*Gladiolus alatus*). Pak. J. Bot. 42:247-258.

[Web of Science ®](#) | [Google Scholar](#)

48. Schuch, U.K., E. Davison, and J. Kelly. 2001. Seed Propagation of *Cordia boissieri* and *Cordia parvifolia*. Turfgrass and Ornamental Research Report.

<http://ag.arizona.edu/pubs/crops/az1246>

[Google Scholar](#)

49. Shabaq, M.N.A. 2013. The role of the different concentrations of GA3 on seed germination and seedling growth of loquat (*Eriobotrya japonica* L.). IOSR J. Agric. Vet. Sci. 4(5):03-06. doi: 10.9790/2380-0450306.

[Google Scholar](#)

50. Shanmugavalli, M. Renganayaki, P.R, and Menaka, C. 2007. Seed dormancy and germination improvement treatments in fodder sorghum. International Crops Research Institute for the Semi-Arid Tropics 3: 1-3.

[Google Scholar](#)

51. Sheoran, O.P., D.S. Tonk, L.S. Kaushik, R.C. Hasija, and R.S. Pannu. 1998. Statistical Software Package for Agricultural Research Workers, p. 139-143. In: D.S. Hooda and R.C. Hasija (eds.). Recent advances in information theory, statistics & computer applications. Department of Mathematics Statistics, CCS HAU, Hisar.

52. Singh, R. 2018. Activities of research wing of Haryana forest department. Retrieved from http://haryanaforest.gov.in/Portals/0/Research_Activities_Final%20as%20on%2031-3-2018.pdf

[Google Scholar](#)

53. Sirova, J., M. Sedlarova, J. Piterkova, L. Luhova, and M. Petrivalsky. 2011. The role of nitric oxide in the germination of plant seeds and pollen. *Plant Sci.* 181:560–572. doi:10.1016/j.plantsci.2011.03.014.

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

54. Song, J.Q., and H. Fujiyama. 1996. Ameliorative effect of potassium on rice and tomato subjected to sodium salinization. *Soil Sci. Plant Nutr.* 42:493–501. doi:10.1080/00380768.1996.10416318.

[Web of Science ®](#)

[Google Scholar](#)

55. Tuan, P.A., R. Kumar, P.K. Rehal, P.K. Toora, and B.T. Ayele. 2018. Molecular mechanisms underlying abscisic acid/gibberellin balance in the control of seed dormancy and germination in cereals. *Front. Plant Sci.* 9:668. doi:10.3389/fpls.2018.00668.

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

56. Varner, J.E. 1965. Enzymes, p. 14–20. In: J. Bonner and J.E. Varner (eds.). *Plant Biochemistry*. Academic Press, London and New York.

[Google Scholar](#)

57. Waman, A.A., P. Bohra, and A. Norman. 2017. Chemical pre-treatments improve seed germination and seedling growth in *Semecarpus kurzii*: An ethnomedicinally important plant. *J. For. Res.* 29:1283. doi:10.1007/s11676-017-0562-9.

[Google Scholar](#)

58. Yücel, E., and G. Yilmaz. 2009. Effects of different alkaline metal salts (Na Cl, KNO₃),

[Google Scholar](#)

59. Zhang, A., D.F. Huang, and Z. Hou. 2002. Effect of potassium nutrient on development and photosynthesis of melon plant. J. Shanghai Agricultural Coll. 20(1):13–17.

[Google Scholar](#)

[Download PDF](#)

Related research

People also read

Recommended articles

Cited by
8

Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research
- Help and information
- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright © 2025 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)
[Accessibility](#)



Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG