

International Journal of Fruit Science >
Volume 20, 2020 - Issue sup2

✓ Free access

3,671 4

Views

CrossRef citations to date

0

Altmetric

Listen

Research Article

Gibberellic Acid and Potassium Nitrate Promote Seed Germination and Growth of Grey-leaved Saucer-berry (*Cordia Sinensis* Lam.) Seedlings

Rahul Dev , Devi Dayal & M. Sureshkumar

Pages S937-S954 | Published online: 11 Jun 2020

“ Cite this article

<https://doi.org/10.1080/15538362.2020.1774465>

Check for updates

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

View PDF

View EPUB

ABSTRACT

Settings

Formulae display: ☒ MathJax

We Care About Your Privacy

We and our 845 partners store and/or access information on a device, such as unique IDs in cookies to process personal data. You may accept or manage your choices by clicking below, including your right to object where legitimate interest is used, or at any time in the privacy policy page. These choices will be signaled to our partners and will not affect browsing data. [Privacy Policy](#)

We and our partners process data to provide:

Use precise geolocation data. Actively scan device characteristics for identification. Store and/or access information on a device. Personalised advertising and content, advertising and content measurement, audience research and services development.

List of Partners (vendors)

I Accept

Essential Only (10–60%) and

Show Purpose

KEYWORDS: Cordia sinensis gibberellic acid potassium nitrate seed germination

Cordia sinensis Lam. (*Cordia gharaf* (Forssk.) Ehrenb. ex Asch.) is a multi-stemmed tree, growing up to 8 to 12 m height. *Cordia sinensis* is a potential underutilized plant which is also known as Gondi, narrow-leaved sebastian or Greyleaved saucer-berry and belongs to the family Boraginaceae. The genus *Cordia* consists of about 250 species and the majority of species are tree or shrub (Barroso and Oliveira, [2009](#)). It is well-adapted plant species in arid to semiarid regions of West Africa to East to the Middle East, India, Sri Lanka, and Pakistan due to its various adaptive features e.g. strong tap root system, waxy leaves, hairiness, sunken and covered stomata in leaves, water binding mechanism, moderate tolerance to salinity and alkalinity. These traits make it hardier than *Cordia myxa*, a popular fruit plant of the Boraginaceae family (Meghwal et al., [2014](#)). The fruits of the *C. sinensis* Lam. are consumed by local inhabitants. The leaves provide the protein (13.12%) rich fodder to cattle (Kuria et al., [2005](#)). Various parts of the plant have ethnomedicinal value and are traditionally used to cure different human and livestock ailments (Gumgumiee and Hajar, [2015](#)). Farmers of arid regions grow *C.*

not and dry
ies for
f *C. sinensis*
on of the
C. sinensis was
is reported
(Haryana)
ut
ous
eri reveals
tification



The proper seedling growth of germinated seeds is the most important factor for successful sapling production under nursery conditions. The earlier studies showed the variable effect of GA₃ and KNO₃ on growth traits of various plants. Application of GA₃ and KNO₃ promotes overall plant growth in *Brassica oleracea Capitata* (Majumdar, [2013](#)), *Cucumis sativus* (Pal et al., [2016](#)), and *Solanum lycopersicum* (Balaguera-Lopez et al., [2009](#)). GA₃ (250 mg L⁻¹) treatment resulted in maximum seed germination (98.75%), subsequent shoot length, root length, leaf area, shoot, and root dry weight (Shabaq, [2013](#)). Similar growth promotion of seedlings was also reported under the GA₃ treated seeds of *Citrus limon* (Dzayi and Rahman, [2010](#)), *Nothapodytes nimmoniana* (Patil et al., [2015](#)), and *Citrus aurantifolia* (Jaiswal et al., [2018](#)). Lay et al. ([2015](#)), found GA₃ and KNO₃ presowing treatment to seeds of papaya improves the seedling growth, fresh and dry weight of seedlings. However, very few studies are available with quantitative data on the subsequent growth of treated plants. The *C. sinensis* is very



The stud
Institute

carried out under the shade net. *Cordia sinensis* fruits were collected from the native population in the Kachchh region of Gujarat (22°84'624 – 23°56'288 N° latitude and 68°95'730 – 70°73'100 E° longitude). Fresh seeds were extracted from the ripened fruits. The mucilaginous pulp was removed from the seeds by rubbing the seeds with sandy soil followed by washing with tap water. Air dried, cleaned seeds were stored in the paper bag for 2 weekss at the room temperature before use.

Experiment 1. Chemical Treatment of Seeds

Before the seed treatment, seeds were disinfected with fungicide (Carbendazim 2%) for 20–30 minutes and rinsed three times with running tap water. For the treatments, seeds were divided into four groups, the total no. of treatment combinations were 16, four levels of each of GA₃ concentration (0, 250, 500, and 1000 ppm) and four KNO₃ concentrations (0%, 0.25%, 0.5%, and 1% KNO₃) purchased from Sigma Aldrich (90% gibberellin A₃ basis) Loba Chemie (99% purity), respectively and were evaluated with 3 replications 1) seeds of first group were transferred into the beaker filled with distilled water and left for 24 h; 2) seeds of second group were soaked in 250, 500, and 1000 mg L⁻¹ (w/v) GA₃ solution for 24 h; 3) similarly, seeds of the third group were soaked in 0.25%, 0.5%, and 1% KNO₃ solution for 24 h; 4) the seeds of the fourth group were treated with the aqueous solution supplemented different combinations of GA₃ and KNO₃ (Table 1).

Table 1. Test design of chemicals (KNO₃, GA₃) and combinations of KNO₃ and GA₃ treatments

Download

The prev
treatme
aver
wool wa
size of b
placed in
three Pe
by frequ

test. The 16
ications. The
yer of cotton
with equal
ton wool
atment with
t uniformly
daily basis

Statistical Analysis

The randomized complete block design (RCBD) was used for conducting the experiment in laboratory and shade net. The critical differences (CD) at $P < .05$ and $P < .01$ level of probability were determined after analysis of variance (ANOVA) for all treatments. The experiment was laid out with 16 treatments including control with three replications and 60 seeds per replicate were used. The mean and analysis of variance were performed to determine significant differences among treatments ($P < .05$) using the Statistical Software Package for Agricultural Research Workers (Sheoran et al., [1998](#)).

Result and Discussion

Experiment 1. Chemical Treatment of Seeds

Mean Germination Time (MGT) and Germination Percentage (GP)

The different concentrations of GA₃ and KNO₃ significantly fasten the seed germination as compared to the germination percentage of control seeds. The KNO₃ (0.5%) treatment significantly shortens the germination initiation duration (4 days), compared to other single KNO₃ applications. Whereas the synergistic effect of GA₃ × KNO₃ treatments was observed on the shortening of germination time. The seeds treated with aqueous solution supplemented with GA₃ 250 mg L⁻¹ and 1% KNO₃, took minimum time for germination initiation (4.33 ± 0.33 days), which was about 1 week less than the control. The maximum germination percentage was observed in 0.25% and 1.0% concentrations of GA₃ and KNO₃ respectively.



Download

The results of final

In this article



Table 3. Mean of germination, growth (MGT, GP, LA, LFW, LDW) and seedling nutrients (NPK and Na) of *Cordia sinensis* under various levels of GA₃ (in ppm) and KNO₃ (in %)

Display Table




germination (66.66%). However, KNO_3 did not influence the seed germination, irrespective of treatment duration and chemical concentration. Gashi et al. (2012) found the higher germination percentage (92.26%) in the *Ramonda* species' seeds treated with 1000 ppm GA_3 + 0.3% KNO_3 . Treatment of *Sorghum bicolor* seeds with GA_3 (100 ppm) and KNO_3 (1%) resulted in increase of germination percentage to a higher extent (94%) (Shanmugavalli et al., 2007). Similar synergistic results of KNO_3 (1%) and GA_3 (500 ppm) seed treatments were reported by Dewir et al. (2011) in *Sabal palmetto*. Whereas, soaking cracked seeds of *Elaeocarpus prunifolius* in the solution of GA_3 and KNO_3 accelerate the germination rate (Iralu and Upadhaya, 2018).

Experiment 2. Subsequent Quantification of Growth

Plant Height and Collar Diameter


Application of $\text{GA}_3 \times \text{KNO}_3$ singly or in combination, significantly ($p = .05$) affected the plant height (Figure 1). However, plant height under different single KNO_3 applications differed significantly. Lower KNO_3 application (0.25% conc.) tends to increase plant height at par to GA_3 500 mg L^{-1} , but at higher KNO_3 application (1%), there was a reduction in the plant height. The longest plants were observed from the GA_3 (500 mg L^{-1}) and KNO_3 (1%) applications, followed by combined application of 500 mg L^{-1} GA_3 and 0.5% KNO_3 . Whereas, the minimum increase in the plant height was seen when plants seeds soaked in the GA_3 1000 mg L^{-1} and KNO_3 1% solutions for 24 h.

Figure 1. Plant height of *Cordia sinensis*, under various levels of GA_3 (in ppm) and KNO_3 (in %) seed treatment, recorded at three month interval



Data presented in Figure 2 illustrated the effect of GA₃ and KNO₃ on the collar diameter of *C. sinensis*. The significant difference was ascertained for *C. sinensis* between seeds treated with GA₃, KNO₃ and H₂O (control). All GA₃ and KNO₃ applications had a positive effect on collar diameter. However, single GA₃ applications were more effective than different single KNO₃ application. Moreover, integrated application of GA₃ and KNO₃ resulted pronounced improvement in collar diameter of plantlets. The maximum increase in the collar diameter in one-year-old plants was observed with the GA₃ 500 mg L⁻¹ and KNO₃ 0.5% treatment followed by GA₃ 1000 mg L⁻¹ and KNO₃ 0.25% treatment. On the contrary, the minimum collar diameter was recorded in the plants of control group. The presowing treatment with lower concentration of GA₃ and KNO₃ improved the seedling height significantly however, reduction in the plant height was observed under the higher concentration of GA₃ and KNO₃. Treatment of *Eriobotrya japonica* seeds with lower concentration of GA₃ (250 mg L⁻¹) resulted maximum collar diameter and plant height comparing to control (Shabaq, 2013). Similarly, maximum plant height and stem diameter of *Citrus aurantifolia* plantlets were observed with GA₃ (200 mg L⁻¹) seed treatment (Meshram et al., 2015).

Figure 2. Collar diameter of *Cordia sinensis*, under various levels of GA₃ (in ppm) and KNO₃ (in %) seed treatment, recorded at three month interval



The GA₃ and KNO₃ treatments significantly enhanced the collar diameter of *C. sinensis* plantlets. The GA₃ 500 mg L⁻¹ and KNO₃ 0.5% treatment resulted in the maximum collar diameter, followed by GA₃ 1000 mg L⁻¹ and KNO₃ 0.25% treatment. The control group showed the minimum collar diameter. The integrated application of GA₃ and KNO₃ resulted in a pronounced improvement in collar diameter of plantlets. The maximum increase in the collar diameter in one-year-old plants was observed with the GA₃ 500 mg L⁻¹ and KNO₃ 0.5% treatment followed by GA₃ 1000 mg L⁻¹ and KNO₃ 0.25% treatment. On the contrary, the minimum collar diameter was recorded in the plants of control group. The presowing treatment with lower concentration of GA₃ and KNO₃ improved the seedling height significantly however, reduction in the plant height was observed under the higher concentration of GA₃ and KNO₃. Treatment of *Eriobotrya japonica* seeds with lower concentration of GA₃ (250 mg L⁻¹) resulted maximum collar diameter and plant height comparing to control (Shabaq, 2013). Similarly, maximum plant height and stem diameter of *Citrus aurantifolia* plantlets were observed with GA₃ (200 mg L⁻¹) seed treatment (Meshram et al., 2015).


The single factors (GA_3 and KNO_3 concentration) significantly influenced the leaf area ($p \leq 0.05$) ([Table 3](#)). Seeds treated with the different concentrations of GA_3 are found to have more leaf area as compare to KNO_3 treated seed and untreated seeds (control). A lower concentration of GA_3 (250 mg L^{-1}) had a favorable effect on leaf expansion. While at a higher concentration of GA_3 treatment leaf area found to be non-significantly decreased. Similar to GA_3 , leaf area of treated KNO_3 seeds was declined with increases of KNO_3 concentration. The GA_3 and KNO_3 treatments significantly affected the leaf area. The maximum leaf area was recorded in the plants treated with $\text{GA}_3 250 \times \text{KNO}_3 1\%$, followed by $\text{GA}_3 500 \times \text{KNO}_3 0.5\%$. While, treatment combination, $\text{GA}_3 1000 \text{ mg L}^{-1} \times \text{KNO}_3 0.25\%$ had some adverse effect on leaf expansion, and it resulted in minimum leaf area. Similar, leaf expansion at lower GA_3 concentration (250 mg L^{-1}) was observed in *Eriobotrya japonica* (Shabaq, [2013](#)), *Citrus limon* (Dzayi and Rahman, [2010](#)) and *Citrus aurantifolia* (Meshram et al., [2015](#)).

In general, numbers of leaves per plant were not improved by the pretreatment of GA_3 and KNO_3 ([Figure 3](#)). However, numbers of leaves per plant were significantly improved with combined application of GA_3 and KNO_3 . The presowing treatment of seeds with $\text{GA}_3 500 \text{ mg L}^{-1} \times \text{KNO}_3 0.5\%$, led to maximum increase in the leaf number per plant, followed by seeds treated with $\text{GA}_3 250 \text{ mg L}^{-1} \times \text{KNO}_3 0.5\%$. In contrast, GA_3 seeds soaked in higher gibberellin concentration (1000 mg L^{-1}) supplemented with medium or high KNO_3 (0.5% or 1%) concentration for 24 h, gave the minimum increase in the leaf numbers per plant. The various previous study reported the opposite relations between leaf growth and plant height increment due to GA_3 treatment (Arney and Mancinelli, [1966](#)). These trends are concurrent with the present study. GA_3

pregerm... a papaya
(Parab... y due to the
stimulat... [2017](#)).

Figure 3. ... GA_3 (in ppm)
and KNO_3





Display full size

Leaf Fresh and Dry Weight

The result indicated that the highest leaf fresh and dry weight was achieved when the *C. sinensis* seeds were treated with the aqueous solution of GA₃ 250 mg L⁻¹, supplemented with the KNO₃ 1%, followed by the solution containing GA₃ 500 mg L⁻¹ and KNO₃ 0.25%. The minimum leaf fresh weight was noted under the highest KNO₃ (1% conc.) treatment. In general, fresh weight of leaf was higher when seeds were treated with GA₃ than a KNO₃ solution; further fresh weight was decreased when GA₃ concentration increased from 0.25% to 1%. While the lowest dry weight was found under maximum concentration of GA₃ treatment and KNO₃. It was observed that higher concentration of both the chemicals (GA₃ and KNO₃) resulted in the decline in the dry weight of plant weight, specific, higher, of Carica, rsha et al., GA₃ (250, mer shoot, and h and dry, mobilization, h ultimately, arsha et al.,

[2017](#)). While, higher mobilization of photosynthetic products was also taken place in the KNO_3 treated seeds in the early stage of seed germination (Lay et al., [2015](#)).

Leaf Nutrient Traits

The different concentration of $\text{GA}_3 \times \text{KNO}_3$ affects the leaf N, P, K, and Na content significantly ([Table 3](#)). The leaf N content was lower under the different single GA_3 and KNO_3 treatments as compared to control. However, combination treatment of GA_3 1000 mg L^{-1} and KNO_3 0.25% resulted in highest leaf N content. In general leaf N content was higher in the plants treated with GA_3 as compare to KNO_3 treatments. The leaf P content was higher under the GA_3 treated plants it was highest under GA_3 250 mg L^{-1} treatment closely followed by GA_3 500 $\text{mg L}^{-1} \times \text{KNO}_3$ 1% treatment. The combined application of GA_3 1000 $\text{mg L}^{-1} \times$ 0.5% KNO_3 resulted minimum leaf P content. In general higher leaf P content was achieved when seeds were treated with GA_3 for 24 h than KNO_3 treatment for the same duration. Percentage leaf K content was lower for GA_3 and KNO_3 treated seeds as compared to seeds treated with distilled water (control). However, among the treatments, the highest K content was noted when seeds were soaked in GA_3 500 mg L^{-1} followed by KNO_3 (0.25%). The lowest leaf K content was reported when the seeds were soaked in the aqueous solution of GA_3 1000 mg L^{-1} supplemented with KNO_3 1% concentration. The treatment of GA_3 resulted significantly high leaf Na content than control. However, there was no significant difference between seeds soaked in H_2O (control) and KNO_3 solution for leaf Na content. The seeds soaked in GA_3 (500 mg L^{-1}) and KNO_3 0.25% had the highest leaf Na content. Whereas, lowest leaf Na content was recorded when seeds were treated with an

0.25% for 24 h.

Increase in leaf N content was observed in GA_3 and KNO_3 wa higher leaf nitrogen content. GA₃ combination of GA_3 (1000 mg L^{-1}) and KNO_3 (0.25%) (semi ([2014](#))) noted the the applicati opersicum leaves. T similar effect was obs on of



were responsible for increase in leaf NPK contents (Marschner, [2012](#)). Leaf sodium content was found to be decreased with combined application of gibberellins and potassium nitrate in the study. The similar antagonistic correlation was also observed by Song and Fujiyama ([1996](#)).

Thus, the external application of KNO_3 and GA_3 showed a synergistic effect in promoting seed germination and seedling vigor of *Cordia sinensis* seed in the present investigation. In conclusion, it can be stated that the application of GA_3 (250 ppm) and KNO_3 (1%) is a simple, effective, and practical method for improving seed germination and seedling growth of *Cordia sinensis*. Although, further work might be useful for in-depth understanding on effects of chemicals in *Cordia sinensis*.

Acknowledgments

Author is thankful to Director, ICAR-CAZRI for proving necessary facility for conducting research.

Disclosure Statement

The authors declare that they have no conflict of interest.

Literature

1. Achar, S. K. (2018). Effect of gibberellins and potassium nitrate on seed germination and seedling vigor of *Cordia sinensis* seed. *Journal of Horticulture and Forestry*, (Vol. 66, No. 1), 1-10. [Google Scholar](#)
2. Al-Hav, S. (2018). Effect of gibberellins and potassium nitrate on seed germination and seedling vigor of *Cordia sinensis* seed. *Journal of Horticulture and Forestry*, (Vol. 66, No. 1), 1-10. [Google Scholar](#)

3. Arney, S.E., and P. Mancinelli. 1966. The basic action of gibberellic acid in elongation of 'Meteor' pea stems. *New Phytol.* 65:161–175.

 | [Web of Science](#)® | [Google Scholar](#)

4. Balaguera-Lopez, H.E., J.F. Cardenas-Hernandez, and J.G. Alvarez-Herrera. 2009. Effect of gibberellic acid (GA3) on seed germination and growth of tomato (*Solanum lycopersicum* L.). *Acta Hort. (ISHS)*. 821:141–148. doi:10.17660/ActaHortic.2009.821.15.

 | [Google Scholar](#)

5. Barroso, I.C.E., and F.D. Oliveira. 2009. Pharmacognostic diagnosis of fruits of *Cordia sellowiana* Cham. and *Cordia myxa* L. (Boraginaceae). *Rev. Bras. Farmacogn.* 19:458–470. doi:10.1590/S0102-695X2009000300021.

 | [Google Scholar](#)

6. Bian, L., L. Yang, J. Wang, and H. Shen. 2013. Effects of KNO₃ pretreatment and temperature on seed germination of *Sorbus pohuashanensis*. *J. Forest Res.* 24:309–316. doi:10.1007/s11676-013-0354-9.

 | [Google Scholar](#)

7. Çetinbaş, M. and Koyuncu, F. 2006. Improving germination of *Prunus avium* L. seeds by gibberellic acid (GA3). *Acta Hort. (ISHS)*. 703:1–5. doi:10.17660/ActaHortic.2006.703.1.

 | [Google Scholar](#)

8. Cárdenas-Hernández, J.F., Balaguera-López, H.E., and Álvarez-Herrera, J.G. 2009. Effect of gibberellic acid (GA3), KNO₃, and temperature on seed germination and growth of tomato (*Solanum lycopersicum* L.). *Acta Hort. (ISHS)*. 821:141–148. doi:10.17660/ActaHortic.2009.821.15.

9. Chen, Y., and J. Wang. 2013. Effects of KNO₃ pretreatment and temperature on seed germination of *Sorbus pohuashanensis*. *J. Forest Res.* 24:309–316. doi:10.1007/s11676-013-0354-9.

- Google Scholar

- Google Scholar

- Google Scholar

- Web of Science ® | Google Scholar

- Goog

- 

6. Fulbright, T. 1992. Temperature Effects on Seed Germination of *Cordia boissieri* A. DC. (Boraginaceae). Southwest. Nat. **37**(2):197–199. doi: 10.2307/3671669.

 | [Web of Science ®](#) | [Google Scholar](#)

7. Gao, N., G.F. Cui, Y.Q. Lai, S.X. Zhang, J. Li, J.H. Wang, and F.H. Liu. 2011. Effects of Different Treatments on the Germination of Oriental lily Seeds. *Acta Agric. Univ. Jiangxi* 33:660–664. (in Chinese with English abstract).

[Google Scholar](#)

8. Gashi, B. Abdullai, K. Mata, V and Kongjika, E. 2012. Effect of gibberellic acid and potassium nitrate on seed germination of the resurrection plants *Ramonda serbica* and *Ramonda nathaliae*. *African Journal of Biotechnology* 11(20): 4537-4542.

[Google Scholar](#)

9. Gniazdowska, A., U. Dobrzyjska, T. Babajczyk, and R. Bogatek. 2007. Breaking the apple embryo dormancy by nitric oxide involves the stimulation of ethylene production. *Planta*. **225**:1051–1057. doi:10.1007/s00425-006-0384-z.

 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

10. Gumgumjee, N.M., and A.S. Hajar. 2015. Antimicrobial efficacy of *Acacia saligna* (Labill.) H.L.Wendl. and *Cordia sinensis* Lam. Leaves extracts against some pathogenic microorganisms. *Int. J. Microbiol. Immunol. Res* 3(4):51–57.

[Google Scholar](#)

11. Gupta, N., and A.S. Hajar. 2015. Antimicrobial efficacy of *Acacia saligna* (Labill.) H.L.Wendl. and *Cordia sinensis* Lam. Leaves extracts against some pathogenic microorganisms. *Int. J. Microbiol. Immunol. Res* 3(4):51–57.

12. Ha, N., and A.S. Hajar. 2015. Antimicrobial efficacy of *Acacia saligna* (Labill.) H.L.Wendl. and *Cordia sinensis* Lam. Leaves extracts against some pathogenic microorganisms. *Int. J. Microbiol. Immunol. Res* 3(4):51–57.

23. Iralu, V., and K. Upadhaya. 2018. Seed dormancy, germination and seedling characteristics of *Elaeocarpus prunifolius* Wall. ex Müll. Berol.: A threatened tree species of north-eastern India. *N.Z. J. Of For. Sci.* **48**:16. doi:10.1186/s40490-018-0121-y.

 | [Web of Science ®](#) | [Google Scholar](#)

24. Jaiswal, S.B., R.V. Nainwad, S.J. Supekar, and S.B. Mane. 2018. Effect of growth regulators and chemicals on growth of Kagzi Lime (*Citrus aurantifolia* Swingle.) seedlings. *Int.J.Curr.Microbiol.App.Sci.* **6**:940–944.

[Google Scholar](#)

25. Kant, G., R.P.S. Dalal, and B.S. Beniwal. 2017. Effect of PGR on leaf nutrients content and root growth in rough lemon (*Citrus jambhiri*) seedling. *Int. J. Pure App. Biosci.* **5**(4):346–351. doi: 10.18782/2320-7051.5386.

 | [Google Scholar](#)

26. Kazemi, M. 2014. Effect of gibberellic acid and potassium nitrate spray on vegetative growth and reproductive characteristics of tomato. *J. Biol. Environ. Sci.* **8**(22):1–9.

[Google Scholar](#)

27. Keskiner, K., and B. Tuncer. 2019. Dormancy breaking treatments for wild *Eremurus spectabilis* M.Bieb seeds. *Fresen. Environ. Bull.* **28**(2A):1167–1173.

[Web of Science ®](#) | [Google Scholar](#)

28. Koyun, M., and A. K. Koyun. 2019. Effect of gibberellic acid (GA3) on vegetative growth and reproductive characteristics of *Crataegus* (Rosaceae) by cold stratification. *Acta Hort.* **1254**:11–16. doi:10.3182/2019051254(01001)11-16.

[Web of Science](#)

29. Kumar, S., and S. Kumar. 2019. Effect of gibberellic acid (GA3) on vegetative growth and reproductive characteristics of *Crataegus* (Rosaceae) by cold stratification. *Acta Hort.* **1254**:11–16. doi:10.3182/2019051254(01001)11-16.

[Google Scholar](#)

30. Lay, P., G.V. Basvaraju, V. Pashte, and M. Gowri. 2015. Studies on effect of gibberellic acid (GA3) and potassium nitrate (KNO3) on breaking of seed dormancy of papaya (*Carica papaya* L.) cv. Surya. *The Ecoscan* 9:109–113.

[Google Scholar](#)

31. Lin, D., and H. Danfeng. 2003. Effects of potassium levels on photosynthesis and fruit quality of muskmelon in culture medium. *Acta Horticulturae Sinica* 30(2):221–223.

[Google Scholar](#)

32. Majumdar, F. 2013. Response of gibberellic acid and potash nutrient on growth and yield of late planting cabbage (*Brassica oleracea Capitata*). MS thesis, Department of Horticulture, Sher-e-Bangla Agricultural University, Dhaka, Bangladesh, pp. 62–84.

[Google Scholar](#)

33. Marschner, P. 2012. Marschner's mineral nutrition of higher plants. 3rd. Academic Press, London. p. 672 p.

[Google Scholar](#)

34. Maundu, P.M., B. Tengnäs, N. Muema, and A. Birnie. (2005). *Cordia sinensis* (C. gharaf, C. rothii) in Useful trees and shrubs for Kenya World Agroforestry Centre, Eastern and Central Africa Regional Programme: Nairobi 173–174 Retrieved December 5, 2018, from

http://www.worldagroforestry.org/usefultrees/pdflib/Cordia_sinensis




[Goog](#)

35. Meena, S.K., and S.K. Singh. 2013. Effect of gibberellic acid on growth and yield of muskmelon (*Cucumis melo* L.). *Acta Horticulturae Sinica* 30(2):248–250.

[Goog](#)

36. Meena, S.K., and S.K. Singh. 2013. Effect of gibberellic acid on growth and yield of muskmelon (*Cucumis melo* L.). In: S.N. Ghosh, S.K. Singh, and J.K. Singh. *Proceedings of the 1st International Conference on Horticulture and Horticulture*. Hore e Bangladesh Agricultural University, Dhaka, Bangladesh, November 2006.

[Goog](#)

37. Meghwal, P.R. 2007. Propagation studies in lehsua (*Cordia myxa*). Indian Journal of Agricultural sciences 77 (11) : 765–767
[Web of Science ®](#) | [Google Scholar](#)
38. Meghwal, P.R., S. Akath, P. Kumar, and B.R. Morwal. 2014. Diversity, distribution and horticultural potential of *Cordia myxa* L.: A promising underutilized fruit species of arid and semi-arid regions of India. Genet. Resour. Crop Evol. 61:1633–1643. doi:10.1007/s10722-014-0161-y.
 | [Web of Science ®](#) | [Google Scholar](#)
39. Mello, A.M., N.A. Streek, E.E. Blankenship, and E.T. Paparozzi. 2009. Gibberellic acid promotes seed germination in *Penstemon digitalis* cv. Husker Red. HortSci 44(3):870–873. doi: 10.21273/HORTSCI.44.3.870.
 | [Web of Science ®](#) | [Google Scholar](#)
40. Meshram, P.C., P.S. Joshi, R.K. Bhoyar, and A.K. Sahoo. 2015. Effect of different plant growth regulators on seedling growth of acid lime (*Citrus aurantifolia* Swingle). Res. Environ. Life Sci. 8(4):725–728.
[Google Scholar](#)
41. Moradi, D.P., F. Sharif-zadeh, and M. Janmohammadi. 2008. Influence of priming techniques on seed germination behavior of maize inbred lines (*Zea mays* L.). J. Agr. Biol. Sci. 3(3):22–25.
[Goog](#)
42. Pal, P., and . . . and
potass
param
Hort
 | [Google Scholar](#)
43. Parab, . . . chemicals on
germi
Int. J. . . .) cv. Solo.

44. Patel, M., R.V. Tank, D.R. Bhanderi, H.M. Patil, V. Patel, and M. Desai. 2018. Response of soaking time and chemicals on germination and growth of tamarind (*Tamarindus indica* L.). *Plant Archives* 18:51–56.

[Google Scholar](#)

45. Patil, A.S., S.R. Patil, and A.S. Kale. 2015. Influence of growth Regulators on germination and growth of endangered medicinal plant *Nothapodytes nimmoniana* j.graham under shade net conditions. *Global J. Res. Med. Plants Indigen. Med.* 4(1):01–09.

[Google Scholar](#)

46. Pradip, K. and Singh H.S. 2010. Tree wealth of the non-forest areas of Gujarat. A report tree census in non-forest areas-2009. Forest department, Gujarat state. p 47

[Google Scholar](#)

47. Ramzan, A., I.A. Hafiz, T. Ahmad, and N.A. Abbasi. 2010. Effect of priming with potassium nitrate and dehusking on seed germination of gladiolus (*Gladiolus alatus*). *Pak. J. Bot.* 42:247–258.

[Web of Science ®](#) | [Google Scholar](#)

48. Schuch, U.K., E. Davison, and J. Kelly. 2001. Seed Propagation of *Cordia boissieri* and *Cordia parvifolia*. Turfgrass and Ornamental Research Report.

<http://ag.arizona.edu/pubs/crops/az1246>

Goog

49. Shaba... seed
germi... J. Agric. Vet.
Sci. 40



50. Shann... cy and
germi... ops
Resea

Goog

51. Sheoran, O.P., D.S. Tonk, L.S. Kaushik, R.C. Hasija, and R.S. Pannu. 1998. Statistical Software Package for Agricultural Research Workers, p. 139–143. In: D.S. Hooda and R.C. Hasija (eds.). Recent advances in information theory, statistics & computer applications. Department of Mathematics Statistics, CCS HAU, Hisar.

[Google Scholar](#)

52. Singh, R. 2018. Activities of research wing of Haryana forest department. Retrieved from http://haryanaforest.gov.in/Portals/0/Research_Activities_Final%20as%20on%2031-3-2018.pdf

[Google Scholar](#)

53. Sirova, J., M. Sedlarova, J. Piterkova, L. Luhova, and M. Petrivalsky. 2011. The role of nitric oxide in the germination of plant seeds and pollen. *Plant Sci.* **181**:560–572. doi:10.1016/j.plantsci.2011.03.014.




[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)


54. Song, J.Q., and H. Fujiyama. 1996. Ameliorative effect of potassium on rice and tomato subjected to sodium salinization. *Soil Sci. Plant Nutr.* **42**:493–501. doi:10.1080/00380768.1996.10416318.



[Web of Science ®](#)

[Google Scholar](#)

55. Tuan, P.A., R. Kumar, P.K. Rehal, P.K. Toora, and B.T. Ayele. 2018. Molecular mechanism of seed dormancy in rice. *Plant Sci.* **271**:1–10. doi:10.1016/j.plantsci.2018.03.014.



[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

56. Varma, R., and H. Fujiyama. 1996. Ameliorative effect of potassium on rice and tomato subjected to sodium salinization. *Soil Sci. Plant Nutr.* **42**:493–501. doi:10.1080/00380768.1996.10416318.



57. Wama, J., and H. Fujiyama. 1996. Ameliorative effect of potassium on rice and tomato subjected to sodium salinization. *Soil Sci. Plant Nutr.* **42**:493–501. doi:10.1080/00380768.1996.10416318.

58. Yücel, E., and G. Yilmaz. 2009. Effects of different alkaline metal salts (Na Cl, KNO3), acid concentrations (H2SO4) and growth regulator (GA3) on the germination of *Salvia cyanescens* Boiss. & Bal. seeds. *Gazi Unive. J. Sci.* 22:123–127.

[Google Scholar](#)

59. Zhang, A., D.F. Huang, and Z. Hou. 2002. Effect of potassium nutrient on development and photosynthesis of melon plant. *J. Shanghai Agricultural Coll.* 20(1):13–17.

[Google Scholar](#)

[Download PDF](#)

Related research

People also read

Recommended articles

Cited by
4



Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Taylor & Francis Group
an informa business

Accessibi

Registered
5 Howick Pl



In this article

