

Free access

5034

CrossRef citations to date


0

Altmetric



Listen


Original Articles


The impact of potential nitrous oxide mitigation strategies on the environmental and economic performance of dairy systems in four New Zealand catchments

C. A. M. De Klein  & R. M. Monaghan

Pages 351-360 | Published online: 16 Feb 2007

 Cite this article  <https://doi.org/10.1080/15693430500402549>

 Reprint




We Care About Your Privacy

We and our 909 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting I Accept enables tracking technologies to support the purposes shown under we and our partners process data to provide. Selecting Reject All or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the Show Purposes link on the bottom of the webpage .Your choices will have effect within our Website. For more details, refer to our Privacy Policy. [Here](#)

We and our partners process data to provide:

Use precise geolocation data. Actively scan device

 I Accept

Reject All

Show Purpose

Abstract

The exposure of the environment and economic performance of New Zealand dairy farm production systems to greenhouse gas emissions, leaching and wintering estimates

feed pads (de Klein et al. [2005](#)). In addition, studies by Di and Cameron ([2002](#), [2003](#)) suggested that the use of a nitrification inhibitor in autumn or spring could be an effective means of reducing N₂O emissions (and nitrate leaching) from animal urine. Other potential N₂O mitigation strategies include diet manipulation to reduce the N content of the diet, increased utilization of effluent N, and improved soil drainage (de Klein & Ledgard [2005](#)).

In 2001, the Best Practice Dairying Catchments project was established to integrate environmentally sustainable practices into dairy farming in New Zealand. This project is carried out in four dairy catchments in New Zealand, two in the North Island, and two in the South Island ([Figure 1](#)), to study farm productivity and catchment-specific environmental issues. Although the initial focus was largely on water-quality issues, estimates of N₂O, CH₄ and carbon dioxide (CO₂) emissions are also made to assess the wider environmental impact of dairy farming in these catchments. The whole-farm system approach of this project enables an evaluation of dairy systems that optimize farm productivity, while minimizing environmental impacts. This paper reports on a desktop assessment of the impact of two of the suggested N₂O mitigation strategies (the use of a wintering pad and of nitrification inhibitors) on the environmental and economic performance of dairy systems in the four catchments.

Figure 1. Location of the four dairy catchments in New Zealand.



×



[Display full size](#)

2 Meth


For each animal p seven to monitori successi descri dairy model (H simulati changes and the OVERSE of nutrie



asture and veys on ality or three ols to ase study mulation duction by ction and om UDDER ne al budgets included

fertilizer use and effluent applications, indirect N₂O emissions from nitrate leaching and ammonia volatilization, and CO₂ emissions associated with fuel and electricity use, processing, and fertilizer use and manufacturing (Wheeler et al. [2003](#)). OVERSEER® estimates the CH₄ and N₂O budgets based on an energy intake model and the IPCC methodology with New Zealand-specific emission factors but also has the ability to assess the impact of on-farm management practices on CH₄ and N₂O emissions (Wheeler et al. [2003](#)). Finally, a purpose-built farm financial model was used to calculate a full farm financial budget based on the current costs of products (milk and meat) and expenses (e.g. imported feed, off-farm grazing) (Ministry of Agriculture and Forestry [2003](#), Dexcel Ltd [2003](#)). Farm profit was expressed as Earnings Before Interest and Tax (EBIT).

Figure 2. Schematic representation of modelling and assessment process.



Display full image

The char... the two... crops du... farm. Th... asse... The imp... of nitrifi... farms w... Table II). can be k... the anim... Table I. In... forage... onto the... the... study farms... and the use... of these... nptions (... the animals... assumed that... were fed

[Download CSV](#)

Display Table

[Download CSV](#)

Display Table



3 Results

The analysis of the four estimates changes from the four estimates (Figure 3). These changes count for the derived.

reductions achieved in the South Island catchments, where off-farm wintering is common practice. Our analysis estimated large N leaching losses from these off-farm wintering areas, which were substantially reduced by keeping the animals on-farm on a wintering pad. The reduction in N leaching had a relatively minor effect on reducing N₂O emissions, because the indirect N₂O emissions from N leaching generally contributed less than 12% of the total emissions. In contrast to the N₂O emissions, total GHG emissions were 1–10% higher than under the current system, with the largest increases occurring in the North Island catchments. The increase in GHG emissions was largely due to an increase in CO₂ emissions associated with fuel use, supplementary feed production, and fertilizer use and manufacturing ([Table III](#)). Methane emissions did not change significantly under the wintering pad option.


Table III. Greenhouse gas emissions (t CO₂ equivalents per farm system per year) from the case-study farm systems and under the wintering pad and nitrification inhibitor mitigation strategies (values in parentheses denote the relative change (%) in total emissions compared with the case-study farm).



[Download CSV](#) [Display Table](#)

Figure 3. Relative changes compared with the case study farms (represented by the dashed line). N leaching losses and N₂O emissions (green bars) and GHG emissions (blue bars) and Earnings (red bars) (bars) and nitrification inhibitor (red bars) based on the annual average.





[Display full size](#)

In terms of financial performance, the wintering pad option had a slightly negative impact on farm profits in three of the catchments, reducing farm EBIT values by 8, 14, and 18% in the Waikakahi, Toenepi, and Waoikura catchments, respectively. These increased costs were largely associated with the increase in imported feed and the capital and operational costs associated with the wintering pad. In the Bog Burn catchment, the wintering pad option did not have any effect on EBIT.

The use of wintering pads for dairy cattle resulted in a small increase in indirect N₂O emissions, with the increase being only 0–9% across the four catchments. This increase was largely due to the increase in methane emissions, which was particularly for the two Southern catchments. The increase in methane emissions was due to the use of wintering pads to utilize methane fermentation. The increase in methane emissions was again due to the use of wintering pads.

4 Discussion

This study suggested that the wintering pad and the nitrification inhibitor strategies could both reduce N_2O emissions and nitrate leaching losses from the case study farms in the four dairy catchments. The reduction in N_2O emission was limited for the wintering pad option, due to the extra feed that was imported to optimize milk production under this management system. However, for the nitrification inhibitor strategy, N_2O emissions were substantially reduced. The reduction in N leaching was slightly less. Our assessment assumed that N leaching from grazed pastures would be reduced by 20%, which is lower than the 60% presented by Di and Cameron ([2002](#)). However, their estimate was based on lysimeter studies of urine patches and is thus likely to represent a maximum potential benefit. Recent field measurements suggest that a 20% reduction better represents the effect of nitrification inhibitors on N leaching losses from grazed pastures (C Smith, unpublished data).

Our analysis further showed that the use of nitrification inhibitors had a limited effect on total GHG emissions reduction, compared with the reduction in N_2O emissions, due to an increase in both CH_4 and CO_2 emissions from the farm systems. Although nitrification inhibitors do not directly affect emissions of these GHGs, their use was estimated to result in an increase in pasture production and thus milk production, which, in turn, increased CH_4 and CO_2 emissions. It should be noted, however, that we assumed that the use of nitrification inhibitors resulted in an increase in stock numbers to utilize the extra pasture production under the inhibitor option. An alternative

scenario was to increase the stocking rate under the wintering pad and/or nitrification inhibitor scenario, which would result in a further reduction in total GHG emissions. However, this alternative scenario was not included in the analysis as it was not a direct result of the nitrification inhibitor strategy. The advantage of the nitrification inhibitor strategy was that it did not require the import of extra feed.

Further research is needed to assess the impact of nitrification inhibitors on the environment. This way, the environmental impact of nitrification inhibitors can be assessed. The environmental impact of nitrification inhibitors can be assessed by comparing the environmental impact of nitrification inhibitors with the environmental impact of the wintering pad and the nitrification inhibitor strategies. This way, the environmental impact of nitrification inhibitors can be assessed by comparing the environmental impact of nitrification inhibitors with the environmental impact of the wintering pad and the nitrification inhibitor strategies.



the reduction in total GHG emissions per unit of product was larger than the reduction in total GHG emissions from the farm systems (c. 15% compared with a maximum of 6%; [Table III](#)). This indicated that nitrification inhibitors are likely to increase a farm's efficiency as the increase in milk production was larger than any increase in total GHG emissions from the farm. An assessment of environmental losses 'per unit of product' can thus identify management strategies that have the largest reduction in environmental emissions for a given production level.

The economic analyses indicated that the wintering pad option generally reduced the total farm earnings due to increased costs associated with imported feed and capital and operating costs of the wintering pad. The exception was for the case-study farm in the Bog Burn catchment, where the wintering pad option did not reduce farm financial performance. However, our calculations did not account for any potential benefits of reduced soil physical damage from grazing in wet winter conditions, which can decrease spring pasture and animal production, and increase N₂O emissions (de Klein et al. [2005](#)). In addition, the economic analysis did not include an assessment of potential savings (or costs) associated with a reduction (or increase) in GHG emissions or N leaching losses. In New Zealand, the value of GHG emissions is currently set at \$25 per tonne of CO₂-equivalent. However, since GHG emissions under the wintering pad option increased in all catchments, this would result in a further reduction in farm earnings. On the other hand, N leaching losses decreased by up to 44%, and potential savings from reduced N leaching losses could offset some of the increased GHG emissions and associated costs.

The economic analysis also indicated that the wintering pad option appeared to be a cost-effective way to reduce GHG emissions and N leaching losses, and enhance farm financial performance. However, the economic analysis did not account for the potential benefits of reduced soil physical damage from grazing in wet winter conditions, which could further reduce GHG emissions and N leaching losses, and increase pasture production. The economic analysis also did not account for the potential benefits of reduced soil physical damage from grazing in wet winter conditions, which could further reduce GHG emissions and N leaching losses, and increase pasture production. The economic analysis also did not account for the potential benefits of reduced soil physical damage from grazing in wet winter conditions, which could further reduce GHG emissions and N leaching losses, and increase pasture production.



5 Conclusions



The ong edged. We

Sustainable Farming Fund, and the New Zealand Fertilizer Manufacturers Research Association, for their financial support of this work, and Duncan Smeaton, Mark Hyslop, and David Stevens, for expert assistance with UDDER modelling.

References

1. de Klein , C AM , Barton , L , Sherlock , R R , Li , Z and Littlejohn , R P . 2003 . Estimating a nitrous oxide emission factor for animal urine from some New Zealand pastoral soils . Australian Journal of Soil Research , 41 : 381 – 399 .
[Web of Science ®](#) | [Google Scholar](#)
2. de Klein , C AM and Ledgard , S F . 2005 . Nitrous oxide emissions from New Zealand agriculture—key sources and mitigation strategies . Nutrient Cycling in Agroecosystems , 72 : 77 – 85 .
[Web of Science ®](#) | [Google Scholar](#)
3. de Klein , C AM , Smith , L C and Monaghan , R M . 2005 . Restricted autumn grazing to red New Zealand. [Agriculture](#) [Google Scholar](#)
4. Dexe [Google Scholar](#) 2002 ,
Hamil [Google Scholar](#)
5. Di , H [Google Scholar](#) andiamide (DCD) [Google Scholar](#) related grazed [Google Scholar](#) 03 .



6. Di , H J and Cameron , K C . 2003 . Mitigation of nitrous oxide emissions in spray-irrigated grazed grassland by treating the soil with dicyandiamide, a nitrification inhibitor . Soil Use and Management , 19 : 284 – 290 .

[Web of Science ®](#) | [Google Scholar](#)

7. Hart , R PS , Larcombe , M T , Sherlock , R A and Smith , L A . 1998 . Optimisation techniques for a computer simulation of a pastoral dairy farm . Computers and Electronics in Agriculture , 19 : 129 – 153 .

[Web of Science ®](#) | [Google Scholar](#)

8. Larcombe , M T . 1999 . UDDER for Windows: A desktop dairyfarm for extension and research—operating manual , Maffra , , Australia : Maffra Herd Improvement Co-op .

[Google Scholar](#)

9. Ledgard , S F and Menneer , J C . 2005 . “ Nitrate leaching in grazing systems and management strategies to reduce losses ” . In Developments in fertiliser application technologies and nutrient management , Edited by: Currie , L D and Hanly , J A . 79 – 92 . Palmerston North , , New Zealand : Fertiliser and Lime Research Centre, Massey University .

[Google Scholar](#)

10. Ministry of Agriculture and Forestry . 2011 . Downloaded on 11/11/2011 11:11:11 AM

November 2011

<http://www.maf.govt.nz/monitoring>

[Google Scholar](#)

11. New Zealand . 2011 . Greenhouse gas inventory report supporting the 2009 Climate Change Act . Office of the Environment .

[Google Scholar](#)



2. New Zealand Climate Change Office . 2005b . Report on revised projections for the Kyoto Protocol—first commitment period. 30 April 2004 , Wellington , , New Zealand : New Zealand Climate Change Office .

[Google Scholar](#)

3. Schils , R LM , Verhagen , A , Aarts , H FM and Sebek , L BJ . 2005 . A farm level approach to define successful mitigation strategies for GHG emissions from ruminant livestock systems . *Nutr Cycl Agroecosyst* , 71 : 163 – 175 .

[Web of Science ®](#)

[Google Scholar](#)

4. Wheeler , D M , Ledgard , S F , de Klein , C AM , Monaghan , R M , Carey , P L , McDowell , R W and Johns , K L . 2003 . OVERSEER® nutrient budgets 2—moving towards on-farm resource accounting . *Proc New Zealand Grassland Assoc* , 65 : 191 – 194 .

[Google Scholar](#)

[Download PDF](#)

Related



Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright

Accessib

Registered
5 Howick Pl

or & Francis Group
orma business

