


[International Journal of Coal Preparation and Utilization](#) >Volume 42, 2022 - [Issue 4](#)415 | 6 | 0
Views | CrossRef citations to date | Altmetric

Research Article

Characterization of rare earth elements by XRT sorting products of a South African coal seam

G. Akdogan , S. Bradshaw, C. Dorfling, C. Bergmann, T. Ghosh  & Q. Campbell

Pages 1071-1087 | Received 25 Sep 2019, Accepted 23 Oct 2019, Published online: 11 Nov 2019

 Cite this article  <https://doi.org/10.1080/19392699.2019.1685506>

Sample our
Engineering & Technology
Journals
>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

 Full Article Figures & data References Citations Metrics Reprints & Permissions[Read this article](#)[Share](#)

ABSTRACT

South Africa is an important participant in the global coal market. The four Upper coal seam situated in Springs-Witbank Coalfield was subjected to XRT sorting and subsequent mineralogical studies to characterize REE content. The average total REE content (LREE and HREE) of the coal were 280 ppm on ash basis. The surrounding elemental composition besides carbon primarily consisted of O, Si, S and Fe with minor quantities of Al, P and Ti. Major minerals, such as quartz, dolomite, illite, gypsum, siderite, pyrite correlated positively with ash content ($r > 0.9$) except kaolinite, and microcline, indicating incongruent source and formation mechanisms. Additionally, the HREEs Lu, Tm, Dy, Ho, Gd and Tb displayed strong correlations ($r > 0.95$) with ash content but were negatively associated with fixed carbon. Sc is negatively associated with coal ash. Fixed carbon is weakly associated with La but negatively associated with

HREEs. La showed a weak association with Ce. Y is strongly correlated with HREE elements especially with Dy, Tm, Lu and Ho while at the same time LREE are strongly associated with Ce, Pr, Nd, La, Eu but negatively with Sc and fixed carbon levels.

KEYWORDS:

Coal

REE

XRT sorting

South Africa

Acknowledgments

This work was partly supported by SAMMRI (South African Minerals to Metals Research Institute) with a seed fund under S1705.

Additional information

Funding

This work was partly supported by the SAMMRI (South African Minerals to Metals Research Institute) with a seed funding under S1705.

Related Research Data

[Estimations of Clarkes for Carbonaceous biolithes: World averages for trace element contents in black shales and coals](#)

Source: International Journal of Coal Geology

[Mineralogical and geochemical compositions of the coal in the Guanbanwusu Mine, Inner Mongolia, China: Further evidence for the existence of an Al \(Ga and REE\) ore deposit in the Jungar Coalfield](#)

Source: International Journal of Coal Geology

[Geochemistry of rare earth elements in a marine influenced coal and its organic solvent extracts from the Antaibao mining district, Shanxi, China](#)

Source: International Journal of Coal Geology

[A high-pyrite semianthracite of Late Permian age in the Songzao Coalfield, southwestern China: Mineralogical and geochemical relations with underlying mafic](#)

tuffs

Source: International Journal of Coal Geology

Related research

People also read

Recommended articles

Cited by
6

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

 Taylor and Francis
Group

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG