







Q

Food Additives & Contaminants: Part A >

Volume 34, 2017 - Issue 2

▶ Volume 34, Issue 2 ▶ Urinary signature of pig carcasses with

217 2

Views CrossRef citations to date Altmetric

Original Articles


Urinary signature of pig carcasses with boar taint by liquid chromatography-highresolution mass spectrometry

Cristina C. Jacob ✓, Gaud Dervilly-Pinel ✓, Yoann Deceuninck, Audrey Gicquiau, Patrick Chevillon, Michel Bonneau & ...show all

Pages 218-227 | Received 04 Oct 2016, Accepted 22 Nov 2016, Published online: 14 Dec 2016

66 Cite this article

A https://doi.org/10.1080/19440049.2016.1265152



Sample our Bioscience journals, sign in here to start your access, Latest two full volumes FREE to you for 14 days

Full Article

Figures & data

References

Supplemental

66 Citations

Metrics

A Reprin

ABSTE

Boar tai

is identif that boa

compou

spectr

androste

Discrimi of multiv

highligh

skatole

urinary

levels in

We Care About Your Privacy

We and our 907 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting "I Accept" enables tracking technologies to support the purposes shown under "we and our partners process data to provide," whereas selecting "Reject All" or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the ["privacy preferences"] link on the bottom of the webpage [or the floating icon on the bottom-left of the webpage, if applicable]. Your choices will have effect within our Website. For more details, refer to our Privacy Policy. Here

We and our partners process data to provide:

I Accept

Reject All

roducts and

widely held Show Purpose

us

ss a mass

ofile of urine

vels of

SI(+)-HRMS.

application

cers to be

tenone and

ent of these

and skatole


GRAPHICAL ABSTRACT BOAR TAINT LC-HRMS URINARY FINGERPRINTING

KEYWORDS:

Metabolomics	boar taint	urinary profile	biomarkers	LC-HRMS

Disclosure statement

No potential conflict of interest was reported by the authors.



Related Research Data

Urinary signature of pig carcasses with boar taint by liquid chromatography-high-resolution mass spectrometry

Source: figshare

Dissemination of metabolomics results: role of MetaboLights and COSMOS

Source: GigaScience

The development of pain in young pigs associated with castration and attempts to prevent castration-induced behavioral changes

Source: Journal of Animal Science

Behavioural responses of piglets to castration: the effect of piglet age

Source: Applied Animal Behaviour Science

Introduction to multivariate calibration in analytical chemistry

Source: The Analyst

Sensory evaluation of boar loins: Trained assessors' olfactory acuity affects the perception of boar taint compounds

Source: Meat Science

An international study on the importance of androstenone and skatole for boar taint: I. Presentation of the programme and measurement of boar taint compounds with different analytical procedures



Visualization of GC/TOF-MS-Based Metabolomics Data for Identification of Biochemically Interesting Compounds Using OPLS Class Models

Source: Analytical Chemistry

Role of 4-Phenyl-3-buten-2-one in Boar Taint: Identification of New Compounds Related

to Sensorial Descriptors in Pig Fat

Source: Journal of Agricultural and Food Chemistry Basics of mass spectrometry based metabolomics

Source: PROTEOMICS

Learning to smell: Repeated exposure increases sensitivity to androstenone, a major

component of boar taint

Source: Meat Science

The effects of method of castration, rearing condition and diet on sensory quality of pork assessed by a trained panel

Source: Meat Science

Boar Taint is Related to Endocrine and Anatomical Changes at Puberty but not to Aggressive Behaviour in Entire Male Pigs

Source: Reproduction in Domestic Animals

Consumer perception of boar meat as affected by labelling information, malodorous compounds and sensitivity to androstenone

Source: Meat Science

Androgen Steroids as an Aid to the Detection of Oestrus in Pig Artificial Insemination



of ultra-high performance liquid chromatography coupled to high resolution mass spectrometry

Source: Food Chemistry

Assessment of PLSDA cross validation

Source: Metabolomics

OPLS discriminant analysis: combining the strengths of PLS-DA and SIMCA

classification

Source: Journal of Chemometrics

XCMS: Processing Mass Spectrometry Data for Metabolite Profiling Using Nonlinear

Peak Alignment, Matching, and Identification

Source: Analytical Chemistry

Performances, meat quality and boar taint of castrates and entire male pigs fed a

standard and a raw potato starch-enriched diet

Source: Animal

LC-MS-based metabonomics analysis Source: Journal of Chromatography B

 5α -androst-16-ene-3-one:—Compound responsible for taint in boar fat

Source: Journal of the Science of Food and Agriculture

Boar taint detection: A comparison of three sensory protocols

Source: Meat Science

Linking provided by Schole plorer



Information for Open access **Authors** Overview R&D professionals Open journals Editors **Open Select** Librarians **Dove Medical Press** Societies F1000Research Opportunities Help and information Reprints and e-prints Advertising solutions Newsroom Accelerated publication Corporate access solutions Books Keep up to date Register to receive personalised research and resources by email Sign me up X or & Francis Group Copyright