

Free access

1,208 Views | 8 CrossRef citations to date | 0 Altmetric

Listen

Author's Views

SESTRINs regulate mTORC1 via RRAGs: The riddle of GATOR

Andrei V Budanov

Article: e997113 | Received 04 Dec 2014, Accepted 08 Dec 2014, Published online: 06 May 2015

Cite this article <https://doi.org/10.1080/23723556.2014.997113>

Check for updates

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

View PDF

Abstract

SESTRINs, proteins encoded by the SESN1-3 genes in mammals, are well-established suppressors of the mechanistic target of rapamycin complex 1 (mTORC1) kinase.

Recently, we have shown that RRAGs regulate mTORC1 activity in this pathway.

We Care About Your Privacy

We and our 848 partners store and/or access information on a device, such as unique IDs in cookies to process personal data. You may accept or manage your choices by clicking below, including your right to object where legitimate interest is used, or at any time in the privacy policy page. These choices will be signaled to our partners and will not affect browsing data. [Privacy Policy](#)

We and our partners process data to provide:

Use precise geolocation data. Actively scan device characteristics for identification. Store and/or access information on a device. Personalised advertising and content, advertising and content measurement, audience research and services development.

List of Partners (vendors)

I Accept

Essential Only

Show Purpose



Key words

SESTRINs, mammalian target of rapamycin (mTORC1), growth

is a... support the... of GATOR2... 3 in... complex 1... growth

factors, and stress to control metabolism, autophagy, and cell growth. mTORC1 activity inversely correlates with the lifespan of most eukaryotes and mTORC1 contributes to several different pathologies including cancer, diabetes, and neurodegenerative diseases.² mTORC1 is activated by 2 groups of small guanine triphosphatases (GTPases): Ras homolog enriched in brain (RHEB) and members of the Ras-related GTP-binding protein (RRAG) family, working as RRAG-A/B and RRAG-C/D heterodimers. RHEB activity is inhibited by the tuberous sclerosis protein complex (TSC). TSC in turn is controlled by several stress insults via activation of 5'-AMP-activated protein kinase (AMPK), and we have shown previously that SESTRINs inhibit mTORC1 via the AMPK-TSC axis.³ Amino acids and glucose control mTORC1 via RRAGs, which stimulate translocation of mTORC1 to the lysosomes where it can be activated by RHEB. RRAG-A/B is activated by its GTPase exchange factor (GEF) called ragulator and is inhibited by the protein complex GATOR1 (GTPase activating protein [GAP] activity toward RRAG complex 1) working as a GAP. GATOR1 in turn is suppressed by the GATOR2 protein complex within the GATOR supercomplex.⁴

Three groups recently reported a novel mechanism of mTORC1 inhibition by SESTRINs via suppression of RRAG-dependent mTORC1 lysosomal translocation.⁵⁻⁷ SESTRINs are critical for mTORC1 inhibition by amino acid withdrawal in mammalian cells. Surprisingly, SESTRINs do not affect RRAG-A/B GDP/GTP loading, suggesting that they are not regulators of RRAG-A/B GAP or GEF. Although all 3 groups agreed on the importance of RRAGs for mTORC1 suppression by SESTRINs, they proposed different mechanisms for this regulatory activity.

Using m... complexes,
our grou... time,
Sabatini... ng similarity
to our da... a GATOR-
and RRA... ATOR2
interacti... way from
GATC...
Li's grou... INs.⁶
Accordin... rk as RRAG-
A/B guar... shares
homolog... GTPase




family members. Mutations in the GDI motif significantly impair the ability of SESTRIN2 to inhibit mTORC1 and delivery of a peptide GDI motif is sufficient to inhibit mTORC1.⁶

There are several major concerns that remain unexplained. First, do SESTRINs directly interact with RRAGs and what is the role of the GATOR complex in the regulation of RRAG-A/B and mTORC1 activities by SESTRINs? Despite the high abundance of GATOR2 peptides among our mass spectrometry data, we did not observe any RRAG peptides, indicating that this interaction is very weak and might be indirect. Second, according to Li's model, SESTRIN2 must have some regulatory role over RRAGs even in the absence of GATOR1 as a result of intrinsic RRAG-A/B GTPase activity,⁶ although data from our group and Sabatini's group indicate a strong requirement for the GATOR complex in the regulation of mTORC1 by SESTRINs.^{5,7} Finally, in contrast to Peng et al., we were not able to detect any colocalization of SESTRIN2 with lysosomes, indicating that SESTRINs might play a major role in the regulation of GATOR and RRAGs in the cytoplasm.^{6,7} To reconcile these studies, we propose that GATOR is required for the interaction between SESTRIN2 and RRAGs, potentially providing a bridge between them and somehow bringing the potential GDI domain of SESTRINs into close proximity of the RRAG-A/B protein. Thus, GATOR might control interactions between SESTRINs and RRAGs and assist in mTORC1 inhibition by SESTRINs (Fig. 1A). Alternatively, SESTRINs might interact solely with GATOR2, and the GDI activity of SESTRIN toward RRAGA/B might be mediated by GATOR (Fig. 1B). We demonstrated that the interaction between SESTRIN2 and GATOR2 is mediated by WD-repeat domain 24 (WDR24) and SEH1-like (SEH1L) proteins, thus it would be important to examine whether the GDI domain of SESTRIN is required for the interaction between SESTRINs and GATOR2.

Figure 1. GATOR and RRAG-A/B interaction and the mTORC1 regulation of RRAG-A/B activity. RRAG-A/B activity is regulated by GATOR2 weakly with RRAG-A/B interactions. SESTRINs on RRAG-A/B activity is mediated by GATOR2. RRAG-A/B activity is mediated by GATOR2.






Display full size

To reconcile the discrepancy in localization studies we propose that the interaction between SESTRINs and GATOR-RRAGs complexes is a dynamic process and under certain conditions SESTRINs can be associated with lysosomes, although most of the proteins associated with GATOR are still located in the cytoplasm in close proximity to lysosomes. Our findings are consistent with recently published data from Avruch's laboratory⁸ showing a predominantly cytoplasmic localization of RRAGs based on protein fractionation studies. In the future it would be important to analyze the localization of endogenous SESTRINs and RRAGs, and the role of SESTRINs in the control of RRAG localization in the cell under normal and stress conditions.

Interestingly, yeast mTORC1 is regulated by amino acids via a mechanism mediated by Gtr1 and Gtr2, orthologs of mammalian RRAG-A/B and RRAG-C/D, and the SEA complex (SEAC), an analog of mammalian GATOR. SEAC functions as a whole complex, suggesting that GATOR might also work as an entire complex to regulate RRAG-A/B activity.^{9,10} However, the fact that yeasts do not express sestrins indicates that they are






Disclo

No potential conflicts of interest were disclosed.

Funding

This work was supported by NIH CA172660 to AB.

References

1. Budanov AV, Lee JH, Karin M. Stressin' Sestrins take an aging fight. *EMBO Mol Med* 2010; **2**:388-400; PMID:20878915; <http://dx.doi.org/10.1002/emmm.201000097>
 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
2. Laplante M, Sabatini DM. mTOR signaling in growth control and disease. *Cell* 2012; **149**:274-93; PMID:22500797; <http://dx.doi.org/10.1016/j.cell.2012.03.017>
 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
3. Budanov AV, Karin M. p53 target genes sestrin1 and sestrin2 connect genotoxic stress and mTOR signaling. *Cell* 2008; **134**:451-60; PMID:18692468; <http://dx.doi.org/10.1016/j.cell.2008.06.028>
 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

4. Bar-Pet... *Cell Biol*
2014;

5. Ch... L, Spooner
E, Isas... negatively
Regul... 2014;
9(1):1



6. Peng M, Yin N, Li MO. Sestrins Function as Guanine Nucleotide Dissociation Inhibitors for Rag GTPases to Control mTORC1 Signaling. Cell 2014; 159:122-33; PMID:25259925; <http://dx.doi.org/10.1016/j.cell.2014.08.038>

 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

7. Parmigiani A, Nourbakhsh A, Ding B, Wang W, Kim YC, Akopiants K, Guan KL, Karin M, Budanov AV. Sestrins Inhibit mTORC1 Kinase Activation through the GATOR Complex. Cell Rep 2014; 9:1281-91; PMID:25457612; <http://dx.doi.org/10.1016/j.celrep.2014.10.019>

 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

8. Oshiro N, Rapley J, Avruch J. Amino acids activate mammalian target of rapamycin (mTOR) complex 1 without changing Rag GTPase guanyl nucleotide charging. J Biol Chem 2014; 289:2658-74; PMID:24337580; <http://dx.doi.org/10.1074/jbc.M113.528505>

 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

9. Dokudovskaya S, Waharte F, Schlessinger A, Pieper U, Devos DP, Cristea IM, Williams R, Salamero J, Chait BT, Sali A, et al. A conserved coatomer-related complex containing Sec13 and Seh1 dynamically associates with the vacuole in *Saccharomyces cerevisiae*. Mol Cell Proteomics 2011; 10:M110 006478; <http://dx.doi.org/10.1074/mcp.M110.006478>

 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

10. Panch... TORC1
activa... 2013;
12:29...



Download

Related

Information for


- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Copyright

Accessib

Registered
5 Howick Pl

or & Francis Group
orma business

