







Q

Home ▶ All Journals ▶ Bioscience ▶ Journal of Environmental Science and Health, Part A ▶ List of Issues ▶ Volume 38, Issue 11 ▶ Concentration of Copper, Iron, Zinc, Cad

Journal of Environmental Science and Health, Part A > Toxic/Hazardous Substances and Environmental Engineering Volume 38, 2003 - Issue 11

160 24
Views CrossRef citations to date Altmetric
Original Articles

Concentration of Copper, Iron, Zinc, Cadmium, Lead, and Nickel in Boar Semen and Relation to the Spermatozoa Quality

Peter Massányi, Jozef Trandžík, Pavol Nad, Beáta Koréneková, Magdaléna Skalická, Robert Toman , ...show all

Pages 2643-2651 | Received 28 Jan 2003, Published online: 06 Feb 2007

Sample our
Engineering & Technology
Journals
>> Sign in here to start your access to the latest two volumes for 14 days

Full A

A Repri

Abstra

The cond

to spern

abso 1.64

boar ser

cadmiun analysis

 0.02 ± 0

total per

determi

We Care About Your Privacy

We and our 880 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting I Accept enables tracking technologies to support the purposes shown under we and our partners process data to provide. Selecting Reject All or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the Show Purposes link on the bottom of the webpage .Your choices will have effect within our Website. For more details, refer to our Privacy Policy. Here

We and our partners process data to provide:

Use precise geolocation data. Actively scan device

I Accept

Reject All

Show Purpose

d by atomic emen was

s its relation

of zinc in

evel of

 $^{-1}$. The

en was

 y^{-1} . The

llysis


88%

flagellum torso, 0.85% flagellum ball, 0.42% broken flagellum, 0.23% retention of the cytoplasmic drop, 0.14% small heads, 0.03% large heads, and 1.83% forms other of pathological changes. Correlation analysis showed significant (p<0.05) positive correlation between copper and lead (r = 0.52). High correlation between small head and knob twisted tail (r = 0.67), small head and broken flagellum (r = 0.88) as well as between small head and total number of pathological spermatozoa (r = 0.73) was determined.



Acknowledgments

We would like to express our gratitude to Ing. P. Cupka and Ing. O. Messinger for statistical analysis, Mrs. P. Sýkorová and Mrs. A. Sobčáková for technical assistance. This was supported by VEGA Scientific Grant 1/9080/02 from the Ministry of Education of Slovak Republic.



Information for Open access Authors Overview R&D professionals Open journals Editors **Open Select** Librarians **Dove Medical Press** Societies F1000Research Opportunities Help and information Reprints and e-prints Advertising solutions Newsroom Accelerated publication Corporate access solutions Books Keep up to date Register to receive personalised research and resources by email Sign me up X or & Francis Group Copyright