





▶ All Journals ▶ Journal of Environmental Science and Health, Part A ▶ List of Issues ▶ Volume 38, Issue 11 ▶ Concentration of Copper, Iron, Zinc, Cad

Journal of Environmental Science and Health, Part A > Toxic/Hazardous Substances and Environmental Engineering Volume 38, 2003 - Issue 11

154 23

Views CrossRef citations to date Altmetric

Original Articles

Concentration of Copper, Iron, Zinc, Cadmium, Lead, and Nickel in Boar Semen and Relation to the Spermatozoa Quality

Peter Massányi, Jozef Trandžík, Pavol Nad, Beáta Koréneková, Magdaléna Skalická, Robert Toman , ... show all

Pages 2643-2651 | Received 28 Jan 2003, Published online: 06 Feb 2007

Sample our Engineering & Technology to the latest two volumes for 14 days

Full Article

Figures & data

References

66 Citations

Metrics

Reprints & Permissions

Read this article

Abstra

The con

to spern

absorpti

 1.64 ± 0

boar 8 cadn

analysis 0.02 ± 0

total per

determi

flagellur

We Care About Your Privacy

We and our 842 partners store and/or access information on a device, such as unique IDs in cookies to process personal data. You may accept or manage your choices by clicking below, including your right to object where legitimate interest is used, or at any time in the privacy policy page. These choices will be signaled to our partners and will not affect browsing data. Privacy Policy

We and our partners process data to provide:

Use precise geolocation data. Actively scan device characteristics for identification. Store and/or access information on a device. Personalised advertising and content, advertising and content measurement, audience research and services development.

List of Partners (vendors)


I Accept s its relation d by atomic **Essential Onl** emen was of zinc in Show Purpose evel of $^{-1}$. The en was $^{-1}$. The lysis 88% tion of the

cytoplasmic drop, 0.14% small heads, 0.03% large heads, and 1.83% forms other of pathological changes. Correlation analysis showed significant (p<0.05) positive correlation between copper and lead (r=0.52). High correlation between small head and knob twisted tail (r=0.67), small head and broken flagellum (r=0.88) as well as between small head and total number of pathological spermatozoa (r=0.73) was determined.

Q Keywords: Copper Iron Zinc Cadmium Lead Nickel Spermatozoa Boar

Acknowledgments

We would like to express our gratitude to Ing. P. Cupka and Ing. O. Messinger for statistical analysis, Mrs. P. Sýkorová and Mrs. A. Sobčáková for technical assistance. This was supported by VEGA Scientific Grant 1/9080/02 from the Ministry of Education of Slovak Republic.



Information for Open access **Authors** Overview R&D professionals Open journals Editors **Open Select** Librarians **Dove Medical Press** Societies F1000Research Opportunities Help and information Reprints and e-prints Advertising solutions Newsroom Accelerated publication Corporate access solutions Books Keep up to date Register to receive personalised research and resources by email Sign me up Taylor & Francis Group Copyright © 2024 Informa UK Limited Privacy policy Cookies Terms & conditions Accessib X

