







Home ► All Journals ► Food Science & Technology ► Food Reviews International ► List of Issues Volume 20, Issue 1 ➤ Omega-6/Omega-3 Essential Fatty Acid Rat ....

#### Food Reviews International >

Volume 20, 2004 - <u>Issue 1</u>

2,986 354 18 Views CrossRef citations to date Altmetric Original Articles

# Omega-6/Omega-3 Essential Fatty Acid Ratio and Chronic Diseases

Artemis P. Simopoulos M.D.

Pages 77-90 | Published online: 16 Aug 2006

**66** Cite this article https://doi.org/10.1081/FRI-120028831



Full Article

Figures & data

References

**66** Citations

**Metrics** 

Reprints & Permissions

Read this article

Share

## **Abstract**

Several sources of information suggest that human beings evolved on a diet with a ratio of omega-6 to omega-3 essential fatty acids (EFA) of  $\sim 1$  whereas in Western diets the ratio is 15/1–16.7/1. Western diets are deficient in omega-3 fatty acids, and have excessive amounts of omega-6 fatty acids compared with the diet on which human beings evolved and their genetic patterns were established. Excessive amounts of omega-6 polyunsaturated fatty acids (PUFA) and a very high omega-6/omega-3 ratio, as is found in today's Western diets, promote the pathogenesis of many diseases, including cardiovascular disease, cancer, and inflammatory and autoimmune diseases, whereas increased levels of omega-3 PUFA (a lower omega-6/omega-3 ratio), exert suppressive effects. In the secondary prevention of cardiovascular disease, a ratio of 4/1 was associated with a 70% decrease in total mortality. A ratio of 2.5/1 reduced

rectal cell proliferation in patients with colorectal cancer, whereas a ratio of 4/1 with the same amount of omega-3 PUFA had no effect. The lower omega-6/omega-3 ratio in women with breast cancer was associated with decreased risk. A ratio of 2-3/1 suppressed inflammation in patients with rheumatoid arthritis, and a ratio of 5/1 had a beneficial effect on patients with asthma, whereas a ratio of 10/1 had adverse consequences. These studies indicate that the optimal ratio may vary with the disease under consideration. This is consistent with the fact that chronic diseases are multigenic and multifactorial. Therefore, it is quite possible that the therapeutic dose of omega-3 fatty acids will depend on the degree of severity of disease resulting from the genetic predisposition. A lower ratio of omega-6/omega-3 fatty acids is more desirable in reducing the risk of many of the chronic diseases of high prevalence in Western societies, as well as in the developing countries, that are being exported to the rest of the world

### Keywords:

Omega-6/omega-3 fatty acid balance genetic patterns chronic disease dietary recommendations

Related research •

People also read

Recommended articles

Cited by 354

Information for

**Authors** 

**R&D** professionals

**Editors** 

Librarians

**Societies** 

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

**Open Select** 

**Dove Medical Press** 

F1000Research

Help and information

Help and contact

Newsroom

All journals

**Books** 

#### Keep up to date

Register to receive personalised research and resources by email



Sign me up











Accessibility



Copyright © 2025 Informa UK Limited Privacy policy Cookies Terms & conditions



Registered in England & Wales No. 01072954 5 Howick Place | London | SW1P 1WG