



Food Reviews International >

Volume 20, 2004 - [Issue 1](#)

2,932

Views

349

CrossRef citations to date

18

Altmetric

Original Articles

# Omega-6/Omega-3 Essential Fatty Acid Ratio and Chronic Diseases

Artemis P. Simopoulos M.D.

Pages 77-90 | Published online: 16 Aug 2006

Cite this article <https://doi.org/10.1081/FRI-120028831>

Sample our  
Food Science & Technology  
Journals  
>> [Sign in here](#) to start your access  
to the latest two volumes for 14 days

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

Read this article

Share

## Abstract

Several sources of information suggest that human beings evolved on a diet with a ratio of omega-6 to omega-3 essential fatty acids (EFA) of ~1 whereas in Western diets the ratio is 15/1–16.7/1. Western diets are deficient in omega-3 fatty acids, and have excessive amounts of omega-6 fatty acids compared with the diet on which human beings evolved and their genetic patterns were established. Excessive amounts of omega-6 polyunsaturated fatty acids (PUFA) and a very high omega-6/omega-3 ratio, as is found in today's Western diets, promote the pathogenesis of many diseases, including cardiovascular disease, cancer, and inflammatory and autoimmune diseases, whereas increased levels of omega-3 PUFA (a lower omega-6/omega-3 ratio), exert suppressive effects. In the secondary prevention of cardiovascular disease, a ratio of 4/1 was associated with a 70% decrease in total mortality. A ratio of 2.5/1 reduced rectal cell proliferation in patients with colorectal cancer, whereas a ratio of 4/1 with the same amount of omega-3 PUFA had no effect. The lower omega-6/omega-3 ratio in

women with breast cancer was associated with decreased risk. A ratio of 2-3/1 suppressed inflammation in patients with rheumatoid arthritis, and a ratio of 5/1 had a beneficial effect on patients with asthma, whereas a ratio of 10/1 had adverse consequences. These studies indicate that the optimal ratio may vary with the disease under consideration. This is consistent with the fact that chronic diseases are multigenic and multifactorial. Therefore, it is quite possible that the therapeutic dose of omega-3 fatty acids will depend on the degree of severity of disease resulting from the genetic predisposition. A lower ratio of omega-6/omega-3 fatty acids is more desirable in reducing the risk of many of the chronic diseases of high prevalence in Western societies, as well as in the developing countries, that are being exported to the rest of the world

Keywords:

- Omega-6/omega-3 fatty acid balance
- genetic patterns
- chronic disease
- dietary recommendations



## Related research

People also read

Recommended articles

Cited by  
349

## Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

## Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

## Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

## Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

## Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2025 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)

 Taylor and Francis Group

Registered in England & Wales No. 01072954  
5 Howick Place | London | SW1P 1WG