



Journal of Asthma >

Volume 41, 2004 - [Issue 3](#)

528 | 89

Views | CrossRef citations to date | 4

Altmetric

Original Article

Ratio of Omega-6 to Omega-3 Fatty Acids and Childhood Asthma

W. H. Oddy, Ph.D. , N. H. de Klerk, Ph.D. , G. E. Kendall, M.P.H. , S. Mihrshahi, M.P.H. & J. K. Peat, Ph.D.

Pages 319-326 | Published online: 26 Aug 2009

🗨️ Cite this article 🔗 <https://doi.org/10.1081/JAS-120026089>

Sample our
Medicine, Dentistry, Nursing
& Allied Health Journals
>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

📄 Full Article

🖼️ Figures & data

📖 References

🗨️ Citations

📊 Metrics

📄 Reprints & Permissions

Read this article

🔗 Share

Abstract

Asthma is a leading cause of morbidity for children and is a major public health problem in Australia. Ecological and temporal data suggest that dietary factors may have a role in recent increases in the prevalence of asthma. Aim: The aim of conducting this study was to investigate whether childhood asthma was associated with the ratio of omega 6 (n-6) to omega 3 (n-3) fatty acids in the diet (n-6:n-3). Method: The Western Australian Pregnancy Cohort Study is a prospective birth cohort of 2602 children. Using a nested case-control cross-sectional study design within this cohort, a group of children were identified as cases with current asthma at 6 or at 8 years of age or as controls with no asthma at 6 or at 8 years. Dietary details including n-6 and n-3 fatty acid intake data were collected by parent response to a questionnaire when the children were 8 years old. Logistical regression was used to compare quartiles of n-6:n-3 intake in cases and controls. Adjustment was made for covariates: gender, gestational age, breastfeeding,

older siblings, maternal smoking during pregnancy, maternal age, maternal asthma, child's current age in months, body mass index, total energy intake, and antioxidant intake (vitamins A, C, E, and zinc). Results: A response rate of 83% was achieved by providing complete data from 335 children [49% cases with current asthma (n = 166), 51% controls (n = 169)]. Following adjustment for covariates the association between the ratio of n-6:n-3 fatty acids and risk for current asthma was statistically significant (p = 0.022). Conclusion: We found evidence for a modulatory effect of the dietary n-6:n-3 fatty acid ratio on the presence of asthma in children. Our results provide evidence that promotion of a diet with increased n-3 fatty acids and reduced n-6 fatty acids to protect children against symptoms of asthma is warranted.

Asthma

Children

Omega 3:omega 6 fatty acid ratio

Related research ⓘ

People also read

Recommended articles

Cited by 89

Information for

Authors
R&D professionals
Editors
Librarians
Societies

Opportunities

Reprints and e-prints
Advertising solutions
Accelerated publication
Corporate access solutions

Open access

Overview
Open journals
Open Select
Dove Medical Press
F1000Research

Help and information

Help and contact
Newsroom
All journals
Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright © 2025 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)

 **Taylor & Francis Group**
an informa business

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG