







Q

Home ▶ All Journals ▶ Medicine ▶ Journal of Asthma ▶ List of Issues ▶ Volume 41, Issue 3 Ratio of Omega-6 to Omega-3 Fatty Acids

Journal of Asthma >

Volume 41, 2004 - Issue 3

528 92 4 Views CrossRef citations to date Altmetric

Original Article

Ratio of Omega-6 to Omega-3 Fatty Acids and Childhood Asthma

W. H. Oddy, Ph.D., N. H. de Klerk, Ph.D., G. E. Kendall, M.P.H., S. Mihrshahi, M.P.H. & J. K. Peat, Ph.D.

Pages 319-326 | Published online: 26 Aug 2009

66 Cite this article https://doi.org/10.1081/JAS-120026089









66 Citations

Metrics

➡ Reprints & Permissions

Read this article

Share

Abstract

Asthma is a leading cause of morbidity for children and is a major public health problem in Australia. Ecological and temporal data suggest that dietary factors may have a role in recent increases in the prevalence of asthma. Aim: The aim of conducting this study was to investigate whether childhood asthma was associated with the ratio of omega 6 (n-6) to omega 3 (n-3) fatty acids in the diet (n-6:n-3). Method: The Western Australian Pregnancy Cohort Study is a prospective birth cohort of 2602 children. Using a nested case-control cross-sectional study design within this cohort, a group of children were identified as cases with current asthma at 6 or at 8 years of age or as controls with no asthma at 6 or at 8 years. Dietary details including n-6 and n-3 fatty acid intake data were collected by parent response to a questionnaire when the children were 8 years old. Logistical regression was used to compare quartiles of n-6:n-3 intake in cases and

controls. Adjustment was made for covariates: gender, gestational age, breastfeeding, older siblings, maternal smoking during pregnancy, maternal age, maternal asthma, child's current age in months, body mass index, total energy intake, and antioxidant intake (vitamins A, C, E, and zinc). Results: A response rate of 83% was achieved by providing complete data from 335 children [49% cases with current asthma (n=166), 51% controls (n=169)]. Following adjustment for covariates the association between the ratio of n-6:n-3 fatty acids and risk for current asthma was statistically significant (p=0.022). Conclusion: We found evidence for a modulatory effect of the dietary n-6:n-3 fatty acid ratio on the presence of asthma in children. Our results provide evidence that promotion of a diet with increased n-3 fatty acids and reduced n-6 fatty acids to protect children against symptoms of asthma is warranted.

Asthma

Children

Omega 3:omega 6 fatty acid ratio

Related research 1



People also read

Recommended articles

Cited by 92

Omega-6/Omega-3 Essential Fatty Acid Ratio and Chronic Diseases >

Artemis P. Simopoulos M.D.

Food Reviews International Published online: 16 Aug 2006 Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up











Accessibility



Copyright © 2025 Informa UK Limited Privacy policy Cookies Terms & conditions



Registered in England & Wales No. 01072954 5 Howick Place | London | SW1P 1WG