

503 Views | 89 CrossRef citations to date | 4 Altmetric

Original Article

Ratio of Omega-6 to Omega-3 Fatty Acids and Childhood Asthma

W. H. Oddy, Ph.D. , N. H. de Klerk, Ph.D. , G. E. Kendall, M.P.H. , S. Mahrshahi, M.P.H. & J. K. Peat, Ph.D.

Pages 319-326 | Published online: 26 Aug 2009

Cite this article <https://doi.org/10.1081/JAS-120026089>

Sample our Medicine, Dentistry, Nursing & Allied Health Journals >> Sign in here to start your access to the latest two volumes for 14 days

Full Article | Figures & data | References | Citations | Metrics

Reprints & Permissions

Read this article

We Care About Your Privacy

We and our 883 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting I Accept enables tracking technologies to support the purposes shown under we and our partners process data to provide. Selecting Reject All or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the Show Purposes link on the bottom of the webpage. Your choices will have effect within our Website. For more details, refer to our Privacy Policy. [Here](#)

We and our partners process data to provide:

Use precise geolocation data. Actively scan device

I Accept

Reject All

Show Purpose



health problem
have a role
g this study
of omega 6
n Australian
g a nested
ren were
ols with no
take data
re 8 years
cases and
astfeeding,

older siblings, maternal smoking during pregnancy, maternal age, maternal asthma, child's current age in months, body mass index, total energy intake, and antioxidant intake (vitamins A, C, E, and zinc). Results: A response rate of 83% was achieved by providing complete data from 335 children [49% cases with current asthma ($n = 166$), 51% controls ($n = 169$)]. Following adjustment for covariates the association between the ratio of n-6:n-3 fatty acids and risk for current asthma was statistically significant ($p = 0.022$). Conclusion: We found evidence for a modulatory effect of the dietary n-6:n-3 fatty acid ratio on the presence of asthma in children. Our results provide evidence that promotion of a diet with increased n-3 fatty acids and reduced n-6 fatty acids to protect children against symptoms of asthma is warranted.

Asthma

Children

Omega 3:omega 6 fatty acid ratio

Related research

People also read

Recommended articles

Cited by
89



Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright

Accessib

Registered
5 Howick Pl

or & Francis Group
orma business

