







Home ▶ All Journals ▶ Medicine ▶ Journal of Asthma ▶ List of Issues ▶ Volume 41, Issue : ▶ Ratio of Omega-6 to Omega-3 Fatty Acids

Journal of Asthma > Volume 41, 2004 - Issue 3

524 89 4

Views CrossRef citations to date Altmetric

Original Article

Ratio of Omega-6 to Omega-3 Fatty Acids and Childhood Asthma

W. H. Oddy, Ph.D., N. H. de Klerk, Ph.D., G. E. Kendall, M.P.H., S. Mihrshahi, M.P.H. & J. K. Peat, Ph.D.

Pages 319-326 | Published online: 26 Aug 2009

Sample our
Medicine, Dentistry, Nursing
& Allied Health Journals
>> Sign in here to start your access
to the latest two volumes for 14 days

Full Article

Figures & data

References

66 Citations

Metrics

Repri

Abstra

Asthma

in Austra

in recen

was to i

Pregna

case-cor

identifie

asthma

were co

old. Log

controls

(n-6)

We Care About Your Privacy

We and our 907 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting "I Accept" enables tracking technologies to support the purposes shown under "we and our partners process data to provide," whereas selecting "Reject All" or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the ["privacy preferences"] link on the bottom of the webpage [or the floating icon on the bottom-left of the webpage, if applicable]. Your choices will have effect within our Website.

For more details, refer to our Privacy Policy. Here

We and our partners process data to provide:

I Accept

Reject All

Show Purpose have a role

this study

alth problem

of omega 6

Australian

g a nested

lren were

ols with no

take data

re 8 years

cases and

astfeeding,

older siblings, maternal smoking during pregnancy, maternal age, maternal asthma, child's current age in months, body mass index, total energy intake, and antioxidant intake (vitamins A, C, E, and zinc). Results: A response rate of 83% was achieved by providing complete data from 335 children [49% cases with current asthma (n=166), 51% controls (n=169)]. Following adjustment for covariates the association between the ratio of n-6:n-3 fatty acids and risk for current asthma was statistically significant (p=0.022). Conclusion: We found evidence for a modulatory effect of the dietary n-6:n-3 fatty acid ratio on the presence of asthma in children. Our results provide evidence that promotion of a diet with increased n-3 fatty acids and reduced n-6 fatty acids to protect children against symptoms of asthma is warranted.

Asthma Children Omega 3:omega 6 fatty acid ratio

Related research (1)



Information for Open access Authors Overview R&D professionals Open journals Editors **Open Select** Librarians **Dove Medical Press** Societies F1000Research Opportunities Help and information Reprints and e-prints Advertising solutions Newsroom Accelerated publication Corporate access solutions Books Keep up to date Register to receive personalised research and resources by email Sign me up X or & Francis Group Copyright