

478	84	4
Views	CrossRef citations to date	Altmetric

“Cite this article” <https://doi.org/10.1081/JAS-120026089>

Reprints & Permissions | Read this article

[illegible]

Show Purpose



child's current age in months, body mass index, total energy intake, and antioxidant intake (vitamins A, C, E, and zinc). Results: A response rate of 83% was achieved by providing complete data from 335 children [49% cases with current asthma (n = 166), 51% controls (n = 169)]. Following adjustment for covariates the association between the ratio of n-6:n-3 fatty acids and risk for current asthma was statistically significant (p = 0.022). Conclusion: We found evidence for a modulatory effect of the dietary n-6:n-3 fatty acid ratio on the presence of asthma in children. Our results provide evidence that promotion of a diet with increased n-3 fatty acids and reduced n-6 fatty acids to protect children against symptoms of asthma is warranted.

Asthma Children Omega 3:omega 6 fatty acid ratio

Related research ⓘ

People also read

Recommended articles

Cited by
84

Effects of supplementation with omega-3 fatty acids during pregnancy on asthma or wheeze of children: a systematic review and meta-analysis >

Jilei Lin et al.
The Journal of Maternal-Fetal & Neonatal Medicine
Published online: 29 Oct 2018

Omega

Artemis
Journal of
Published

On

Timothy
Expert R
Published



view more



Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Open access


- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up

