



Journal of Environmental Science and Health, Part A >

Toxic/Hazardous Substances and Environmental Engineering

Volume 39, 2004 - [Issue 11-12](#)

229 | 42

Views | CrossRef citations to date | 0 Altmetric

Original Articles

Concentration of Copper, Iron, Zinc, Cadmium, Lead, and Nickel in Bull and Ram Semen and Relation to the Occurrence of Pathological Spermatozoa

P. Massányi , J. Trandzik, P. Nad, B. Koreneková, M. Skalická, R. Toman, ...[show all](#)

Pages 3005-3014 | Received 13 Feb 2004, Published online: 24 Jun 2011

Cite this article

Sample our
Earth Sciences
Journals



>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

Read this article

Share

Abstract

In this study the concentration of copper, iron, zinc, cadmium, lead, and nickel in bull and ram semen and relation of these metals to spermatozoa morphology was investigated. Analysis by atomic absorption spectrophotometry showed that copper concentration was significantly higher ($p < 0.0001$) in ram semen in comparison with bull semen. The zinc concentration was higher in bull semen in comparison with ram semen. The iron and cadmium concentrations in the semen were similar. Higher concentration of lead was found in ram semen. Higher levels of nickel were found in ram semen in comparison with bulls. In bull semen $11.79 \pm 4.88\%$ of pathological spermatozoa was found. Higher occurrence of pathological spermatozoa was in ram semen (17.17 ± 3.76) in comparison with the semen of bulls. Separated tail, tail torso,

and knob twisted tail were the most frequent forms of pathological spermatozoa in both species. Correlation analysis in bulls showed high positive relation between iron and zinc ($r = 0.72$), nickel and separated tail ($r = 0.76$), separated tail and tail torso ($r = 0.71$), tail torso and total number of pathological spermatozoa ($r = 0.72$), and between tail ball and total number of pathological spermatozoa ($r = 0.78$). In rams high positive correlation between cadmium and lead ($r = 0.98$), nickel and separated tail ($r = 0.77$), separated tail and total number of pathological spermatozoa ($r = 0.69$), knob twisted tail and retention of cytoplasmic drop ($r = 0.78$), and between knob twisted tail and other pathological spermatozoa ($r = 0.71$) was found. High negative correlation in ram semen was observed between copper and nickel ($r = 0.71$), copper and separated tail ($r = 0.70$), and between iron and tail torso ($r = 0.67$). The results suggest that the studied metals have a direct effect on spermatozoa quality.

Keywords:

- Trace elements
- Semen
- Pathological spermatozoa
- Bull
- Ram

Acknowledgment

We would like to express our gratitude to A. Sobcakova, P. Sykorova and Ing. M. Cinkanic for technical assistance. This study was supported by VEGA Scientific Grant 1/9080/02 from the Ministry of Education of Slovak Republic.

Related research ⓘ

People also read

Recommended articles

Cited by 42

Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2025 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)



Taylor & Francis Group
an informa business

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG