



Quality Engineering >

Volume 14, 2002 - [Issue 3](#)

10,677 | 912 | 3
Views | CrossRef citations to date | Altmetric

Original Articles

Model-Dependent Variance Inflation Factor Cutoff Values

Trevor A. Craney & James G. Surles

Pages 391-403 | Published online: 15 Feb 2007

Cite this article <https://doi.org/10.1081/QEN-120001878>

Sample our Engineering & Technology Journals
>> [Sign in here](#) to start your access to the latest two volumes for 14 days

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

Read this article

Share

Abstract

When creating designed experiments, it is not always possible to run the experiment at the exact settings required to maintain orthogonal effects. However, this is not measurement error when precise measurements of the settings can be made once the experiment begins. A comparison is made for a 15-run Box-Behnken design using both the intended design settings and the actual design settings. Variance inflation factors are used to measure the induced collinearity in the effects. Two cutoff values are suggested for use to determine when an effect's variance inflation factor is too large to keep that effect in the model.

Keywords:

[Collinearity](#)

[Orthogonal effects](#)

[Variable noise](#)

[Box-Behnken](#)

[Experimental design](#)

Acknowledgments

Related research

People also read

Recommended articles

Cited by
912

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources
by email

 Sign me up

  

  