Separation Science and Technology ›

Volume 38, 2003 - Issue 2

249 Views | 32 CrossRef citations to date | 0 Altmetric

Original Articles

Selective Extraction and Separation of Titanium(IV) from Multivalent Metal Chloride Solutions Using 2-Ethylhexyl Phosphonic Acid Mono-2-ethylhexyl Ester

J. Saji & M. L. P. Reddy

Pages 427-441 | Received 01 Jan 2002, Published online: 15 Feb 2007

Cite this article https://doi.org/10.1081/SS-120016583

Sample our Physical Sciences Journals

>> Sign in here to start your access to the latest two volumes for 14 days

Full Article | Figures & data | References | Citations | Metrics

Reprints & Permissions | Read this article | Share

Abstract

The extraction behavior of titanium(IV) from hydrochloric acid solutions was investigated using 2-ethylhexyl phosphonic acid mono-2-ethylhexyl ester (EHEHPA=HX) in kerosene as an extractant. The results demonstrate that the extraction of titanium(IV) follows the cation exchange mechanism:

$$TiO^{2+}_{aq} + 2(HX)_{2org} \Leftrightarrow TiO(HX_2)_{2org} + 2H^{+}_{aq}$$

where $(HX)_2$refers to the dimeric form of EHEHPA. The equilibrium constant of the extracted complex was deduced by nonlinear regression analysis by taking into account the aqueous phase complexation of metal ion with inorganic ligands and all plausible complexes extracted into the organic phase. The effect of the nature of diluent on the

extraction of titanium(IV) with EHEHPA was investigated and found that the extraction efficiency varies in the order: chloroform<benzene~toluene<xylene<kerosene. IR spectral studies of the extracted complex were used to further clarify the nature of extracted complex. The separation possibilities of titanium(IV) from other associated multivalent metal ions, that is, magnesium(II), aluminum(III), vanadium(V), chromium(III), manganese(II), and iron(III), which are associated with titanium in the waste chloride liquors of titanium minerals processing industry was discussed.

Keywords:

Selective extraction | Separation | Titanium(IV) | 2-Ethylhexyl phosphonic acid mono-2-ethylhexyl ester | Multivalent metal chlorides | Titania wastes

Acknowledgments

One of the authors, JS would like to thank CSIR, New Delhi, for the award of senior research fellowship.

Related research

People also read | Recommended articles | Cited by 32

Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email

Sign me up

Copyright © 2026 Informa UK Limited Privacy policy

Cookies Terms & conditions Accessibility

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG