



Separation Science and Technology >

Volume 38, 2003 - [Issue 2](#)

240 | 32

Views | CrossRef citations to date | Altmetric | 0

Original Articles

# Selective Extraction and Separation of Titanium(IV) from Multivalent Metal Chloride Solutions Using 2-Ethylhexyl Phosphonic Acid Mono-2-ethylhexyl Ester

J. Saji & M. L. P. Reddy

Pages 427-441 | Received 01 Jan 2002, Published online: 15 Feb 2007

Cite this article <https://doi.org/10.1081/SS-120016583>

Sample our  
Physical Sciences  
Journals



>> [Sign in here](#) to start your access  
to the latest two volumes for 14 days

Full Article

Figures & data

References

Citations

Metrics


Reprints & Permissions

Read this article

Share

## Abstract

The extraction behavior of titanium(IV) from hydrochloric acid solutions was investigated using 2-ethylhexyl phosphonic acid mono-2-ethylhexyl ester (EHEHPA=HX) in kerosene as an extractant. The results demonstrate that the extraction of titanium(IV) follows the cation exchange mechanism:



where (HX)<sub>2</sub> refers to the dimeric form of EHEHPA. The equilibrium constant of the extracted complex was deduced by nonlinear regression analysis by taking into account the aqueous phase complexation of metal ion with inorganic ligands and all plausible complexes extracted into the organic phase. The effect of the nature of diluent on the

extraction of titanium(IV) with EHEHPA was investigated and found that the extraction efficiency varies in the order: chloroform<benzene~toluene<xylene<kerosene. IR spectral studies of the extracted complex were used to further clarify the nature of extracted complex. The separation possibilities of titanium(IV) from other associated multivalent metal ions, that is, magnesium(II), aluminum(III), vanadium(V), chromium(III), manganese(II), and iron(III), which are associated with titanium in the waste chloride liquors of titanium minerals processing industry was discussed.

Keywords:

Selective extraction

Separation

Titanium(IV)

2-Ethylhexyl phosphonic acid mono-2-ethylhexyl ester

Multivalent metal chlorides

Titania wastes

## Acknowledgments

One of the authors, JS would like to thank CSIR, New Delhi, for the award of senior research fellowship.

## Related research

People also read

Recommended articles

Cited by  
32

LIQUID-LIQUID EXTRACTION OF TETRAVALENT TITANIUM FROM ACIDIC CHLORIDE SOLUTIONS BY BIS(2,4,4-TRIMETHYLPENTYL)PHOSPHINICACID >

J. Saji et al.

Solvent Extraction and Ion Exchange

Published online: 10 May 2007

## Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

## Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

## Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

## Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

## Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954  
5 Howick Place | London | SW1P 1WG



**Taylor & Francis**  
by informa