



Transactions of the American Fisheries Society >

Volume 130, 2001 - [Issue 5](#)

302 | 72 | 0
Views | CrossRef citations to date | Altmetric

Articles

Response of Yellow Perch to Changes in the Benthic Invertebrate Community of Western Lake Erie

Jeff T. Tyson & Roger L. Knight

Pages 766-782 | Received 19 Sep 2000, Accepted 10 May 2001, Published online: 09 Jan 2011

🗨️ Cite this article 🔗 [https://doi.org/10.1577/1548-8659\(2001\)130<0766:ROYPTC>2.0.CO;2](https://doi.org/10.1577/1548-8659(2001)130<0766:ROYPTC>2.0.CO;2)

Sample our
Economics, Finance,
Business & Industry Journals
>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

Read this article

Share

Abstract

In the western basin of Lake Erie, benthic invertebrate abundance and community composition have changed dramatically over the past five decades, as have abundance and growth of yellow perch *Perca flavescens*. Before 1950, large benthic invertebrates dominated the benthic community of the western basin. Yellow perch readily consumed *Hexagenia* larvae, caddisfly larvae, amphipods, chironomids, and zooplankton. From 1960 through 1980, large-bodied benthic invertebrates were eliminated from the western basin, and yellow perch relied primarily on smaller chironomids and zooplankton as forage. Growth rates and abundance of yellow perch declined through this period in the western basin, in part because of food limitation. From the late 1980s through the 1990s, large benthic invertebrates, including *Hexagenia* spp., caddisfly larvae, and amphipods recolonized the basin, and yellow perch again readily consumed

them. Additionally, daily food consumption by adult yellow perch in the 1990s was marginally more than in the 1980s, suggesting that submaintenance feeding episodes were less frequent. In recent years, yellow perch growth rates have increased modestly, and yellow perch abundance has rebounded. The growth rate of age-3 yellow perch during the year before spawning explained 49% of the variation in age-0 recruitment, indicating that adult growth and condition may influence recruitment. We suggest that increases in benthic macroinvertebrate abundance are responsible, in part, for the increases in yellow perch growth and recruitment. We also suggest that yellow perch diets are a useful indicator of changes in the benthic community.

Related research

People also read

Recommended articles

Cited by
72

Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG

 Taylor and Francis
Group