







Q

Home ► All Journals ► Environment & Agriculture ► Transactions of the American Fisheries Society ► List of Issues ► Volume 130, Issue 5 ► Response of Yellow Perch to Changes in t

Transactions of the American Fisheries Society >

Volume 130, 2001 - <u>Issue 5</u>

 $\begin{array}{c|c} 302 & 71 \\ \text{Views} & \text{CrossRef citations to date} & \text{Altmetric} \end{array}$

Articles

Response of Yellow Perch to Changes in the Benthic Invertebrate Community of Western Lake Erie

Jeff T. Tyson & Roger L. Knight

Pages 766-782 | Received 19 Sep 2000, Accepted 10 May 2001, Published online: 09 Jan 2011



Full Article

Figures & data

Metrics

Reprints & Permissions

Read this article

Share

Abstract

In the western basin of Lake Erie, benthic invertebrate abundance and community composition have changed dramatically over the past five decades, as have abundance and growth of yellow perch Perca flavescens. Before 1950, large benthic invertebrates dominated the benthic community of the western basin. Yellow perch readily consumed Hexagenia larvae, caddisfly larvae, amphipods, chironomids, and zooplankton. From 1960 through 1980, large-bodied benthic invertebrates were eliminated from the western basin, and yellow perch relied primarily on smaller chironomids and zooplankton as forage. Growth rates and abundance of yellow perch declined through this period in the western basin, in part because of food limitation. From the late 1980s through the 1990s, large benthic invertebrates, including Hexagenia spp., caddisfly larvae, and amphipods recolonized the basin, and yellow perch again readily consumed

them. Additionally, daily food consumption by adult yellow perch in the 1990s was marginally more than in the 1980s, suggesting that submaintenance feeding episodes were less frequent. In recent years, yellow perch growth rates have increased modestly, and yellow perch abundance has rebounded. The growth rate of age-3 yellow perch during the year before spawning explained 49% of the variation in age-0 recruitment, indicating that adult growth and condition may influence recruitment. We suggest that increases in benthic macroinvertebrate abundance are responsible, in part, for the increases in yellow perch growth and recruitment. We also suggest that yellow perch diets are a useful indicator of changes in the benthic community.



People also read

Recommended articles

Cited by 71

Spatial and Temporal Patterns of Food Use by White Perch and Yellow Perch in Lake Erie >

Donna L. Parrish et al.

Journal of Freshwater Ecology Published online: 6 Jan 2011

Eutrophication Effects on Prey Size and Food Available to Yellow Perch in Lake Erie >

Robert S. Hayward et al.

Transactions of the American Fisheries Society

Published online: 9 Jan 2011

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up











Accessibility



Copyright © 2025 Informa UK Limited Privacy policy Cookies Terms & conditions



Registered in England & Wales No. 01072954 5 Howick Place | London | SW1P 1WG