On Wednesday 11 November 05:00-22:00 GMT, we'll be making some site updates on Taylor & Francis Online. You'll still be able to search, browse and read our articles, where access rights already apply. Registration, purchasing, activation of tokens, eprints and other features of Your Account will be unavailable during this scheduled work.

Transactions of the American Fisheries Society ›

Volume 130, 2001 - Issue 5

303 Views | 74 CrossRef citations to date | 0 Altmetric

Articles

Response of Yellow Perch to Changes in the Benthic Invertebrate Community of Western Lake Erie

Jeff T. Tyson & Roger L. Knight

Pages 766-782 | Received 19 Sep 2000, Accepted 10 May 2001, Published online: 09 Jan 2011

Cite this article https://doi.org/10.1577/1548-8659(2001)130<0766:ROYPTC>2.0.CO;2

Sample our Environment & Agriculture Journals

>> Sign in here to start your access to the latest two volumes for 14 days

Full Article | Figures & data | References | Citations | Metrics

Reprints & Permissions | Read this article | Share

Abstract

In the western basin of Lake Erie, benthic invertebrate abundance and community composition have changed dramatically over the past five decades, as have abundance and growth of yellow perch Perca flavescens. Before 1950, large benthic invertebrates dominated the benthic community of the western basin. Yellow perch readily consumed Hexagenia larvae, caddisfly larvae, amphipods, chironomids, and zooplankton. From 1960 through 1980, large-bodied benthic invertebrates were eliminated from the western basin, and yellow perch relied primarily on smaller chironomids and zooplankton as forage. Growth rates and abundance of yellow perch declined through

this period in the western basin, in part because of food limitation. From the late 1980s through the 1990s, large benthic invertebrates, including Hexagenia spp., caddisfly larvae, and amphipods recolonized the basin, and yellow perch again readily consumed them. Additionally, daily food consumption by adult yellow perch in the 1990s was marginally more than in the 1980s, suggesting that submaintenance feeding episodes were less frequent. In recent years, yellow perch growth rates have increased modestly, and yellow perch abundance has rebounded. The growth rate of age-3 yellow perch during the year before spawning explained 49% of the variation in age-0 recruitment, indicating that adult growth and condition may influence recruitment. We suggest that increases in benthic macroinvertebrate abundance are responsible, in part, for the increases in yellow perch growth and recruitment. We also suggest that yellow perch diets are a useful indicator of changes in the benthic community.

Related research

People also read | Recommended articles | Cited by 74

Spatial and Temporal Patterns of Food Use by White Perch and Yellow Perch in Lake Erie

Donna L. Parrish et al.
Journal of Freshwater Ecology
Published online: 6 Jan 2011

Eutrophication Effects on Prey Size and Food Available to Yellow Perch in Lake Erie

Robert S. Hayward et al.
Transactions of the American Fisheries Society
Published online: 9 Jan 2011

Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email

Sign me up

Copyright © 2026 Informa UK Limited Privacy policy

Cookies Terms & conditions Accessibility

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG