

Expert Review of Gastroenterology & Hepatology >

Volume 7, 2013 - [Issue 5](#)

94 Views | 11 CrossRef citations to date | 0 Altmetric

Editorial

The hepatitis C virus NS5A inhibitor daclatasvir has a dual mode of action and leads to a new virus half-life estimate

Jeremie Guedj, Harel Dahari, Susan L Uprichard & Alan S Perelson

Pages 397-399 | Published online: 10 Jan 2014

 Cite this article <https://doi.org/10.1586/17474124.2013.811050>

Sample our
Medicine, Dentistry, Nursing
& Allied Health Journals
>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

 Full Article

 Figures & data

 References

 Citations

 Metrics

 Reprints & Permissions

[Read this article](#)

[Share](#)

The hepatitis C virus NS5A inhibitor daclatasvir has a dual mode of action and leads to a new virus half-life estimate

Expert Rev. Gastroenterol. Hepatol. 7(5), 397–399 (2013)



Jeremie Guedj
INSERM, UMR 738, Université Paris Diderot, Sorbonne Paris Cité, Paris 75018, France



Harel Dahari
Division of Hepatology, Department of Medicine, Loyola University Medical Center, Maywood, IL 60153, USA and Theoretical Biology and Biophysics, Los Alamos National Laboratory, Los Alamos, NM 87545, USA



Susan L Uprichard
Division of Hepatology, Department of Medicine, Loyola University Medical Center, Maywood, IL 60153, USA



Alan S Perelson
Author for correspondence: Theoretical Biology and Biophysics, Los Alamos National Laboratory, Los Alamos, NM 87545, USA
Tel.: +1 505 667 6829
asp@lanl.gov

“We employed mathematical modeling to decipher the origin of the very rapid viral decline observed in the first hours following initiation of daclatasvir, a first-in-class NS5A inhibitor, where one dose of daclatasvir led to a 1000-fold decrease in viral levels within approximately 12 h.”

Chronic infection with hepatitis C virus (HCV) affects approximately 160 million people worldwide [1] and is the leading cause of cirrhosis, liver cancer and liver transplantation [2]. Until 2011, the standard of care in the western world was weekly injections of pegylated-interferon (peg-IFN)- α and daily oral ribavirin (RBV), but viral eradication was achieved in less than 50% in treatment-naïve HCV genotype 1 patients, the most prevalent genotype in western world [3].

Since the approval in 2011 of two protease inhibitors (PIs), telaprevir (TVR) and boceprevir, in combination with peg-IFN/RBV, the landscape of anti-HCV therapy has been rapidly changing. Dozens of direct antiviral agents targeting different stages of the viral lifecycle are currently in clinical development, holding the promise that a universal cure can be achieved. Although results from clinical trials, often obtained in small populations of highly selected patients, should be taken with caution, short, tolerable, pan-genotype and IFN-free regimens appear to be attainable within the next couple of years.

In this highly dynamical context, there has been an increasing appeal from the industry and academics to use viral kinetic mathematical modeling to anticipate, evaluate and rationalize the effectiveness of these new antiviral strategies [4]. This is particularly true in the case of agents that target proteins with complex or ill-defined

functions, such as the HCV nonstructural protein 5A. In a recent paper published in *Proceedings of the National Academy of Sciences* [5], we employed mathematical modeling to decipher the origin of the very rapid viral decline observed in the first hours following initiation of daclatasvir (DCV), a first-in-class NS5A inhibitor, where one dose of DCV led to a 1000-fold decrease in viral levels within approximately 12 h [6].

Hepatitis C viral dynamics, virion half-life & drug antiviral effectiveness

One of the most important results of viral kinetic modeling was to introduce the idea that viral load level observed in a patient reflects the balance between the antagonist processes of production and clearance of the virus. By blocking viral production from infected cells, the initiation of antiviral treatment disrupts this equilibrium and the clearance of the virus is no longer efficiently compensated. Thus, the rate of viral decline observed early after treatment initiation can be used to calculate the virus half-life. This basic idea was used by Neumann *et al.* to fit the viral decline observed during interferon therapy and to estimate the antiviral effectiveness of interferon, as well as to obtain the original estimate of the HCV half-life of 2.7 h [7]. Importantly, if HCV is rapidly eliminated from the circulation, it implies that large quantities of the virus

Keywords: daclatasvir • hepatitis C virus • multiscale model • NS5A inhibitor • viral kinetics

EXPERT
REVIEWS

Financial & competing interests disclosure

Portions of this work were performed under the auspices of the US Department of Energy (under contract DE-AC52-06NA25396) and supported by NIH grants (R56/R01-AI078881, P20-GM103452, OD011095, AI028433 and HL109334). AS Perelson has been a consultant for Bristol-Myers Squibb. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

Related research

People also read

Recommended articles

Cited by
11

Information for

Authors
R&D professionals
Editors
Librarians
Societies

Opportunities

Reprints and e-prints
Advertising solutions
Accelerated publication
Corporate access solutions

Open access

Overview
Open journals
Open Select
Dove Medical Press
F1000Research

Help and information

Help and contact
Newsroom
All journals
Books

Keep up to date

Register to receive personalised research and resources by email




Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG

 Taylor and Francis
Group