

397 | 17 | 5
Views | CrossRef citations to date | Altmetric


Review Article

Is dental caries experience increased in HIV-infected children and adolescents? A meta-analysis

Cristiana Aroeira Guimarães Rosa Oliveira, Patricia Nivoloni Tannure, Ivete Pomarico Ribeiro de Souza, Lucianne Cople Maia, Maristela Barbosa Portela & Gloria Fernanda B. de A. Castro 

Pages 481-487 | Received 26 Apr 2014, Accepted 08 Aug 2014, Published online: 12 Mar 2015

 Cite this article  <https://doi.org/10.3109/00016357.2014.958874>

 Check for updates

Sample our
Medicine, Dentistry, Nursing
& Allied Health Journals
>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

 Full Article

 Figures & data

 References

 Citations

 Metrics

 Reprints & Permissions

Read this article

We Care About Your Privacy

We and our 845 partners store and/or access information on a device, such as unique IDs in cookies to process personal data. You may accept or manage your choices by clicking below, including your right to object where legitimate interest is used, or at any time in the privacy policy page. These choices will be signaled to our partners and will not affect browsing data. [Privacy Policy](#)

We and our partners process data to provide:

Use precise geolocation data. Actively scan device characteristics for identification. Store and/or access information on a device. Personalised advertising and content, advertising and content measurement, audience research and services development.

List of Partners (vendors)

 I Accept

Essential Only

Show Purpose

scores in primary teeth in the HIV-infected patients with mean dmft/dmfs scores of 3.8–4.1/7.8–11.0 compared to the control group 1.5–2.4/3.4–5.1. No differences in caries index were found for permanent dentition. The meta-analysis excluded caries data of permanent teeth and showed a significant association between caries experience in primary dentition and HIV infection considering cavitated and non-cavitated lesions (OR = 2.33, 95% CI = 1.48–3.68) or only cavitated lesions (OR = 2.98, 95% CI = 1.59–5.59). Conclusion. Evidence exists that suggests HIV-infected children/adolescents have an increased caries experience in primary dentition.

🔍 Key Words: [child](#) [dental caries](#) [HIV infections](#)

Acknowledgments

This work was supported by CNPq (Comissão Nacional de Desenvolvimento Científico e Tecnológico) and FAPERJ (Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro).

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Related research



Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2024 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)



Taylor & Francis Group
an informa business

Accessibility



Registered
5 Howick Place

