







Home ► All Journals ► Pathology ► List of Issues ► Volume 42, Issue 7 ► A variant in microRNA-196a2 is associate

Pathology > Volume 42, 2010 - Issue 7

Views CrossRef citations to date Altmetric

Molecular Pathology

A variant in microRNA-196a2 is associated with susceptibility to hepatocellular carcinoma in Chinese patients with cirrhosis

Xiao-Dong Li Z, Zhi-Gao Li, Xian-Xu Song & Chun-Fu Liu Z

Pages 669-673 | Received 11 May 2010, Accepted 13 Jul 2010, Published online: 17 Nov 2010

66 Cite this article

Full Article Figure

Figures & data

References

66 Citations

Metrics

➡ Reprints & Permissions

Read this article

Abstract

Aims: Ci while litt

approac

protein

play an

rs11614

lung

Methods without

polymor polymor from the

We Care About Your Privacy

We and our 855 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting "I Accept" enables tracking technologies to support the purposes shown under "we and our partners process data to provide," whereas selecting "Reject All" or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the ["privacy preferences"] link on the bottom of the webpage [or the floating icon on the bottom-left of the webpage, if applicable]. Your choices will have effect within our Website. For more details, refer to our Privacy Policy. Here

We and our partners process data to provide:

I Accept C) in China, raditional used on


Show Purpose t the grancer, small cell on can

2 patients miR-196a2 ment length obtained rels. Results: The frequency of the CC genotype among HCC patients was higher than that in the control group, implying that the cirrhotic patients with the CC genotype or C allele containing genotypes (CT and CC) may have a higher risk of HCC. However, in a subsequent analysis of the association between this polymorphism and clinicopathological characteristics, there was an association between rs11614913 genotype and tumour size (p = 0.046), but not with tumour number, grade, stage, invasiveness or Child-Pugh grade. In a genotype-phenotype correlation analysis using 59 tumour tissues of liver, rs11614913 CC or carrying at least one C allele was associated with significantly increased mature miR-196a expression (p = 0.006 or = 0.002).

Conclusions: Our results suggest, for the first time, that miR-196a2 polymorphism may contribute to cirrhosis-related HCC susceptibility in Chinese patients through influencing mature miR-196a expression.

Q Key words: Cirrhosis hepatocellular carcinoma MiR-196a2 single nucleotide polymorphism





Information for Open access Authors Overview R&D professionals Open journals Editors **Open Select** Librarians **Dove Medical Press** Societies F1000Research Opportunities Help and information Reprints and e-prints Advertising solutions Newsroom Accelerated publication Corporate access solutions Books Keep up to date Register to receive personalised research and resources by email Sign me up X or & Francis Group Copyright Registered 5 Howick Pl