







Home ▶ All Journals ▶ Scandinavian Journal of Gastroenterology ▶ List of Issues ▶ Volume 46, Issue 1
▶ Nurse-administered propofol sedation for

Scandinavian Journal of Gastroenterology > Volume 46, 2011 - Issue 12

362 | 33

0

Views CrossRef citations to date Altmetric

Endoscopy

Nurse-administered propofol sedation for gastrointestinal endoscopic procedures: first Nordic results from implementation of a structured training program

Charlotte Slagelse, Peter Vilmann

→, Pernille Hornslet, Anne Hammering & Teit Mantoni

Pages 1503-1509 | Received 03 Jun 2011, Accepted 25 Aug 2011, Published online: 04 Nov 2011

Sample our
Medicine, Dentistry, Nursing
& Allied Health Journals
>> Sign in here to start your access
to the latest two volumes for 14 days

Full Article

Figures & data

References

66 Citations

Metrics

Reprints & Permissions

Read this article

Abstra/

Introduc

program

sedation

for endo

give

including endosco including

2527 pa

The pati

We Care About Your Privacy

We and our 842 partners store and/or access information on a device, such as unique IDs in cookies to process personal data. You may accept or manage your choices by clicking below, including your right to object where legitimate interest is used, or at any time in the privacy policy page. These choices will be signaled to our partners and will not affect browsing data. Privacy Policy

We and our partners process data to provide:

Use precise geolocation data. Actively scan device characteristics for identification. Store and/or access information on a device. Personalised advertising and content, advertising and content measurement, audience research and services development.

List of Partners (vendors)

Essential Onlintroduced

Show Purpose

Tof
fol was
ek course
r
strict criteria

Results.


e included.

ely. Median

dose of propofol was 300 mg. No mortality was noted. 119 of 2527 patients developed short lasting hypoxia (4.7%); 61 (2.4%) needed suction; 22 (0.9%) required bag-mask ventilation and 8 (0.3%) procedures had to be discontinued. In 11 patients (0.4%), anesthetic assistance was called due to short lasting desaturation. 34 patients (1.3%) experienced a change in blood pressure greater than 30%. Conclusion. NAPS provided by properly trained nurses according to the present protocol is safe and only associated with a minor risk (short lasting hypoxia 4.7%). National or international structured training programs are at present few or non-existing. The present training program has documented its value and is suggested as the basis for the current development of guidelines.

Q Key Words:: endoscopy-general endoscopy-interventional general

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.



Information for Open access Authors Overview R&D professionals Open journals Editors **Open Select** Librarians **Dove Medical Press** Societies F1000Research Opportunities Help and information Reprints and e-prints Advertising solutions Newsroom Accelerated publication Corporate access solutions Books Keep up to date Register to receive personalised research and resources by email Sign me up Taylor & Francis Group Copyright © 2024 Informa UK Limited Privacy policy Cookies Terms & conditions Accessib X

