



387 | 34

Views | CrossRef citations to date | 6

Altmetric

Endoscopy

Nurse-administered propofol sedation for gastrointestinal endoscopic procedures: first Nordic results from implementation of a structured training program

Charlotte Slagelse, Peter Vilmann , Pernille Hornslet, Anne Hammering & Teit Mantoni

Pages 1503-1509 | Received 03 Jun 2011, Accepted 25 Aug 2011, Published online: 04 Nov 2011

 Cite this article  <https://doi.org/10.3109/00365521.2011.619274>

Sample our
Medicine, Dentistry, Nursing
& Allied Health Journals
>> **Sign in here** to start your access
to the latest two volumes for 14 days

 Full Article

 Figures & data

 References

 Citations

 Metrics

 Reprints & Permissions

Read this article

 Share

Abstract

Introduction. Proper training to improve safety of NAPS (nurse-administered propofol sedation) is essential. **Objective.** To communicate our experience with a training program of NAPS. **Materials and methods.** In 2007, a training program was introduced for endoscopists and endoscopy nurses in collaboration with the Department of Anaesthesiology. During a 2.5-year period, eight nurses were trained. Propofol was given as monotherapy. The training program for nurses consisted of a 6-week course including theoretical and practical training whereas the training program for endoscopists consisted of 2.5 h of theory. Patients were selected based on strict criteria including patients in ASA (American Society of Anesthesiologists) group I-III. **Results.** 2527 patients undergoing 2.656 gastrointestinal endoscopic procedures were included.

The patients were ASA group I, II and III in 34.7%, 56% and 9,3%, respectively. Median dose of propofol was 300 mg. No mortality was noted. 119 of 2527 patients developed short lasting hypoxia (4.7%); 61 (2.4%) needed suction; 22 (0.9%) required bag-mask ventilation and 8 (0.3%) procedures had to be discontinued. In 11 patients (0.4%), anesthetic assistance was called due to short lasting desaturation. 34 patients (1.3%) experienced a change in blood pressure greater than 30%. Conclusion. NAPS provided by properly trained nurses according to the present protocol is safe and only associated with a minor risk (short lasting hypoxia 4.7%). National or international structured training programs are at present few or non-existing. The present training program has documented its value and is suggested as the basis for the current development of guidelines.

Key Words::

- endoscopy-general
- endoscopy-interventional
- general

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Related Research Data

[Automated graphic assessment of respiratory activity is superior to pulse oximetry and visual assessment for the detection of early respiratory depression during therapeutic upper endoscopy](#)
Source: Gastrointestinal Endoscopy

[Endoscopist-Directed Administration of Propofol: A Worldwide Safety Experience](#)
Source: Gastroenterology

[Sedation, surveillance, and preparation](#)
Source: Endoscopy

[Patient-maintained sedation for ERCP with a target-controlled infusion of propofol: A pilot study](#)
Source: Gastrointestinal Endoscopy

[A randomized controlled trial on use of propofol alone versus propofol with midazolam, ketamine, and pentazocine “sedato-analgesic cocktail” for sedation during ERCP](#)

Source: Endoscopy

Risk stratification and safe administration of propofol by registered nurses supervised

Related research

People also read

Recommended articles

Cited by
34

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2025 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)



Taylor & Francis Group
an informa business

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG