



Open access

1,305 Views | 22 CrossRef citations to date | 10 Altmetric

Listen

ORIGINAL ARTICLE

"I'm Dr Jekyll *and* Mr Hyde": Are GPs' antibiotic prescribing patterns contextually dependent? A qualitative focus group study

Eva Lena Strandberg , Annika Brorsson, Charlotta Hagstam, Margareta Troein & Katarina Hedin

Pages 158-165 | Received 27 Mar 2013, Accepted 08 May 2013, Published online: 14 Aug 2013

Cite this article <https://doi.org/10.3109/02813432.2013.824156>

Full Article

Figures & data

References

Citations

Metrics

Licensing

Reprints

We Care About Your Privacy

We and our 912 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting I Accept enables tracking technologies to support the purposes shown under we and our partners process data to provide. Selecting Reject All or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the Show Purposes link on the bottom of the webpage. Your choices will have effect within our Website. For more details, refer to our Privacy Policy. [Here](#)

We and our partners process data to provide:

Use precise geolocation data. Actively scan device

I Accept

Reject All

Show Purposes



Several factors influence the meeting and contribute to enhancing the conditions for rational prescribing. These conditions are connected to the GP, the relationship, and the setting; organization as well as professional culture. The findings indicate synergies between the factors, and that one factor can sometimes compensate for lack of another. Continuity and mutual trust can make a brief consultation successful, but lack of continuity can eliminate the effects of knowledge and professional skills.

Conclusions. The findings emphasize the importance of the encounter between the GP and the patient for prudent antibiotic prescribing. Furthermore, the importance of an appropriate organization of primary care, which promotes continuity and encourages professional autonomy, is demonstrated.

Key Words::

- Antibiotic prescribing
- consultation
- doctor-patient relationship
- general practice
- primary health care
- respiratory tract infections
- Sweden

In spite of numerous efforts to reduce unnecessary antibiotic prescribing, the problem

- The
- A



Introduction

Respiratory encounters in primary care are characterized by a high prevalence of bacterial infections, which often leads to over-use of antibiotics.

has increased antibiotic resistance, and there is evidence that resistance develops not only on the aggregated level but also in the individual [2].

The reasons for prescribing antibiotics contrary to current guidelines have been discussed and studied extensively [3–6]. In a Canadian historical cohort it was shown that inappropriate antibiotic prescribing increased with time in practice and was also more frequent among foreign medical graduates and among primary care physicians with a high practice volume [3].

A well-known problem is the constant prescribing pattern on an individual basis. Cars and Håkansson concluded that “Doctors have an individual and very constant pattern of prescribing antibiotics, and it seems that the diagnoses are often given to justify the treatment, rather than the other way around” [4]. Some other studies concluded that a high frequency of prescriptions of antibiotics may reflect a general disposition among GPs to give priority to maintaining good relations with the patients [5,6]. They suggest that low antibiotic prescribing and high patient satisfaction could be combined if sufficient time were spent on listening to patients. Patient satisfaction was not related to the prescription of antibiotics, but to a better understanding of their illness.

Numerous efforts have been made to reduce unnecessary and potentially harmful antibiotic prescribing. These include education for providers, near-patient tests, and the introduction of financial incentives, as well as information to patients and public [7–10].

Case-based prescribing, where the physician is supported by a decision support system, has been shown to improve prescribing. A study in Sweden showed that physicians who used the system responded with altered prescribing patterns, and the system was able to detect and prevent inappropriate prescribing. A follow-up study after implementation showed that the system was used, and the prescribing patterns had changed.

A recent study in Sweden showed that a decision support system can help to improve prescribing. The system was able to detect and prevent inappropriate prescribing, and the prescribing patterns had changed. The system was used, and the prescribing patterns had changed.

According to the study, the system was able to detect and prevent inappropriate prescribing, and the prescribing patterns had changed. The system was used, and the prescribing patterns had changed.



Material and methods

The APO method

This paper reports on one component of the EU project HAPPY AUDIT [[13](#)]. The Audit Project Odense (APO) method was tested to change antibiotic prescribing habits in six countries, chosen for their differing prevalence of antibiotic resistance. The audit was performed twice, with a year's interval during which an intervention took place. Considerable changes in prescribing habits were registered [[13,14](#)]. However, in Sweden, substantial differences between practices remained after the intervention. This suggests that the application of guidelines varies due to factors other than medical.

Study design

Focus groups were used to explore GPs' perceptions of antibiotic prescribing for RTIs in primary care. Focus groups are effective for exploring the meaning of conceptions and definitions of important phenomena in health care [[15,16](#)].

The interactive discussions of focus groups generate valuable details of complex experiences and reasons behind actions, beliefs, perceptions, and attitudes [[17,18](#)]. Focus-group interaction facilitates in-depth discussions through a sequence of open-ended questions that encourage participation within the group. Group dynamics create a synergy of ideas, where the interaction of individuals produces ideas that are not likely to emerge from individual contributions. A central element in the focus group method is the interaction between participants and the group as a whole.

All authors contributed equally and significantly to the research. The research was supervised by the first author (J.B.). The research was conducted in various settings, including primary care, hospital, and research institutions, with experienced researchers and clinicians.

Data collection

Two focus groups were arranged in each country (Sweden, Norway, Denmark, Finland, and the Netherlands). The participants were GPs with at least 5 years of experience in primary care. The focus groups were conducted once for



questions [20]. Topics for the focus group discussions were (i) Experiences of antibiotic prescribing for RTIs and personal behaviour, (ii) Experiences of deviating from national and local guidelines, (iii) The impact of educational interventions such as the Happy Audit project.

Table I. Distribution of participating GPs.

[Download CSV](#) [Display Table](#)



A moderator introduced each focus, and made sure that participants did not deviate from the research question and that everyone was able to take part in the discussion. An assistant took notes, which were added to the data from the audiotaped discussion that was transcribed verbatim. To validate the focus-group data, ELS and AB did three individual telephone interviews with Happy Audit participants from a third Swedish region.

Data analysis

An editing analysis style according to Miller and Crabtree [21] was used starting with a naive reading followed by repeated readings of all transcripts that were examined independently and on their own and prior to categorising the data into categories in an iterative process. The exploratory analysis was continued until ELS and AB met and discussed the data until they were satisfied that they contained



When referring in the results to the participants’ statements, these are written in the past tense, while interpretations of statements are in the present tense.

Ethical considerations

According to Swedish legislation, ethical approval was not necessary for this study since it was considered a quality improvement project. All participants were informed that participation was voluntary, that they could withdraw at any time, that all data would be handled confidentially, and that the results would be presented in a non-identifiable way.

Results

We found that the most significant and important factor concerning antibiotic prescribing for RTIs is what happens when the GP meets the patient ([Table II](#)). Our main finding concerns two dimensions of this encounter: characteristics of the meeting depending on the interaction with the patient ([Figure 1](#)) and influencing factors ([Figure 2](#)).

Figure 1. Characteristics of the encounter between the GP and the patient.



Agree

Display full

Figure 2. Factors influencing the encounter between the GP and the patient.



Display full size

Charac

When th
recogniz
compror
fight wh
sometim

Fact

When lo
influenci
the GPs
encount

x

ter is
r
rized as a
omise,
d factors
main factors:
ich the

of evidence-based medicine, as a GP always has to take into account the patient in front of him/her. Professional performance is about being able to judge what is best for the patient.

Setting of the encounter. The organization of the work may facilitate or complicate the rational use of antibiotics for RTIs.

A facilitating factor that was mentioned is the overall consensus at the primary health care centre (PHC) about how to treat patients with RTIs. Overall consensus and a common professional culture are achieved through local professional discussions and exchange of experience. Teamwork with primary nurse assessment and return visits for follow-up are also considered facilitating.

The availability of ordinary GPs is a facilitating factor as well. A primary care practice with ordinary GPs promotes continuity of care. Many Swedish physicians today are hired for a shorter period of time without a list of patients of their own and they are, according to our participants, the ones who usually meet the RTI patients, a kind of fragmentation of the workload. According to the participants, they seem to have a tendency to prescribe more antibiotics for RTIs than the ordinary GP.

GPs who reported that they normally follow the guidelines admitted that they prescribe more antibiotics when on call or at the emergency centre. This is due to lack of time and possibly lack of information about the patient:



Discus

Summa



The GP and the process disagree condition relations well as t

of the
between the
the ensuing
e, or
ancing the
P, the
nization as
es between

Strengths and limitations

This study is part of a multi-method approach to deepen the understanding of circumstances favourable to rational antibiotic prescribing. Focus groups were chosen for data collection. We planned for and conducted two groups and were prepared for a third if the data achieved proved to be insufficient. This method has the advantage of producing ample data in a limited time, and the group process between the participants stimulates the sharing of experiences. The participants were strategically chosen among those who participated in both registrations in the Happy Audit project, indicating that they were actively and practically engaged in rational prescribing and thus could be expected to prescribe antibiotics more prudently. We recruited participants to get as broad a representation as possible from the Happy Audit material, which means that the invited participants represented both urban and rural areas as well as low-prescribing and high-prescribing districts/areas. According to the literature there is no magic number concerning sample size in focus-group interviews. A common recommendation is two to five groups [22]. Sample size is not, according to Crabtree and Miller [21], “the determinant of research significance in a qualitative study; the major concern is with information richness”.

The GPs had voluntarily participated in the Happy Audit project including the intervention. It is therefore likely that they have more knowledge and are more reflective. This is, they share nu is, they believe that these ex

The data expectations and exp cannot be generaliz aim of which is to con assume that

Compa

The find expectat ted by similar r



One result was the importance of the knowledge level of the GP, also corroborated in a previous study [24]. The findings indicate that continuity is favourable for the relationship between GP and patient, but it is obvious from another study [25] that when patients do not get well, continuity is not sufficient.

The participants in this study revealed experiences concerning the consequences of the organization for prescribing of antibiotics. The organization of primary care may be important in at least two ways. It can facilitate a good relationship between the GP and the patient by promoting continuity. A stable relationship based on continuity increases the chances of a smooth encounter. Thus, inappropriate organization may indirectly contribute to increased antibiotic resistance.

Second, stable staff at a PHC is a prerequisite for the formation of a professional culture with a local common policy. This can be labelled continuity on the aggregated level. The findings of this study indicate that a common policy in a primary care centre is a success factor in the task of improving prescribing habits. Corresponding results have been found in previous research as well as in a focus-group study on similar topics among Happy Audit participants in Lithuania and the Kaliningrad region [26,27].

A professional culture with a common policy may not simply be regarded as a result of adherence to guidelines, but also as an autonomous attitude: a genuine desire to do things right and make a difference [28].

An alternative way to improve prescribing habits is through continuing medical education (CME) for GPs. A common policy in a primary care centre entails synergies between GPs and the organization, which includes both GPs and the organization. A professional culture is created.

In Sweden, the use of antibiotics is high. For instance, the use of antibiotics is widespread in primary care. The purpose of the study was to investigate the use of antibiotics in primary care [29,30].

In future research, the importance of a professional culture and the role of the organization in Sweden should be investigated.



Our findings emphasize the importance of the encounter between the GP and the patient for decisions concerning antibiotic prescribing. A stable relationship between the two and the GP's knowledge of guidelines contribute to prudent prescribing. Furthermore, the importance of appropriate organization of primary care, which promotes continuity of care and encourages professional autonomy, is demonstrated.

Authors' contributions

ELS and AB were equally involved in study design, conducted and analysed the focus groups and interviews, wrote the paper, and will serve as guarantors for the integrity of the data. CH and KH did the literature search. MT contributed to editing the paper. All authors have read and approved the final manuscript.

Acknowledgements

The authors would like to thank the GPs who chose to take part in the focus groups for this study for their time and contribution. This project was prepared as part of the EU project HAPPY AUDIT, a Specific Targeted Research Project (STREP) funded by the European Commission: DG SANCO under the Framework Programme 6 (SP5A-CT-2007-044154). Contract Number 044154.

Special thanks to the GPs who participated in the focus groups for their time and contribution. The authors also thank the European Commission for the funding of this project.



Related research
Effectiveness of antibiotic prescribing in primary care

cluster randomised trial

Source: BMJ

Capturing the Group Effect in Focus Groups: A Special Concern in Analysis

Source: Qualitative Health Research

Improvements in antimicrobial prescribing for treatment of upper respiratory tract infections through provider education

Source: Journal of General Internal Medicine

Antibiotics Prescription in Primary Care: A 5-Year Follow-up of an Educational Programme

Source: Family Practice

Intervention with educational outreach at large scale to reduce antibiotics for respiratory tract infections: a controlled before and after study

Source: Family Practice

Pay-for-virtue: an option to improve pay-for-performance?

Source: Journal of Evaluation in Clinical Practice

General practitioners' strategies to identify alcohol problems: A focus group study

Source: Scandinavian Journal of Primary Health Care

Awareness of antibiotic resistance and antibiotic prescribing in UTI treatment: A qualitative study among primary care physicians in Sweden

Source:

Focus

Source:

The p

distrib

Source:

Health

tr

in

Source:

An o

chan

Source:

To pr

great



Antibiotic Use for Emergency Department Patients With Upper Respiratory Infections: Prescribing Practices, Patient Expectations, and Patient Satisfaction

Source: Annals of Emergency Medicine

Einnahme von Antibiotika führt teilweise zu längerfristiger Resistenz

Source: Unknown Repository

Intrinsic and Extrinsic Motivations: Classic Definitions and New Directions

Source: Contemporary Educational Psychology

What about N? A methodological study of sample-size reporting in focus group studies

Source: BMC Medical Research Methodology

Is my child sick? Parents' management of signs of illness and experiences of the medical encounter: Parents of recurrently sick children urge for more cooperation

Source: Scandinavian Journal of Primary Health Care

Interventions to improve antibiotic prescribing practices in ambulatory care

Source: Cochrane Database of Systematic Reviews

Revenge of the killer microbe

Source: CMAJ

The effect of financial incentives on the quality of health care provided by primary care physicians

Source:

Trends

tract

Cour

Sour

The

in ge

So

T

Sour

Predi

phys

Sour

A stu

Sour




Source: Swiss Medical Weekly

Health Alliance for Prudent Prescribing, Yield and Use of Antimicrobial Drugs in the Treatment of Respiratory Tract Infections (HAPPY AUDIT)

Source: BMC Family Practice

Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: systematic review and meta-analysis

Source: BMJ

Linking provided by 

References

1. Neumark T, Brudin L, Engström S, Mölsted S. Trends in number of consultations and antibiotic prescriptions for respiratory tract infections between 1999 and 2005 in primary health care in Kalmar County, Southern Sweden. Scand J Prim Health Care 2009;27:3-7.

[Web of Science ®](#)

[Google Scholar](#)

2. Costelloe C, Metcalfe C, Lovering A, Mant D, Hay AD. Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: Systematic review and meta-

3. Cadieu P, et al. Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: Systematic review and meta-

4. Cars H, et al. Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: Systematic review and meta-

5. Lundkvist M, et al. Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: Systematic review and meta-



[PubMed](#)[Web of Science ®](#)[Google Scholar](#)

6. Ong S, Nakase J, Moran GJ, Karras DJ, Kuehnert MJ, Talan DA. Antibiotic use for emergency department patients with upper respiratory infections: Prescribing practice, patient expectations, and patient satisfaction. *Ann Emerg Med* 2007;50:213-20.

[PubMed](#)[Web of Science ®](#)[Google Scholar](#)

7. Juzych NS, Banerjee M, Essenmacher L, Lerner SA. Improvements in antimicrobial prescribing for treatment of upper respiratory tract infections through provider education. *J Gen Intern Med* 2005;20: 901-5.

[PubMed](#)[Web of Science ®](#)[Google Scholar](#)

8. Smeets HM, Kuyvenhoven MM, Akkerman AE, Welschen I, Schouten GP, van Essen GA, et al. Intervention with educational outreach at large scale to reduce antibiotics for respiratory tract infections: A controlled before and after study. *Fam Pract* 2009, 26: 183-7.

[PubMed](#)[Web of Science ®](#)[Google Scholar](#)

9. Briel M, et al. Community-based educational interventions: A cluster randomized controlled trial. *Ann Intern Med* 2006;144:100-10.

[PubMed](#)

10. Carr SA, et al. Are we testing for C reactive protein in lower respiratory tract infections? *Arch Dis Child* 2004;89:100-104.

11. Arnolds J, et al. Antibiotic practice in

2. Mölstað S, Ekedahl A, Hovelius B, Thimansson H. Antibiotics prescription in primary care: A 5-year follow-up of an educational programme. *Fam Pract* 1994;11:282-6.

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

3. Bjerrum L, Munck A, Gahrn-Hansen B, Plejdrup Hansen M, Jarboel DE, Cordoba G, et al. Health Alliance for prudent antibiotic prescribing in patients with Respiratory Tract Infections (HAPPY AUDIT) – impact of a non-randomised multifaceted intervention programme. *BMC Fam Pract* 2011;12:52.

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

4. Landstinget Blekinge. Happy Audit Report: Swedish registrations 2008 and 2009. Karlskrona, Sweden: Landstinget Blekinge; 2010.

[Google Scholar](#)

5. Strandberg EL, Ovhed I, Borgquist L, Wilhelmsson S. The perceived meaning of a (w)holistic view among general practitioners and district nurses in Swedish primary care: A qualitative study. *BMC Fam Pract* 2007;8:8.

6. Lid TG. Problems: A focus

7. Ca. al concern in ana

8. Krueg Thousand Oaks,



19. Parker A, Tritter J. Focus group method and methodology: Current practice and recent debate. *Int J Research & Method in Education* 2006;29:23–37.

[Google Scholar](#)

20. Kvale S. *Den kvalitativa forskningsintervjun* [The qualitative research interview]. Lund: Studentlitteratur; 2009.

[Google Scholar](#)

21. Miller WL, Crabtree BF. Primary care research: A multi method typology and qualitative road map. In: Crabtree BF, Miller WL, editors. *Doing qualitative research: Research methods for primary care*. Vol. 3. Newbury Park, CA: Sage Publications; 1992.

[Google Scholar](#)

22. Carlsen B, Glenton C. What about N? A methodological study of sample-size reporting in focus group studies. *BMC Med Res Methodol* 2011;11:26–35.

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

23. Hedin K, Andre M, Håkansson A, Mölsted S, Redhe N, Petersson C. A population-based study of...

2006;56:680–

5.

[PubMed](#)

24. Björk... resistance
and at... primary care
ph...



25. Ertma... gement of
signs... currently sick
childre... 7.

26. Jacoby A, Smith M, Eccles M. A qualitative study to explore influences on general practitioners' decisions to prescribe new drugs. *Br J Gen Pract* 2003;53:120-5.

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

27. Jaruseviciene L, Radzeviciene-Jurgute R, Lazarus JV, Jurgutis A, Ovhed I, Strandberg EL, et al. A study of antibiotic prescribing: The experience of Lithuanian and Russian GPs. *Cent Eur J Med*. 2012;7:790-9.

[Web of Science ®](#) | [Google Scholar](#)

28. Ryan RM, Deci EL. Intrinsic and extrinsic motivations: Classic definitions and new directions. *Contemp Educ Psychol* 2000;25:54-67.

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

29. Scott A, Sivey P, Ait Ouakrim D, Willenberg L, Naccarella L, Furler J, et al. The effect of financial incentives on the quality of health care provided by primary care physicians. *Cochrane Database Syst Rev* 2011;7:9: CD008451. Review.

[PubMed](#) | [Google Scholar](#)

30. Bueton S, Entwistle V. Pay for virtue: An option to improve pay for performance? *J Eval C*



Download



Rel

Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright

Accessib

Registered
5 Howick Pl

or & Francis Group
orma business

