

Home ► All Journals ► Medicine ► Disability and Rehabilitation ► List of Issues ► Volume 37, Issue 7 ► Effect of trunk-restraint training on fu ....


Q

### Disability and Rehabilitation >

Volume 37, 2015 - Issue 7

1,507401ViewsCrossRef citations to dateAltmetric

Review



Liza M. Pain , Ross Baker, Denyse Richardson & Anne M. R. Agur Pages 553-562 | Received 01 Dec 2013, Accepted 04 Jun 2014, Published online: 25 Jun 2014




### Abstract

Purpose: The purpose of this systematic review was to determine the effect of trunk restraint (TR) training on post-stroke compensatory trunk movements during functional reach, and to identify functional gains of TR within the context of the International Classification of Functioning, Disability and Health (ICF). Methods: Six databases were searched prior to 20 February 2013. Randomized control trials (RCTs) comparing the effects of a post-stroke reaching intervention with and without TR in the adult poststroke population were included in this review. Data related to study inclusion/exclusion criteria, methodology, study participants and between-group outcomes (p < 0.05) were extracted. All outcomes were categorized according to functional domains defined within the ICF. Results: Five RCTs met the inclusion criteria. TR groups demonstrated significant between-group differences (p < 0.05) for decreased compensatory trunk displacement (4/5 RCTs), increased elbow extension (3/5 RCTs) and increased shoulder flexion (2/5 RCTs). All significant between-group differences fell within the Body Structure/Function domain of the ICF. Conclusions: Trunk restraint is a simple, costeffective technique that may help to reduce compensatory trunk/shoulder/elbow movements in the post-stroke adult population. Synthesis of study outcomes also highlights applications of TR to clinical practice and areas for further research.

Implications for Rehabilitation

- The ability to use the shoulder and elbow to perform functional reach is a primary goal in post-stroke recovery; however, compensatory trunk movements are often used to achieve the reaching goal.
- Long-term use of compensatory strategies may contribute to secondary impairments, such as learned non-use, joint contractures and pain.
- Trunk restraint enables functional reach practice, while limiting compensatory strategies in the moderately to severely impaired stroke population.



## Acknowledgements

The authors would like to acknowledge Shannon Roberts and Tapas Pain for their assistance with manuscript review and formatting.

# Declaration of interest

The authors declare no potential conflicts of interest with respect to the authorship and/or publication of this article.

A review protocol was determined a priori by the review authors in terms of the type of study design, study population, types of interventions and outcome measures; however, a formal protocol was not registered with a registry service. The authors received no financial support for the research and/or authorship of this article.



Related research 1

People also read	Recommended articles	Cited by 40
------------------	----------------------	----------------

Information for	Open access
Authors	Overview
R&D professionals	Open journals
Editors	Open Select
Librarians	Dove Medical Press
Societies	F1000Research
Opportunities	Help and information
Reprints and e-prints	Help and contact
Advertising solutions	Newsroom
Accelerated publication	All journals
Corporate access solutions	Books

#### Keep up to date

Register to receive personalised research and resources by email





Copyright © 2025 Informa UK Limited Privacy policy Cookies Terms & conditions

Taylor & Francis Group an informa business



Registered in England & Wales No. 01072954 5 Howick Place | London | SW1P 1WG