

Free access

2,343 Views | 39 CrossRef citations to date | 12 Altmetric

Listen

RESEARCH COMMUNICATIONS

Effects of molybdenum on sperm quality and testis oxidative stress

Xiao-Wei Zhai, Yu-Ling Zhang, Qiao Qi, Yu Bai, Xiao-Li Chen, Li-Jun Jin, ... show all
Pages 251-255 | Received 23 Nov 2012, Accepted 12 Feb 2013, Published online: 08 May 2013

Cite this article <https://doi.org/10.3109/19396368.2013.791347>

Full Article | Figures & data | References | Citations | Metrics

Reprints & Permissions | View PDF | Share

We Care About Your Privacy

We and our 887 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting I Accept enables tracking technologies to support the purposes shown under we and our partners process data to provide. Selecting Reject All or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the Show Purposes link on the bottom of the webpage. Your choices will have effect within our Website. For more details, refer to our Privacy Policy. [Here](#)

We and our partners process data to provide:

Use precise geolocation data. Actively scan device

I Accept

Reject All

Show Purpose

Introduction

A variety of endocrine disrupting chemicals have been released into the environment in the rapid industrial progress, which may exert adverse health effects in human and animals [Carlsen et al. [1992](#); Khurana et al. [2000](#); Friedmann [2002](#)]. Molybdenum (Mo) is an essential trace element in animals and humans. It has been identified as part of the active sites of over 50 enzymes, and may promote normal cell function possibly by catalyzing a variety of hydroxylation, oxygen atom transfer and other oxidation-reduction reactions [Hille et al. [1998](#)]. Molybdenum is also an endocrine disruptor and has been widely present and detected in our food and water [Underwood [1981](#); Mills and Davis [1987](#); Kargar et al. [2011](#); Yu et al. [2011](#)]. In addition, Mo is broadly used in industrial production, such as metallurgical processes, the manufacture of electronic products, glass, ceramics, lubricants, catalysts, pigments and nano materials [Pandey and Singh [2002](#); CDC 2005; Braydich-Stolle et al. [2005](#); Ema et al. [2010](#)]. Furthermore, Mo is also an environmental pollutant discharged from uranium processing, combustion processing, contact lens solutions, and the color additives in cosmetics [ACGIH 1995]. This wide distribution greatly increases the risk of animals and humans exposed to the high level of Mo in the environment. For example, molybdenum concentrations have

risen to maximum levels in the environment. Excessive exposure to Mo in animals and humans through food and water ingestion can lead to various abnormal effects [Pandey and Singh [2002](#)]. In the study of Meeker et al. [2008](#), molybdenum concentrations in the environment of male animals were found to be significantly higher than those of female animals [Meeker et al. [2008](#)]. In rats, molybdenum exposure led to morphologic changes in the testes [Pandey and Singh [2002](#)]. A reduction in the number of sperm was observed in the study of Bersényi et al. [2008](#). Meeker et al. [2008](#) also reported that molybdenum exposure led to a reduction in the number of male animals. Oxidative stress and even DNA damage were observed in the study of Meeker et al. [2008](#).



titanium-molybdenum generated oxidative stress in mouse fibroblasts cultures.

Antioxidants provide a defense against oxidative stress.

Seminal plasma contains three main enzymatic antioxidants: superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPx). Spermatozoa possess primarily enzymatic antioxidants, with SOD being the most predominant [Makker et al. [2009](#)]. Superoxide dismutase in conjugation with CAT and GPx scavenge both intracellular and extracellular superoxide radicals and inhibit lipid peroxidation [Agarwal and Prabhakaran [2005](#)]. Malondialdehyde (MDA) is one of the byproducts of lipid peroxidation that indirectly reflects the level of peroxidation and the degree of cell injury [Sharma et al. [2004](#)]. This byproduct has been used in various biochemical assays to monitor the degree of oxidative damage sustained by spermatozoa [Aitken et al. [1989](#); Aitken and Fisher [1994](#)]. Oxidative stress can be evaluated by detecting the activities of SOD and GPx, as well as the MDA level in the tissue. Up to now, there are no data available for the oxidative stress of testicular tissue caused by Mo on mice.

This study has been undertaken to evaluate the effect of orally administered Mo on the sperm parameters of the epididymis index, sperm motility, count, and morphology changes. The oxidative stress of testis was considered as a function of the levels of MDA, SOD, and GPx in mice.

Result

The epididymis index, sperm count, and morphology changes in morphological parameters of sperm at 25 mg/L molybdenum affected the sperm quality significantly.




Figure 1 shows the effect of 25 mg/L molybdenum on the epididymis index, sperm count, and morphology changes in mice. The epididymis index, sperm count, and morphology changes in mice at 25 mg/L molybdenum were significantly different from the 100 mg/L molybdenum group (p < 0.05). The epididymis index, sperm count, and morphology changes in mice at 25 mg/L molybdenum were significantly different from the 100 mg/L molybdenum group (p < 0.05). The epididymis index, sperm count, and morphology changes in mice at 25 mg/L molybdenum were significantly different from the 100 mg/L molybdenum group (p < 0.05).



The change in epididymis index, sperm count, and morphology changes in mice at 25 mg/L molybdenum affected the sperm quality significantly.

from the 100 mg/L molybdenum group (p < 0.05). The epididymis index, sperm count, and morphology changes in mice at 25 mg/L molybdenum were significantly different from the 100 mg/L molybdenum group (p < 0.05). The epididymis index, sperm count, and morphology changes in mice at 25 mg/L molybdenum were significantly different from the 100 mg/L molybdenum group (p < 0.05).



Display full size

Table 1. Effects of molybdenum treatments on sperm parameters in mice.

Download CSV

Display Table



As shown in Table 2, compared to control, Mo at ≥ 100 mg/L decreased the activities of SOD and GPx, yet the content of MDA significantly increased. At 25 mg/L Mo markedly improved the activities of SOD and GPx, but did not change MDA. At a concentration that ranged from 12.5 and 50 mg/L only GPx activity decreased significantly. The level of SOD and MDA did not markedly change.

Table
and M

Downlo

×



The resu
acco
doses of
the 25 m
damage

Oral adn
a signific
well as a

are
us oral
eased within
nd oxidative

as shown by
count as
s probably

oxidative stress as shown by a significant decrease in the activity of both SOD and GPx, and a considerable increase in MDA in testicular tissue (Table 2). The effects of Mo on reproductive improvement have been described in several in vitro studies. Our previous study showed that at 5 µg/ml Mo is likely to improve the development of mouse embryos cultured in vitro [Bi et al. 2012]. In contrast, Braydich-Stolle et al. [2005] observed that in vitro 5 µg/ml and 10 µg/ml of Mo nano-particles seem to promote plasma membrane leakage of mouse spermatogonial stem cell lines. Molybdenum at ≥100 mg/L negatively impacted sperm quality and increased the oxidative damage in testicular tissue. There is little information in the literature on the in vivo effect of Mo on male mouse reproductive parameters. However, similar phenomena have been observed in other animals and humans. Pandey and Singh [2002] reported a dose-dependent degeneration of testicular morphology and function with declining sperm concentration, motility, normal morphology, and epididymides in rats after oral administration of sodium molybdate at a dose level of ≥ 30 mg/kg body weight. Similarly, Lyubimov et al. [2004] observed that a significant reduction of epididymal weight, sperm count, motility, morphologic abnormalities, and histopathologic changes in testis and epididymis occurred in the rats treated by tetrathiomolybdate at 12 mg/kg/day for 2 months. Bersényi et al. [2008] revealed a reduction in the number of germ cells and mature spermatocytes in the testes, and an appearance of a large number of syncytial giant cells and degenerated cells among the spermatogenic cells in

the sem
compare
depende
In additi
calves fe
lack of li
interstiti
The r
epidid
oxidative
and an i
damage
that a hi
oxidative
decrease

matter as
found dose-
in humans.
port in bull
d a complete
mage of the
the testes.
the
ed through
Px level,
ative
dicating
y increased
significant
in the



oxygen and hydrogen peroxide to protect the structures and functions of cell membranes from the interference and damage by peroxides is apparent [Ola-Mudathir et al. [2008](#); Portugal-Cohen et al. [2010](#)]. Similar phenomena were observed in rabbits. Bersényi et al. [2008] observed that high dietary Mo (39 mg Mo/kg dry matter) can generate free radicals or reactive intermediates, resulting in altering MDA and GPx activity.

In conclusion, molybdenum affects sperm quality through regulating the testicular oxidative stress in a complex manner. Male reproductive parameters apparently improved at moderate doses (25 mg/L), but were significantly repressed at high doses (≥ 100 mg/L). The change in the levels of SOD, GPx, and MDA indicate that the dual functions of Mo on sperm quality are likely to be mediated through oxidative stress in testicular tissue.

Materials and Methods

Chemicals

Unless otherwise stated, all components used in the present study were procured from Sigma-Aldrich Corp. (St. Louis, MO, USA).

Animal

All of the
Henan U
male mi
in this st
mice ha



mittee of
eks of age)
All mice used
ons. The
ellet diet.

Exp



A sub-ac
proven f
different
the Mo (
200 mg/

in mice of
he mice in
containing
0, 100, or
and

Collection of testes and epididymides

The testicular tissues of mice from each group were used for determination of SOD, GPx, and MDA, and epididymides for collecting the sperm. The mice were sacrificed by cervical dislocation on day 14 of the experiment. Testes and epididymides were quickly removed and weighed. The epididymides were placed into 37°C preheated saline, the cauda epididymis was lacerated for incubation of sperm. The testes were put in 4°C precooled saline in a refrigerator, then transferred into -20°C before homogenate preparation.

Evaluation of sperm parameters

Semen samples were collected after incubation for 30 min, and semen analysis was conducted following the World Health Organization protocol [WHO 1999]. Sperm concentration (million sperm per milliliter), percent motile sperm, and sperm morphology were investigated in this study. The concentration of immobilized sperm was determined on a hemacytometer. Sperm motility was evaluated within 1 hr after collection. Percent motile was the sum of the percentages with rapid linear progression (3 to ≥ 4) and slow linear progression (≥ 2). Sperm morphology (percent normal forms) was determined using air-dried smears stained with a modified Wright-Giemsa stain. At least 200 sperm in four different areas of the slide were evaluated according to Kruger's strict criteria [Kruger et al. 1988].

Detection

The testis was weighed and homogenized in 1 mL of precooled 0.9% saline. The homogenate was centrifuged at 1000g for 5 min. The supernatant was used for the assay of SOD, GPx, and MDA. The assay was performed according to the method of Jiancheng [1984].



Statistical analysis

Five replicates were performed for each group. The data are expressed as the mean \pm SD. The statistical analysis was performed by single factor ANOVA. The differences between groups were considered significant at $P < 0.01$.

Acknowledgments

Dr. Chunjie Zhang, Dr. Yinju Li, and Dr. Yumei Liu are thanked for assistance in usage of instruments. Ms. Kerry Alicia Webber is thanked for assistance in preparing the manuscript.

Declaration of interest: This work was supported by the National Natural Science Foundation of China (Project No. 30901026) and the Higher Education Finance Scheme for Young Backbone Teachers in Henan Province (Project No. 2010GGJS-071), China. The authors report no declarations of interest.

Author contributions: Conceived and designed the experiments: F-JL, Z-JY, X-WZ, Y-LZ; Performed the experiments: X-WZ, QQ, YB; Analyzed the data: X-LC, L-JJ, X-GM; Contributed reagents/materials/analysis tools: RS; Wrote the manuscript: F-JL, X-WZ, Y-LZ, RS.

Related Research Data

Mine

Alber

Sour

Linki



Ref










1. ACGI

1996

biolog

Goog

2. Agarw

3. Aitken, J. and Fisher, H. (1994) Reactive oxygen species generation and human spermatozoa: the balance of benefit and risk. *Bioessays* 16:259-267.
 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
4. Aitken, R.J., Clarkson, J.S. and Fishel, S. (1989) Generation of reactive oxygen species, lipid peroxidation, and human sperm function. *Biol Reprod* 41:183-197.
 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
5. Bersényi, A., Berta, E., Kádár, I., Glávits, R., Szilágyi, M. and Fekete, S.G. (2008) Effects of high dietary molybdenum in rabbits. *Acta Vet Hung* 56:41-55.
 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
6. Bi, C.M., Zhang, Y.L., Liu, F.J., Zhou, T.Z., Yang, Z.J., Gao, S.Y. (2013) The effect of molybdenum on the in vitro development of mouse preimplantation embryos. *Syst Biol Reprod Med* 59:69-73.
 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
7. Braydich-Stolle, L., Hussain, S., Schlager, J.I. and Hofmann, M.C. (2005) In vitro cytotoxicity of molybdenum. *Toxicol Sci* 88:412-419.
 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
8. Carlse, J. (2005) Molybdenum: a review of its role in human health and its potential for toxicity. *Toxicol Lett* 162:1-10.
 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
9. CDC (2005) Molybdenum. *Report on the Toxicology and Carcinogenesis Studies of Molybdenum Trioxide in F344/N Rats and B6C3F1 Mice*. [Google Scholar](#)
10. Ema, M. (2005) Molybdenum: a review of its role in human health and its potential for toxicity. *Toxicol Lett* 162:1-10.
 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)



1. Friedmann, A.S. (2002) Atrazine inhibition of testosterone production in rat males following peripubertal exposure. *Reprod Toxicol* 16:275–279.

2. Hille, R., Rétey, J., Bartlewski-Hof, U. and Reichenbecher, W. (1998) Mechanistic aspects of molybdenum-containing enzymes. *FEMS Microbiol Rev* 22:489–501.

3. Huang, J., Wu, J., Li, T., Song, X., Zhang, B., Zhang, P. (2011) Effect of exposure to trace elements in the soil on the prevalence of neural tube defects in a high-risk area of China. *Biomed Environ Sci* 24:94–101.

4. Kargar, M., Khorasani, N., Karami, M., Rafiee, G.R. and Naseh, R. (2011) Study of aluminum, copper and molybdenum pollution in groundwater sources surrounding (Miduk) Shahr-E-Babak copper complex tailings dam. *World Academy of Science, Engineering and Technology* 76:412–416.

5. Khura... in male and female... etinemia and altera... 7.

6. Kruge... nninger, S. (1... tion. Fertil Steril

7. Lyubin... J.E., Smith, A.C., (opper pro... pro... Toxicol



8. Makker, K., Agarwal, A. and Sharma, R. (2009) Oxidative stress & male infertility. Indian J Med Res 129:357-367

PubMed | Google Scholar

9. Meeker, J.D., Rossano, M.G., Protas, B., Diamond, M.P., Puscheck, E., Daly, D. (2008) Cadmium, lead, and other metals in relation to semen quality: human evidence for molybdenum as a male reproductive toxicant. Environ Health Perspect 116:1473-1479.

PubMed | Web of Science ® | Google Scholar

10. Mills, C.F. and Davis, G.K. (1987) Molybdenum, In: Trace Elements in Human and Animal Nutrition, 5th edn, ed. Mertz, W., Academic Press, San Diego, CA pp. 429-463

Google Scholar

11. Ola-Mudathir, K.F., Suru, S.M., Fafunso, M.A., Obioha, U.E. and Faremi, T.Y. (2008) Protective roles of onion and garlic extracts on cadmium-induced changes in sperm characteristics and testicular oxidative damage in rats. Food Chem Toxicol 46:3604-3611.

12. Pande... male rats.
Biome...

13. Portu... ces
ox... J Dermatol
Sci 50

14. Schroe... on the
repro...



25. Sharma, S., Kaur, R. and Sandhu, H.S. (2004) Effect of subacute oral toxicity of molybdenum on antioxidant status in cross bred cow calves. *Ind J Anim Sci* 74:734-736.

[Google Scholar](#)

26. Spalj, S., Mlacovic Zrinski, M., Tudor Spalj, V. and Ivankovic Buljan, Z. (2012) In-vitro assessment of oxidative stress generated by orthodontic archwires. *Am J Orthod Dentofacial Orthop* 141:583-589.

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

27. Thomas, J.W. and Moos, S. (1951) The effect of orally administered molybdenum on growth spermatogenesis and testes histology of young dairy bulls. *J Dairy Sci* 34:929-934.

[Web of Science ®](#) | [Google Scholar](#)

28. Underwood, E.J. (1981) Trace metals in human and animal health. *J Hum Nutr* 35:37-48.

[PubMed](#) | [Google Scholar](#)

29. WHO (World Health Organization) (1993) Guidelines for drinking water quality.

Second

[Goog](#)

30. WHO (World Health Organization) (1996) Guidelines for drinking water quality. Examination and testing of water. Geneva: WHO. New York: CRC Press.

York: CRC

[Goog](#)

31. Wira, D. (1996) The effect of molybdenum exposure on male reproductive health. *Environ Health Perspect* 104:11-15.

32. Yu, C., et al. (2000) Molybdenum exposure and its effects on human health: a review. *Environ Health Perspect* 108:11-15.

Download PDF

Related research

People also read

Recommended articles

Cited by
39



Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research
- Help and information
- Help and contact
- Newsroom
- All journals
- Books

Copyright

Accessib

Registered
5 Howick Pl

or & Francis Group
orma business

