



Free access

2,406 41 15  
Views CrossRef citations to date Altmetric

Listen

RESEARCH COMMUNICATIONS

# Effects of molybdenum on sperm quality and testis oxidative stress

Xiao-Wei Zhai, Yu-Ling Zhang, Qiao Qi, Yu Bai, Xiao-Li Chen, Li-Jun Jin, ... show all

Pages 251-255 | Received 23 Nov 2012, Accepted 12 Feb 2013, Published online: 08 May 2013

Cite this article <https://doi.org/10.3109/19396368.2013.791347>

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

View PDF

Share

## We Care About Your Privacy

We and our 899 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting "I Accept" enables tracking technologies to support the purposes shown under "we and our partners process data to provide," whereas selecting "Reject All" or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the ["privacy preferences"] link on the bottom of the webpage [or the floating icon on the bottom-left of the webpage, if applicable]. Your choices will have effect within our Website. For more details, refer to our Privacy Policy. [Here](#)

We and our partners process data to provide:

...

I Accept

Reject All

Show Purpose



# Introduction

A variety of endocrine disrupting chemicals have been released into the environment in the rapid industrial progress, which may exert adverse health effects in human and animals [Carlsen et al. [1992](#); Khurana et al. [2000](#); Friedmann [2002](#)]. Molybdenum (Mo) is an essential trace element in animals and humans. It has been identified as part of the active sites of over 50 enzymes, and may promote normal cell function possibly by catalyzing a variety of hydroxylation, oxygen atom transfer and other oxidation-reduction reactions [Hille et al. [1998](#)]. Molybdenum is also an endocrine disruptor and has been widely present and detected in our food and water [Underwood [1981](#); Mills and Davis [1987](#); Kargar et al. [2011](#); Yu et al. [2011](#)]. In addition, Mo is broadly used in industrial production, such as metallurgical processes, the manufacture of electronic products, glass, ceramics, lubricants, catalysts, pigments and nano materials [Pandey and Singh [2002](#); CDC 2005; Braydich-Stolle et al. [2005](#); Ema et al. [2010](#)]. Furthermore, Mo is also an environmental pollutant discharged from uranium processing, combustion processing, contact lens solutions, and the color additives in cosmetics [ACGIH 1995]. This wide distribution greatly increases the risk of animals and humans exposed to the high level of Mo in the environment. For example, molybdenum concentrations have risen to a maximum of 100 µg/L in drinking water in some areas of the world [Bersényi et al. [2008](#)].

Excessive exposure to Mo can cause adverse health effects in animals and humans. In rats, high-dose Mo exposure caused morphologic abnormalities in the reproductive system [Pandey and Singh [2002](#)]. A reduction in the number of spermatozoa was observed in male rats [Bersényi et al. [2008](#)]. In humans, Mo exposure has been associated with Meeker et al. [2008](#)]. However, the effects of Mo on male reproductive system are still unclear.

Oxidative stress is a key factor in the development of reproductive deformity, and even infertility. In this study, we investigated the effects of Mo on male reproductive system and the underlying mechanism. We showed that Mo induced oxidative stress and damaged the reproductive system of male rats.



titanium-molybdenum generated oxidative stress in mouse fibroblasts cultures.

Antioxidants provide a defense against oxidative stress.

Seminal plasma contains three main enzymatic antioxidants: superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPx). Spermatozoa possess primarily enzymatic antioxidants, with SOD being the most predominant [Makker et al. [2009](#)]. Superoxide dismutase in conjugation with CAT and GPx scavenge both intracellular and extracellular superoxide radicals and inhibit lipid peroxidation [Agarwal and Prabhakaran [2005](#)]. Malondialdehyde (MDA) is one of the byproducts of lipid peroxidation that indirectly reflects the level of peroxidation and the degree of cell injury [Sharma et al. [2004](#)]. This byproduct has been used in various biochemical assays to monitor the degree of oxidative damage sustained by spermatozoa [Aitken et al. [1989](#); Aitken and Fisher [1994](#)]. Oxidative stress can be evaluated by detecting the activities of SOD and GPx, as well as the MDA level in the tissue. Up to now, there are no data available for the oxidative stress of testicular tissue caused by Mo on mice.


This study has been undertaken to evaluate the effect of orally administered Mo on the sperm parameters of the epididymis index, sperm motility, count, and morphology changes. The oxidative stress of testis was considered as a function of the levels of MDA, SOD, and GPx in mice.

## Result

The epididymis index and sperm count in morphological parameters of mice treated with 25 mg/L molybdenum were significantly lower than those of the control group.

Figure 1. The change in morphological parameters of mice treated with 25 mg/L molybdenum (A) epididymis index; B) sperm count; C) sperm motility; D) sperm morphology. The change in morphological parameters of mice treated with 25 mg/L molybdenum at 25 mg/L molybdenum affected the sperm count and motility significantly.





Display full size

Table 1. Effects of molybdenum treatments on sperm parameters in mice.

Download CSV   Display Table

As shown in Table 2, compared to control, Mo at  $\geq 100$  mg/L decreased the activities of SOD and GPx, yet the content of MDA significantly increased. At 25 mg/L Mo markedly improved the activities of SOD and GPx, but did not change MDA. At a concentration that ranged from 12.5 and 50 mg/L only GPx activity decreased significantly. The level of SOD and MDA did not markedly change.

Table  
and M

Downlo

The re  
acco  
doses of  
the 25 m  
damage  
  
Oral adn  
a signific  
well as a

are  
us oral  
eased within  
nd oxidative  
  
as shown by  
count as  
s probably

oxidative stress as shown by a significant decrease in the activity of both SOD and GPx, and a considerable increase in MDA in testicular tissue (Table 2). The effects of Mo on reproductive improvement have been described in several in vitro studies. Our previous study showed that at 5 µg/ml Mo is likely to improve the development of mouse embryos cultured in vitro [Bi et al. 2012]. In contrast, Braydich-Stolle et al. [2005] observed that in vitro 5 µg/ml and 10 µg/ml of Mo nano-particles seem to promote plasma membrane leakage of mouse spermatogonial stem cell lines. Molybdenum at  $\geq 100$  mg/L negatively impacted sperm quality and increased the oxidative damage in testicular tissue. There is little information in the literature on the in vivo effect of Mo on male mouse reproductive parameters. However, similar phenomena have been observed in other animals and humans. Pandey and Singh [2002] reported a dose-dependent degeneration of testicular morphology and function with declining sperm concentration, motility, normal morphology, and epididymides in rats after oral administration of sodium molybdate at a dose level of  $\geq 30$  mg/kg body weight. Similarly, Lyubimov et al. [2004] observed that a significant reduction of epididymal weight, sperm count, motility, morphologic abnormalities, and histopathologic changes in testis and epididymis occurred in the rats treated by tetrathiomolybdate at 12 mg/kg/day for 2 months. Bersényi et al. [2008] revealed a reduction in the number of germ cells and mature spermatocytes in the testes, and an appearance of a large number of syncytial giant cells and degenerated cells among the spermatogenic cells in the seminiferous tubules. In addition, Pandey and Singh [2002] found dose-dependent degeneration of testicular morphology and function in humans. In addition, Pandey and Singh [2002] reported a complete degeneration of the testes in bull calves fed with sodium molybdate at a dose level of  $\geq 30$  mg/kg body weight. The lack of lipid peroxidation in the testes of the rats treated by tetrathiomolybdate at 12 mg/kg/day for 2 months. The interstitial cells of Leydig in the testes of the rats treated by tetrathiomolybdate at 12 mg/kg/day for 2 months. The epididymis of the rats treated by tetrathiomolybdate at 12 mg/kg/day for 2 months. The oxidative damage in the testes of the rats treated by tetrathiomolybdate at 12 mg/kg/day for 2 months. and an increase in the activity of SOD and GPx in the testes of the rats treated by tetrathiomolybdate at 12 mg/kg/day for 2 months. damage in the testes of the rats treated by tetrathiomolybdate at 12 mg/kg/day for 2 months. that a high dose of Mo can cause oxidative damage in the testes of the rats treated by tetrathiomolybdate at 12 mg/kg/day for 2 months. oxidative damage in the testes of the rats treated by tetrathiomolybdate at 12 mg/kg/day for 2 months. decrease in the activity of SOD and GPx in the testes of the rats treated by tetrathiomolybdate at 12 mg/kg/day for 2 months.



oxygen and hydrogen peroxide to protect the structures and functions of cell membranes from the interference and damage by peroxides is apparent [Ola-Mudathir et al. [2008](#); Portugal-Cohen et al. [2010](#)]. Similar phenomena were observed in rabbits. Bersényi et al. [2008] observed that high dietary Mo (39 mg Mo/kg dry matter) can generate free radicals or reactive intermediates, resulting in altering MDA and GPx activity.

In conclusion, molybdenum affects sperm quality through regulating the testicular oxidative stress in a complex manner. Male reproductive parameters apparently improved at moderate doses (25 mg/L), but were significantly repressed at high doses ( $\geq 100$  mg/L). The change in the levels of SOD, GPx, and MDA indicate that the dual functions of Mo on sperm quality are likely to be mediated through oxidative stress in testicular tissue.

## Materials and Methods

### Chemicals

Unless otherwise stated, all components used in the present study were procured from Sigma-Aldrich Corp. (St. Louis, MO, USA).

### Animal

All of the experiments were approved by the Animal Care and Use Committee of Henan University. The mice were 6-8 weeks of age. All mice used in this study were male. The mice were housed in cages with a 12 h light/dark cycle and a pellet diet.

### Expe

A sub-acute study was conducted in mice of 6-8 weeks of age. The mice in this study were divided into four groups containing 10, 100, or 200 mg/L of Mo. The mice were housed in cages with a 12 h light/dark cycle and a pellet diet.

## Collection of testes and epididymides

The testicular tissues of mice from each group were used for determination of SOD, GPx, and MDA, and epididymides for collecting the sperm. The mice were sacrificed by cervical dislocation on day 14 of the experiment. Testes and epididymides were quickly removed and weighed. The epididymides were placed into 37°C preheated saline, the cauda epididymis was lacerated for incubation of sperm. The testes were put in 4°C precooled saline in a refrigerator, then transferred into -20°C before homogenate preparation.

## Evaluation of sperm parameters

Semen samples were collected after incubation for 30 min, and semen analysis was conducted following the World Health Organization protocol [WHO 1999]. Sperm concentration (million sperm per milliliter), percent motile sperm, and sperm morphology were investigated in this study. The concentration of immobilized sperm was determined on a hemacytometer. Sperm motility was evaluated within 1 hr after collection. Percent motile was the sum of the percentages with rapid linear progression (3 to  $\geq 4$ ) and slow linear progression ( $\geq 2$ ). Sperm morphology (percent normal forms) was determined using air-dried smears stained with a modified Wright-Giemsa stain. At least 200 sperm in four different areas of the slide were evaluated according to Kruger's strict cri

# Detecti

The test was performed using pre-cooled 0.9% saline homogenate for the assay of SOD, Jiancheng

# Statisti

Five replicates were performed for each treatment and the mean values were calculated. The data were analyzed by single factor ANOVA. The differences between treatments were considered significant at  $P < 0.01$ .




# Acknowledgments

Dr. Chunjie Zhang, Dr. Yinju Li, and Dr. Yumei Liu are thanked for assistance in usage of instruments. Ms. Kerry Alicia Webber is thanked for assistance in preparing the manuscript.

Declaration of interest: This work was supported by the National Natural Science Foundation of China (Project No. 30901026) and the Higher Education Finance Scheme for Young Backbone Teachers in Henan Province (Project No. 2010GGJS-071), China. The authors report no declarations of interest.

Author contributions: Conceived and designed the experiments: F-JL, Z-JY, X-WZ, Y-LZ; Performed the experiments: X-WZ, QQ, YB; Analyzed the data: X-LC, L-JJ, X-GM; Contributed reagents/materials/analysis tools: RS; Wrote the manuscript: F-JL, X-WZ, Y-LZ, RS.



## Related Research Data

Effect

Source

Cadr

Evide

Source

The I

and T

S

Trace

Source

Evide

Source

Moly

Source



esis



Reproductive and developmental toxicity studies of manufactured nanomaterials

Source: Reproductive Toxicology

Mechanistic aspects of molybdenum-containing enzymes

Source: FEMS Microbiology Reviews

Reprint of: Predictive value of abnormal sperm morphology in in vitro fertilization

Source: Fertility and Sterility

In-vitro assessment of oxidative stress generated by orthodontic archwires

Source: American Journal of Orthodontics and Dentofacial Orthopedics

World Health Organization Laboratory Manual for the Examination of Human Semen and Sperm-Cervical Mucus Interaction, 4th ed.

Source: Journal of Andrology

Effect of exposure to trace elements in the soil on the prevalence of neural tube defects in a high-risk area of China.

Source: Biomedical and Environmental Sciences

In Vitro Cytotoxicity of Nanoparticles in Mammalian Germline Stem Cells

Source: Toxicological Sciences

Toxic Effects of Trace Elements on the Reproduction of Mice and Rats

Source: Archives of Environmental Health An International Journal

Gene × Sperm

Func

Sour

Expo

Delat

Rece

1015

th

at

2000

Sour

Coca

oxide

Sour

Prote

Sperm

, ES-

from

sented

June

ic

es in



Source: Food and Chemical Toxicology

Reactive oxygen species generation and human spermatozoa: The balance of benefit and risk

Source: BioEssays

The effect of molybdenum on the in vitro development of mouse preimplantation embryos

Source: Systems Biology in Reproductive Medicine

Effects of high dietary molybdenum in rabbits

Source: Acta Veterinaria Hungarica

Adverse Effects of Low Level Heavy Metal Exposure on Male Reproductive Function

Source: Systems Biology in Reproductive Medicine

Atrazine inhibition of testosterone production in rat males following peripubertal exposure

Source: Reproductive Toxicology

The effects of tetrathiomolybdate (TTM, NSC-714598) and copper supplementation on fertility and early embryonic development in rats

Source: Reproductive Toxicology

Predictive value of abnormal sperm morphology in in vitro fertilization

Source: Fertility and Sterility







Link




## Refer

1. ACGIH (1995) 1995-1996 Threshold Limit Values for Chemical Agents and Biological Agents. Cincinnati, OH: American Conference of Governmental Industrial Hygienists. Google Scholar
2. Agarwal, A.K., et al. (1994) Prevention of oxidative damage to spermatozoa by antioxidants. J. Androl. 15: 974. PubMed



3. Aitken, J. and Fisher, H. (1994) Reactive oxygen species generation and human spermatozoa: the balance of benefit and risk. *Bioessays* 16:259–267.
-  | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
4. Aitken, R.J., Clarkson, J.S. and Fishel, S. (1989) Generation of reactive oxygen species, lipid peroxidation, and human sperm function. *Biol Reprod* 41:183–197.
-  | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
5. Bersényi, A., Berta, E., Kádár, I., Glávits, R., Szilágyi, M. and Fekete, S.G. (2008) Effects of high dietary molybdenum in rabbits. *Acta Vet Hung* 56:41–55.
-  | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
6. Bi, C.M., Zhang, Y.L., Liu, F.J., Zhou, T.Z., Yang, Z.J., Gao, S.Y. (2013) The effect of molybdenum on the in vitro development of mouse preimplantation embryos. *Syst Biol Reprod Med* 59:69–73.
-  | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
7. Braydich-Stella, L., Hussain, S., Schlager, L. and Hefeman, M.C. (2005) In vitro cytotoxicity of molybdenum compounds. *Toxicol Appl Pharmacol* 88:412–419.
-  | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
8. Carlsson, L. and Carlsson, L. (2005) Molybdenum: a review of its role for human health. *Ann NY Acad Sci* 1062:1–10.
-  | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
9. CDC (Centers for Disease Control and Prevention) (2005) Molybdenum. *Report on the status of the periodic table of elements*. [Google Scholar](#)
10. Ema, P. (2005) Molybdenum: a review of its role for human health. *Ann NY Acad Sci* 1062:1–10.






[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

1. Friedmann, A.S. (2002) Atrazine inhibition of testosterone production in rat males following peripubertal exposure. *Reprod Toxicol* 16:275-279.




[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

2. Hille, R., Rétey, J., Bartlewski-Hof, U. and Reichenbecher, W. (1998) Mechanistic aspects of molybdenum-containing enzymes. *FEMS Microbiol Rev* 22:489-501.



[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

3. Huang, J., Wu, J., Li, T., Song, X., Zhang, B., Zhang, P. (2011) Effect of exposure to trace elements in the soil on the prevalence of neural tube defects in a high-risk area of China. *Biomed Environ Sci* 24:94-101.

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

4. Kargar, M., Khorasani, N., Karami, M., Rafiee, G.R. and Naseh, R. (2011) Study of aluminum, copper and molybdenum pollution in groundwater sources surrounding (Midul

*Engine*


[Google](#)

5. Khura... in male and female... etinemia and alter... 7.



6. Kruge... nninger, S. (1988) ... tion. *Fertil Steril*


7. Lyubimov, A.V., Smith, J.A., Rousselle, S.D., Mercieca, M.D., Tomaszewski, J.E., Smith, A.C., (2004) The effects of tetrathiomolybdate (TTM, NSC-714598) and copper supplementation on fertility and early embryonic development in rats. *Reprod Toxicol* 19:223-233.

[PubMed](#)[Web of Science ®](#)[Google Scholar](#)


8. Makker, K., Agarwal, A. and Sharma, R. (2009) Oxidative stress & male infertility. *Indian J Med Res* 129:357-367

[PubMed](#)[Google Scholar](#)

9. Meeker, J.D., Rossano, M.G., Protas, B., Diamond, M.P., Puschek, E., Daly, D. (2008) Cadmium, lead, and other metals in relation to semen quality: human evidence for molybdenum as a male reproductive toxicant. *Environ Health Perspect* 116:1473-1479.

[PubMed](#)[Web of Science ®](#)[Google Scholar](#)


10. Mills, C.F. and Davis, G.K. (1987) Molybdenum, In: *Trace Elements in Human and Animal Nutrition*, 5th edn, ed. Mertz, W., Academic Press, San Diego, CA pp. 429-463



11. Ola-M... (2008)  
Protect... s in sperm  
charac... ol 46:3604-  
3611.



12. Pande... ale rats.  
Biome...



13. Portug... ces  
evidet... Dermatol

[PubMed](#)[Web of Science ®](#)[Google Scholar](#)

24. Schroeder, H.A. and Mitchener, M. (1971) Toxic effects of trace elements on the reproduction of mice and rats. *Arch Environ Health* 23:102–106.

[PubMed](#)[Web of Science ®](#)[Google Scholar](#)

25. Sharma, S., Kaur, R. and Sandhu, H.S. (2004) Effect of subacute oral toxicity of molybdenum on antioxidant status in cross bred cow calves. *Ind J Anim Sci* 74:734–736.

[Google Scholar](#)

26. Spalj, S., Mlacovic Zrinski, M., Tudor Spalj, V. and Ivankovic Buljan, Z. (2012) In-vitro assessment of oxidative stress generated by orthodontic archwires. *Am J Orthod Dentofacial Orthop* 141:583–589.

[PubMed](#)[Web of Science ®](#)[Google Scholar](#)

27. Thomas, J.W. and Moos, S. (1951) The effect of orally administered molybdenum on growth spermatogenesis and testes histology of young dairy bulls. *J Dairy Sci* 34:929–934.

28. Underwood EJ (1976) Molybdenum deficiency in sheep. *Nutr* 35:37–48.

[PubMed](#)

29. WHO (1996) *World Health Statistics Quarterly*. Second Report. Geneva: WHO.

[Google Scholar](#)

30. WHO (1996) *World Health Statistics Quarterly*. Second Report. Geneva: WHO.

31. Wirth, J.J. and Mijal, R.S. (2010) Adverse effects of low level heavy metal exposure on male reproductive function. Syst Biol Reprod Med 56:147–167.

PubMed

Web of Science ®

Google Scholar

32. Yu, C., Xu, S., Gang, M., Chen, G. and Zhou, L. (2011) Molybdenum pollution and speciation in Nver River sediments impacted with Mo mining activities in western Liaoning, northeast China. Int J Environ Res 5:205–212.

Web of Science ®

Google Scholar

Download PDF

Related research ⓘ

People also read

Recommended articles

Cited by 41

Dual eff...

Yu-Ling Systems Publishe



## Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

## Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

## Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research
- Help and information
- Help and contact
- Newsroom
- All journals
- Books

## Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright © 2024 John Wiley & Sons, Inc. All rights reserved. Wiley, Wiley Online Library, and Wiley InterScience are trademarks of John Wiley & Sons, Inc. or John Wiley & Sons Group, Inc. or one of its subsidiaries. All other trademarks are the property of their respective owners. For more information, please visit [wiley.com/termsandconditions](#).

Accessibility

Registered  
5 Howick Place

