

Autophagy >

Volume 8, 2012 - Issue 6

Free access

1,184 Views | 22 CrossRef citations to date | 0 Altmetric

Listen

Autophagic Punctum

Long time-lapse imaging reveals unique features of PARK2/Parkin-mediated mitophagy in mature cortical neurons

Qian Cai , Hesham Mostafa Zakaria & Zu-Hang Sheng

Pages 976-978 | Published online: 01 Jun 2012

Cite this article <https://doi.org/10.4161/auto.20218>

Full Article

Figures & data

Citations

Metrics

Reprints & Permissions

View PDF

Share

We Care About Your Privacy

We and our 880 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting I Accept enables tracking technologies to support the purposes shown under we and our partners process data to provide. Selecting Reject All or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the Show Purposes link on the bottom of the webpage. Your choices will have effect within our Website. For more details, refer to our Privacy Policy. [Here](#)

We and our partners process data to provide:

Use precise geolocation data. Actively scan device

I Accept


Reject All

Show Purpose



This article refers to:

Mitochondria are essential organelles for neuronal function, development and survival. Throughout a neuron's lifetime, aged and damaged mitochondria undergo dynamic recycling via fusion/fission or are ultimately eliminated via mitophagy, an autophagic pathway specific for mitochondrial degradation. Dysfunctional mitochondria not only produce energy less efficiently, but also release harmful reactive oxygen species and initiate apoptotic signaling cascades, which have been linked to the pathogenesis of several major neurodegenerative diseases. Proper sequestration of damaged mitochondria into autophagosomes and subsequent degradation within the lysosomal system constitute a key cellular pathway in mitochondrial quality control mechanisms. Mutations in PINK1 and PARK2 are linked to autosomal recessive early onset Parkinson disease. Recent studies demonstrate that the PINK1-PARK2 pathway mediates mitophagy, and that PARK2 translocation onto mitochondria induces mitochondrial degradation via mitophagy in several non-neuronal cell types where PARK2 is overexpressed, such as in HEK293, HeLa, MEF and neuroblastoma cells. However, evidence of PARK2 translocation onto mitochondria is missing or controversial. Mitophagy has some unique features that are not addressed by conventional autophagy (1) whether this translocation of PARK2 onto mitochondrial mobility is targeted to the autophagosome. First, we observed that mitochondrial membrane potential is depolarized in mitophagy. We established that PARK2 translocation onto mitochondria is such that they survive long enough to exhibit PARK2 translocation. CCCP treatment for 24 h



induces 26.67% of neurons to undergo PARK2 translocation onto depolarized mitochondria. Co-treatment with CCCP and lysosomal inhibitors (LIs) results in a doubling in the percentage (55.87%) of neurons with PARK2 translocation. Thus, proper lysosomal function is critical to avoid the accumulation of PARK2-associated mitochondria in neurons upon dissipating $\Delta\psi_m$. Second, we examined PARK2 translocation kinetics by imaging neurons at various time points during CCCP treatment. PARK2 translocation between 0.5–6 h was exceptionally rare, occasionally observed as early as 12 h, and became increasingly frequent after 18 h of CCCP treatment. We further imaged the dynamic recruitment of endogenous PARK2 to mitochondria in mature neurons following CCCP/LIs treatment. To confirm this, we alternatively isolated the mitochondria-enriched membrane fraction from mature cortical neurons following the same treatment. While the majority of PARK2 is present in the cytosolic fraction, mitochondrial depolarization results in the association of substantial levels of endogenous PARK2, along with autophagic markers LC3-II and SQSTM1/p62 with the mitochondrial membrane. These results consistently indicate the recruitment of both endogenous and exogenous PARK2 onto depolarized mitochondria in mature cortical neurons.

Next, we utilized a $\Delta\psi_m$ -sensitive dye, TMRE, to examine whether PARK2-targeted mitochondria display dissipated $\Delta\psi_m$. Healthy mitochondria will sequester and accumulate TMRE, thus resulting in a higher TMRE fluorescence intensity. In control

neurons, electrocatalytic targets that required impairs mitochondria that mitochondria. Intriguingly, mitochondria axons and translocated retrograde be attributed to the unique distribution pattern. PARK2-tagged depolarized




mitochondria are restricted to the soma for degradation, where lysosomes are predominately located, while healthy mitochondria are distributed distally to support synaptic functions. To test this hypothesis, we utilized syntaphilin, an axonal mitochondria docking protein, to artificially immobilize mitochondria in distal processes. To our surprise, we found that PARK2 is recruited to stationary mitochondria anchored by syntaphilin in distal processes. Therefore, the PARK2-mediated process prevents dysfunctional mitochondria from traveling peripherally, leading to their accumulation in somatodendritic regions. In addition, our long time-lapse imaging exhibits dynamic PARK2 recruitment and degradation of depolarized mitochondria within the autophagy-lysosomal system. CCCP exposure results in LC3-labeled ring-like structures surrounding fragmented mitochondria in the somadendritic regions and enhances the recruitment of mitochondria to lysosomes for degradation in live cortical neurons.

In summary, our study reveals several unique features of PARK2-mediated mitophagy in mature cortical neurons. First, PARK2 is selectively recruited to depolarized mitochondria to form ring-like structures, a process occurring more slowly than in non-neuronal cells. Second, following 24 h CCCP incubation, PARK2 translocation only occurs in a small percentage of neurons. Third, PARK2 translocation is restricted to the somatodendritic regions, where mature lysosomes are predominantly located. This spatial and dynamic process allows neurons to efficiently eliminate dysfunctional mitochondria via the autophagy-lysosomal pathway (Fig. 1).

Figure 1.  control via
PARK2-mitochondria
to form a
dendritic
dendritic
of mitoc
Altered
health
targeted
mature l
neurons
pathway
dysfunct

control via
mitochondria
proximal
distal
al transport
ased.
while
PARK2-
ions, where
cess allows
lysosomal
mination of





Display full size

Acknowledgments

This work was supported by the NIH Program

(Z.-H.S.),
Scholars



Download



Related

Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright

Accessib

Registered
5 Howick Pl

or & Francis Group
orma business

