

Autophagy > Volume 8, 2012 - Issue 10

Free access

929 Views | 17 CrossRef citations to date | 1 Altmetric

Listen

Autophagic Punctum

Curbing autophagy and histone deacetylases to kill cancer cells

Noor Gammoh, Paul A. Marks & Xuejun Jiang

Pages 1521-1522 | Received 01 Jun 2012, Accepted 18 Jun 2012, Published online: 16 Aug 2012

Cite this article <https://doi.org/10.4161/auto.21151>

Full Article

Figures & data

Citations

Metrics

Reprints & Permissions

View PDF

Abstract

Cells respond to cytotoxicity by activating a variety of signal transduction pathways. One pathway frequently upregulated during cytotoxic response is macroautophagy

(hereafter deacetylase (SA inhibition kinase U and ne auto of quest have dis kinase M what the

-histone hydroxamic HDAC phagic apoptotic potential of ed a series ual HDACs sensing autophagy, Tackling

We Care About Your Privacy

We and our 843 partners store and/or access information on a device, such as unique IDs in cookies to process personal data. You may accept or manage your choices by clicking below, including your right to object where legitimate interest is used, or at any time in the privacy policy page. These choices will be signaled to our partners and will not affect browsing data. [Privacy Policy](#)

We and our partners process data to provide:

Use precise geolocation data. Actively scan device characteristics for identification. Store and/or access information on a device. Personalised advertising and content, advertising and content measurement, audience research and services development.

List of Partners (vendors)

I Accept

Essential Only

Show Purpose



these questions should lead to a better understanding of autophagy and HDAC biology and contribute to the development of novel therapeutic strategies.

Keywords: : HDAC inhibitors autophagy apoptosis nonapoptotic cell death MTOR ULK1 SAHA


This article refers to:

HDAC expression is deregulated in a wide range of human cancer types. Targeting HDAC activity can impose antitumor effects, possibly through both chromatin-dependent and -independent mechanisms. Previous work by us and others demonstrates that HDAC inhibitors, including SAHA, can induce transcriptional upregulation of multiple genes that function to promote growth arrest or apoptosis, and downregulation of genes that may serve to facilitate cancer development. HDAC inhibitors can also enhance acetylation of various nonhistone proteins, and many of these proteins are involved in fundamental cellular processes whose malfunction can have an impact on tumorigenesis.

In addition to the induction of cell death preferentially in transformed cells, we found that HDAC inhibitors can also induce autophagy, a cellular catabolic process highly implicated in cancer development and treatment (Fig. 1). Our recent study shows that SAHA does so by suppressing the MTOR complex 1 (MTORC1), a master regulator of cellular metabolism and a therapeutic target for anticancer treatment. MTOR suppresses autophagy by phosphorylating and inactivating the ULK1 complex, an upstream regulator of autophagy. SAHA restores the function of ULK1, highlighting the importance of this pathway in inducing autophagy. ULK1 expression is upregulated in SAHA-treated cells, and its activity is restored by SAHA. Inactivation of MTOR, the primary regulator of acetylation, leads to increased acetylation of histone and nonhistone proteins, and its regulation is restored by SAHA. We demonstrate that SAHA induces autophagy when MTORC1 activity is inhibited by SAHA.



Figure 1. HDAC inhibitor-induced autophagy and cell death. HDAC inhibitors can induce autophagy by inactivating MTORC1 and consequently activating the upstream component of the autophagy pathway, the ULK1 complex. Additionally, HDAC inhibition can also lead to the transcriptional upregulation of LC3, and possibly enhanced acetylation of autophagy proteins, such as ULK1, ATG3 and ATG7. These events may further augment autophagy once autophagy is activated through the ULK1 complex. Autophagy in turn plays a survival role in attenuating both apoptotic and nonapoptotic cell death induced by HDAC inhibitors.



Display full size

Intriguingly, the recent findings that acetylation of a number of autophagy-related (Atg) gene products, such as ULK1 and ATG3, can potentiate autophagy, have provided further mechanistic insights into the role of HDACs and HDAC inhibitors in autophagy. Presumably, by deacetylating these ATG proteins, HDACs can function as negative

modulat
autoph
deacety
shuttle b
strictly c
Further
play
utilize tr
might ev
autoph
mitocho
autoph
by RNAi should be employed.

The obvious therapeutically relevant question is: Since HDAC inhibitors can trigger autophagy, what is the role of autophagy in HDAC inhibitor-induced cancer cell death? In our assays, HDAC inhibitors can induce robust caspase activity and apoptotic cell death in various cancer cells, including glioblastoma cells, which are resistant to multiple anticancer agents. We examined the effect of autophagy on SAHA-induced apoptosis in these malignant cells. Because specific inhibitors against autophagy are presently not available and all the currently used inhibitors affect both autophagy and endocytosis, we used an RNAi approach to tackle this question. Caspase activation and cell death are greatly increased when autophagy is impeded by RNAi. Because autophagy is a major survival mechanism upon stress, this result is not a surprise. However, mechanistically, how exactly autophagy protects cells from apoptosis (as well as nonapoptotic cell death as described below) is not well defined. This continues to be an important question, especially considering that under certain specific contexts, autophagy can be a cell death-promoting mechanism.

Perhaps the more interesting question concerns how SAHA triggers nonapoptotic cell death, and the role of autophagy in this death process. We found that autophagy plays a similar protective role in nonapoptotic cell death. The nature of this nonapoptotic cell death observed during combinational treatment using SAHA and the caspase inhibitor zVAD remains unknown. We have previously shown that the generation of reactive oxygen species, which is deliberately regulated by HDACs, may play a role in nonapoptotic cell death. However, our preliminary studies show that when the necrotic factor RIPK1/RIP1 kinase is inhibited using necrostatin-1, the induced nonapoptotic cell death still occurs. Therefore, is the SAHA-induced nonapoptotic cell death dependent on

other re...
death pa...
death m...
Obvious...
and the...
point...



Related research

People also read

Recommended articles

Cited by
17

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact


Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright

Access

Register
5 Howick Pl

or & Francis Group
orma business

