

[Plant Signaling & Behavior](#) >Volume 7, 2012 - [Issue 6](#) Free access

1,455

Views

24

CrossRef citations to date

1

Altmetric

[Listen](#)

Mini Review

N-rich protein (NRP)-mediated cell death signaling

A new branch of the ER stress response with implications for plant biotechnology

Pedro A.B. Reis & Elizabeth P.B. Fontes 

Pages 628-632 | Published online: 14 May 2012

 Cite this article  <https://doi.org/10.4161/psb.20111> Full Article Figures & data References Citations Metrics Reprints & Permissions View PDF Share

Abstract

Upon disruption of ER homeostasis, plant cells activate at least two branches of the unfolded protein response (UPR) through IRE1-like and ATAF6-like transducers, resulting in the upregulation of ER-resident molecular chaperones and the activation of the ER-associated degradation protein system. Here, we discuss a new ER stress response pathway in plants that is associated with an osmotic stress response in transducing a cell death signal. Both ER and osmotic stress induce the expression of the novel transcription factor GmERD15, which binds and activates N-rich protein (NRP) promoters to induce NRP expression and cause the upregulation of GmNAC6, an effector of the cell death response. In contrast to this activation mechanism, the ER-resident molecular chaperone binding protein (BiP) attenuates the propagation of the cell death signal by modulating the expression and activity of components of the ER and osmotic stress-induced NRP-mediated cell death signaling. This interaction

attenuates dehydration-induced cell death and promotes a better adaptation of BiP-overexpressing transgenic lines to drought.

Keywords: :

ERD15 GmNAC6 N-rich proteins NRP-A NRP-B cell death response endoplasmic reticulum stress osmotic stress

The endoplasmic reticulum is a key signaling organelle involved in the activation of cellular stress responses in eukaryotic cells. One such well-characterized signaling event is the unfolded protein response (UPR), which is activated to cope with the disruption of ER homeostasis that results in the accumulation of unfolded or misfolded proteins in the lumen of the organelle. In mammalian cells, the UPR has been associated with other stress response pathways through shared components.¹ In plants, the potential of the ER stress response to accommodate adaptive pathways and its connection with other environmentally induced responses have been the subjects of studies in recent years.^{2,3} One plant-specific, ER stress-shared response is ER and osmotic stress-integrated signaling, which converges on N-rich proteins (NRPs) to transduce a cell death signal.⁴ As an integrated pathway, NRP-mediated cell death signaling is activated by either ER or osmotic stress but requires both signals for full activation.

ER stress response and NRP-mediated cell death signaling

Any condition that disrupts ER homeostasis to induce the accumulation of unfolded or misfolded proteins in the ER lumen activates a signal transduction system that connects the ER lumen with the cytoplasm and nucleus, a process called the unfolded protein response (UPR). In mammalian cells, this signaling pathway is transduced through three branches of ER stress sensors: the ER-associated receptors PERK (protein kinase RNA-like ER kinase), IRE1 (inositol-requiring protein-1) and ATF6 (activating transcription factor-6).^{5,1} The activation mechanism of these proteins has not been completely resolved, but it is known that certain molecular chaperones in the ER

transducers. Under normal conditions, the molecular chaperone BiP is bound to the luminal domain of these receptors, rendering them inactive. The accumulation of misfolded proteins in the ER increases the need for the molecular chaperone activity of BiP, and thus, upon ER stress, BiP is released from these receptors. BiP dissociation causes the activation of the three transducers. The activation of PERK suppresses protein synthesis through the phosphorylation of translation initiation factor 2 α (IF2 α). Upon BiP release, Ire1 undergoes dimerization, which sequentially activates its kinase and ribonuclease activity to induce the spliceosome-independent splicing of XBP1 (X-box binding protein-1) mRNA to generate mature mRNA that encode an active transcription factor. The activation of ATF6 promotes its translocation to the Golgi, where it is specifically cleaved by the proteolytic enzymes SP1 and SP2. Upon proteolytic hydrolysis of the transmembrane, the transcriptional activation domain of ATF6 detaches from the membrane and is directed to the nucleus. In the nucleus, XBP1 and ATF6 act in concert to activate the expression of ER-resident molecular chaperones, foldases and components of the ER-associated degradation (ERAD) system to increase the protein folding and processing capacity of the ER under stress conditions. Thus, on a physiological level, the activation of UPR triggers three protective cellular responses: (i) attenuation of protein translation through the PERK-mediated phosphorylation of IF2 α , (ii) upregulation of ER chaperones and (iii) degradation of misfolded proteins by the proteasome (ERAD). However, the failure to restore ER homeostasis under stress conditions results in the activation of programmed cell death signaling pathways as an ultimate attempt for survival. In plants, the UPR seems to operate as a bipartite module, as the ER stress signal is transduced through homologs of the IRE1 and ATF6 transducers, but a PERK-mediated branch of the UPR has not been shown.

In Arabidopsis, two IRE1-related genes, IRE1a and IRE1b, encode transmembrane domain-containing proteins that function as ER stress sensors or transducers. The GFP fusions of these receptors have been localized to the perinuclear ER, and the luminal sensor domain of these receptors can functionally replace the yeast IRE1 gene in complementation assays.⁶ Furthermore, both IRE1a and IRE1b display ribonuclease activity in vitro,⁷ and the functional characterization of an *atire1a atire1b* double mutant has revealed an essential and overlapping role for AtIRE1A and AtIRE1B as plant UPR regulators.⁸ Recently, a second component of this branch of ER stress signaling has been identified as the ER membrane-associated transcription factor bZIP60.^{7,9} Under normal conditions, the transcription factor bZIP60 is associated with microsomes,

and activates the stress-inducible BiP3 promoter.¹⁰ More recently, it has been demonstrated that upon ER stress, bZIP60 mRNA is spliced in an IRE1b-mediated process to generate an alternatively spliced transcript that lacks the transmembrane domain-encoding sequences.^{7,9} This splicing leads to the synthesis of a soluble and functional bZIP60 transfactor that can be translocated to the nucleus, where it activates ER stress inducible promoters, such as the BiP3 promoter. Likewise, OsbZIP74 or OsbZIP50 from rice, an ortholog of Arabidopsis AtbZIP60, is regulated through the IRE1-mediated splicing of its RNA to render the activation of ER stress-inducible promoters.^{11,12}

The second branch of UPR in plants mechanistically resembles the ATF6-mediated transduction of the ER stress signal. The membrane-associated Arabidopsis ATF6 homologs are bZIP17 and bZIP28, and they have canonical S1P sites in their C-terminal tails that face the ER lumen and appear to be processed by S1P and S2P. Upon ER stress, bZIP17 and zZIP28 are relocated to the Golgi, where their transcriptional domains are proteolytically released from the membrane by SP2.¹³ The transcriptional activator components of these transmembrane transfactors are then translocated to the nucleus, where they promote the expression of stress response genes.^{14-16,2} Thus, the bipartite cellular response to ER stress in plants is mediated through Ire1-like receptors and ATAF6 analog transducers to induce ER-resident molecular chaperones and the ER-associated protein degradation machinery as a protective measure. However, if ER stress is sustained, an apoptotic pathway is activated. Nevertheless, little is known about the pathways that can contribute to ER stress-induced cell death in plants. Recently, an ER membrane-associated G β -G γ heterodimer protein has been shown to be involved in the signaling events that trigger UPR-associated cell death in Arabidopsis.¹⁷ The study demonstrates that inactivation of the G β subunit (AGB1) of the heterotrimeric G protein protects plant cells against cell death induced by tunicamycin (Tm), a potent inducer of ER stress. Discrepant results were later reported with the demonstration that an AGB1 loss-of-function mutation causes over-sensitivity to ER stress and enhances the Tm-sensitive phenotype in the *atire1a atire1b* double mutant.⁸ Although the involvement of the GTP-binding protein β 1 (AGB1) in mediating ER stress-induced cell death is still a matter of debate, the latter study clearly demonstrated through reverse genetics that AtIRE1 and AGB1 independently and antagonistically control two essential plant UPR pathways.

UPR mediated cell death signaling is a distinct plant specific branch of the ER stress

and osmotic stress signals into a full response.¹⁸ This integrative pathway was first identified through genome-wide approaches and expression profiling, which revealed the existence of a modest overlap of the ER and osmotic stress-induced transcriptomes in soybean seedlings treated with PEG (an inducer of osmotic stress) or tunicamycin and AZC (potent inducers of ER stress).¹⁸ The co-regulated genes were first considered to be downstream targets of the integrated pathway based on similar induction kinetics and a synergistic response to the combination of osmotic and ER stress-inducing treatments. Based on these criteria, the selected downstream components of this ER and osmotic stress response-integrating pathway demonstrated the strongest synergistic induction, encoding proteins with diverse roles, such as plant-specific development and cell death (DCD) domain-containing proteins (NRP-A and NRP-B), an ubiquitin-associated (UBA) protein homolog and NAC domain-containing proteins (GmNAC6). NRP-A and NRP-B share a highly conserved C-terminal DCD (development and cell death) domain and possess a high number of asparagine residues at their more divergent N terminus.¹⁹ NRPs are critical mediators of ER and osmotic stress-induced cell death in soybeans.⁴ The cell death response mediated by NRPs resembles a programmed cell death event. The overexpression of NRPs in soybean protoplasts induces caspase-3-like activity and promotes extensive DNA fragmentation. Furthermore, the transient expression of NRPs in plants causes leaf yellowing, chlorophyll loss, malondialdehyde production, ethylene evolution and the induction of senescence marker genes, which are hallmarks of leaf senescence.

Similar to NRPs, GmNAC6 is another target of the integrated pathway that is strongly induced by cycloheximide, a potent inducer of cell death in soybean suspension cells. It is synergistically activated by a combination of ER and osmotic stress signals, and induces a senescence-like response in planta and cell death in soybean protoplasts.^{20,21} GmNAC6 belongs to a class of NAC (NAM, ATAF1, ATAF2 and CUC2) domain-containing proteins, which make up a large family of plant-specific transcription factor genes represented by at least 101 sequences in the soybean genome that are clustered into 15 different sub-families.²⁰ Members of this family are involved in the development and stress response. The NAC transactors are organized into a general structure that consists of a highly conserved N-terminal domain involved in DNA binding (called the NAC domain) and a C-terminal region that is highly divergent in sequence and length, which functions as the activation domain.²² NRPs and GmNAC6 are coordinately regulated through a variety of biotic and abiotic stresses, but induction of NRPs

expression of NRPs activates the GmNAC6 promoter and induces GmNAC6 expression, suggesting that GmNAC6 is located downstream of NRP in the ER and osmotic stress-induced cell death response.²¹

An upstream component of the NRP-mediated cell death response, GmERD15 (Glycine max Early Responsive to Dehydration 15), has been recently identified using one-hybrid screening that targeted the NRP-B promoter in yeast.²³ GmERD15 is induced by ER and osmotic stress to activate the expression of NRP genes. GmERD15 binds to and activates the NRP-B promoter in vitro and in vivo, exhibits transcriptional activity, is localized in the nucleus and induces the expression of the NRP-B gene when transiently expressed in soybean protoplasts. GmERD15 belongs to a class of ssDNA binding proteins and specifically recognizes the 12-bp palindromic sequence -511AGCAnnnnTGCT-500 in both SS (single-stranded) and DS (double-stranded) configurations of the NRP-B promoter. As an upstream component of ER stress-induced NRP-mediated signaling, GmERD15 associates stress in the ER with an osmotic stress-induced cell death signal.

The binding protein inhibits NRP-mediated cell death signaling and confers tolerance to drought

Although the activation mechanism of the three UPR transducers has not been totally deciphered in mammalian cells, the molecular chaperone BiP has been shown to play a pivotal role in controlling the activation status of IRE1, PERK and ATF6.^{24,1} Because BiP is the sole molecular chaperone involved in the activation of UPR, changes in its expression might indicate changes in the folding environment and ER processing capacity. Accordingly, the overexpression of BiP in mammals attenuates ER stress and suppresses the activation of UPR.²⁵ Mammalian BiP also exhibits protective properties that prevent oxidative stress, Ca²⁺ disturbances and cell death.²⁶⁻²⁹ In plants, there is no compelling evidence that BiP functions as a regulator of UPR transducers. Nevertheless, the overexpression of BiP in tobacco and soybean attenuates the activation of the UPR, suggesting a role for BiP in controlling the activation of UPR.^{30,4} More recently, it has been shown that the BiP-mediated inhibition of UPR activation appears to be dependent on the duration and intensity of the ER stress, such that BiP inhibition seems to be relieved if the stress persists.³¹ This observation might explain

In addition to alleviating ER stress,³⁰ the overexpression of BiP in plants has also been shown to increase their tolerance to water deficits.³²⁻³⁴ The apparent increase in BiP-mediated drought tolerance was not associated with typical short and long-term avoidance responses or with other known tolerance mechanisms.³⁴ The only variations observed in the BiP-overexpressing (OE) lines are a delay in drought-induced leaf senescence and an inhibition of the drought-mediated downregulation of ER molecular chaperone transcripts that occurs under prolonged osmotic stress. Recently, it has been shown that the enhanced expression of BiP in soybean (*Glycine max*) attenuated ER and osmotic-stress-mediated cell death.³¹ In transgenic lines, BiP overexpression attenuates ER and osmotic stress-induced cell death phenotypes, such as foliar necrotic lesions and wilting, percentage of dead cells, induction of senescence-associated gene markers, DNA fragmentation, caspase activity, lipid peroxidation and induction of the cell death marker genes NRP-A, NRP-B and GmNAC6. Accordingly, the prosurvival effect of BiP was associated with the modulation of the ER and osmotic stress-induced NRP-mediated cell death signaling, as determined in transgenic tobacco lines with enhanced and suppressed BiP levels. The enhanced expression of BiP prevented NRP- and NAC6-mediated cell death, whereas the silencing of endogenous BiP accelerated the onset of leaf senescence mediated through the ectopic expression of NRPs and GmNAC6 in tobacco leaves. These results implicate BiP as a negative regulator of the stress-induced NRP-mediated cell death response. Thus, it is not surprising that the overexpression of BiP delays drought-induced senescence in tobacco and soybean plants and confers the increased adaptation of these transgenic lines under water deprivation conditions compared with wild-type controls.

NRP-mediated cell death signaling represents a general plant response to multiple stresses

The components of the NRP-mediated cell death response, which include GmERD15, NRP-A, NRP-B and GmNAC6, are coordinately induced by several biotic and abiotic stimuli.^{20,21} It is likely that the NRP-mediated cell death signaling pathway represents a common response of plant cells to a variety of different stimuli. At least three lines of evidence implicate the stress-induced NRP-mediated cell death response as part of the hypersensitive response elicited by pathogen incompatibility interactions. First, the expression of both NRPs and GmNAC6 is induced through incompatibility interactions

necrotic lesions, resembling those of the hypersensitive response phenotype, and induces the expression of pathogenesis-related genes.²¹ Finally, ERD15, the transcriptional activator of NRP expression, was first described in Arabidopsis as a dehydration-induced gene³⁶ that functions as a negative regulator of the abscisic acid (ABA)-mediated response and a positive regulator of the salicylic acid (SA)-dependent defense pathway.³⁷

Conclusions

Since the discovery of the stress-induced NRP-mediated cell death response, many aspects of this signaling pathway induced by prolonged ER and osmotic stress have been elucidated. We now know that this pathway integrates the ER and osmotic-stress signals to synergistically increase the expression of N-rich proteins (NRP-A and NRP-B) and the NAC domain-containing protein GmNAC6, which are critical mediators of stress-induced cell death in plants. We also know that NRP-B expression is controlled by the novel ER and osmotic stress-induced transcriptional factor GmERD15 (Glycine max Early Responsive to Dehydration 15). However, several key players of this stress-induced signaling pathway are unknown, and many questions remain unanswered. What is the ER receptor that molecularly links the ER stress signal with the cell death response? How is the cell death signal propagated from induction of the plasma membrane-associated NRPs to the activation of the GmNAC6 promoter? What are the molecular events downstream of GmNAC6 that promote the cell death response? The synergistic induction of NRPs and NAC6 expression through the combination of ER and osmotic stress inducers indicates that the osmotic and ER stress-mediated cell death signals are transduced through separate pathways that converge on NRP expression. Which upstream components of NRPs are activated by the osmotic stress signal? The identification of NRP-interacting partners and downstream targets of the GmNAC6 transcriptional factor combined with the use of reverse genetics to examine plant ER stress transducers will be crucial to decipher this stress-induced cell death signaling response.

Recent data show that BiP, an ER molecular chaperone, acts as a negative regulator of NRP-mediated cell death signaling and that the manipulation of BiP expression protects plants against drought. We propose that the underlying mechanism of BiP-mediated

to modulate the osmotic stress-induced NRP-mediated cell death response. The positive effect of the modulation of the NRP-mediated cell death pathway on plant adaptation to stress might implicate this pathway as an excellent target for engineering superior crops. Because the NRP-mediated cell death signaling pathway represents a shared response to multiple stress signals in plants, it might permit coordinate adaptive cellular responses under a large array of stress conditions.

References

1. Malhotra JD, Kaufman RJ. The endoplasmic reticulum and the unfolded protein response. *Semin Cell Dev Biol* 2007; 18:716 - 31; <http://dx.doi.org/10.1016/j.semcdb.2007.09.003>; PMID: 18023214

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

2. Liu JX, Howell SH. Endoplasmic Reticulum Protein Quality Control and Its Relationship to Environmental Stress Responses in Plants. *Plant Cell* 2010; 22: 1–13. PMID: 20876830; DOI: 10.1105/tpc.110.078154.

[Google Scholar](#)

3. Moreno AA, Orellana A. The physiological role of the unfolded protein response in plants. *Biol Res* 2011; 44:75 - 80; <http://dx.doi.org/10.4067/S0716-97602011000100010>; PMID: 21720684

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

4. Costa MDL, Reis PAB, Valente MAS, Irsigler AST, Carvalho CM, Loureiro ME, et al. A new branch of endoplasmic reticulum stress signaling and the osmotic signal converge on plant-specific asparagine-rich proteins to promote cell death. *J Biol Chem* 2008; 283:20209 - 19; <http://dx.doi.org/10.1074/jbc.M802654200>; PMID: 18490446

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

5. Ron D, Walter P. Signal integration in the endoplasmic reticulum unfolded protein response. *Nat Rev Mol Cell Biol* 2007; 8:519 - 29; <http://dx.doi.org/10.1038/nrm2199>; PMID: 17565364

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

6. Koizumi N, Martine IM, Kimata Y, Kohno K, Sano H, Chrispeels MJ. Molecular characterization of two Arabidopsis Ire1 homologs, endoplasmic reticulum-located transmembrane protein kinases. *Plant Physiology* 2001; 127: 949-962. PMID: 11706177; DOI: 10.1104/pp.010636.

[Google Scholar](#)

7. Deng Y, Humbert S, Liu JX, Srivastava R, Rothstein SJ, Howell SH. Heat induces the splicing by IRE1 of a mRNA encoding a transcription factor involved in the unfolded protein response in Arabidopsis. *Proc Natl Acad Sci U S A* 2011; 108:7247 - 52; <http://dx.doi.org/10.1073/pnas.1102117108>; PMID: 21482766

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

8. Chen Y, Brandizzi F. AtIRE1A/AtIRE1B and AGB1 independently control two essential unfolded protein response pathways in Arabidopsis. *Plant J* 2012; 69:266 - 77; <http://dx.doi.org/10.1111/j.1365-313X.2011.04788.x>; PMID: 21914012

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

9. Nagashima Y, Mishiba K, Suzuki E, Shimada Y, Iwata Y and Koizumi N. Arabidopsis IRE1 catalyses unconventional splicing of bZIP60 mRNA to produce the active transcription factor. *Scientific reports* 2011; 1:29. PMID: 22355548; DOI: 10.1038/srep00029.

[Google Scholar](#)

10. Iwata Y, Fedoroff NV, Koizumi N. Arabidopsis bZIP60 is a proteolysis-activated transcription factor involved in the endoplasmic reticulum stress response. *Plant Cell* 2008; 20:3107 - 21; <http://dx.doi.org/10.1105/tpc.108.061002>; PMID: 19017746

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

1. Hayashi S, Wakasa Y, Takahashi H, Kawakatsu T, Takaiwa F. Signal transduction by IRE1-mediated splicing of bZIP50 and other stress sensors in the endoplasmic reticulum stress response of rice. *Plant J* 2012; 69:946 - 56; <http://dx.doi.org/10.1111/j.1365-313X.2011.04844.x>; PMID: 22050533

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

2. Lu S-J, Yang Z-T, Sun L, Sun L, Song Z-T, Liu J-X. Conservation of IRE1-Regulated bZIP74 mRNA Unconventional Splicing in Rice (*Oryza sativa* L.) Involved in ER Stress Responses. *Molecular Plant* 2011. PMID: 22199238; DOI:10.1093/mp/ssr115

[Google Scholar](#)

3. Che P, Bussell JD, Zhou W, Estavillo GM, Pogson BJ, Smith SM. Signaling from the endoplasmic reticulum activates brassinosteroid signaling and promotes acclimation to stress in *Arabidopsis*. *Sci Signal* 2010; 3:ra69; <http://dx.doi.org/10.1126/scisignal.2001140>; PMID: 20876872

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

4. Liu JX, Srivastava R, Che P, Howell SH. An endoplasmic reticulum stress response in *Arabidopsis* is mediated by proteolytic processing and nuclear relocation of a membrane-associated transcription factor, bZIP28. *Plant Cell* 2007; 19:4111-4119. PMID: 18156219; DOI: 10.1105/tpc.106.050021.

[Google Scholar](#)

5. Gao H, Brandizzi F, Benning C, Larkin RM. A membrane-tethered transcription factor defines a branch of the heat stress response in *Arabidopsis thaliana*. *Proc Natl Acad Sci U S A* 2008; 105:16398 - 403; <http://dx.doi.org/10.1073/pnas.0808463105>; PMID: 18849477

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

6. Tajima H, Iwata Y, Iwamoto M, Takayama S, Skoizumi. Identification of an *Arabidopsis* transmembrane bZIP transcription factor involved in the endoplasmic reticulum stress response. *Biochemical and Biophysical Research Communications* 2008; 374: 242-247. PMID: 18624751; DOI:10.1016/j.bbrc.2008.07.021

7. Wang S, Narendra S, Fedoroff N. Heterotrimeric G protein signaling in the Arabidopsis unfolded protein response. *Proc Natl Acad Sci U S A* 2007; 104:3817 - 22; <http://dx.doi.org/10.1073/pnas.0611735104>; PMID: 17360436

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

8. Irsigler AST, Costa MDL, Zhang P, Reis PAB, Dewey RE, Boston RS, et al. Expression profiling on soybean leaves reveals integration of ER- and osmotic-stress pathways. *BMC Genomics* 2007; 8:431; <http://dx.doi.org/10.1186/1471-2164-8-431>; PMID: 18036212

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

9. Tenhaken R, Doerks T, Bork P. DCD - a novel plant specific domain in proteins involved in development and programmed cell death. *BMC Bioinformatics* 2005; 6:169; <http://dx.doi.org/10.1186/1471-2105-6-169>; PMID: 16008837

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

10. Pinheiro GL, Marques CS, Costa MDBL, Reis PAB, Alves MS, Carvalho CM, et al. Complete inventory of soybean NAC transcription factors: sequence conservation and expression analysis uncover their distinct roles in stress response. *Gene* 2009; 444:10 - 23; <http://dx.doi.org/10.1016/j.gene.2009.05.012>; PMID: 19497355

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

11. Faria JAQA, Reis PAB, Reis MTB, Rosado GL, Pinheiro GL, Mendes GC, et al. The NAC domain-containing protein, GmNAC6, is a downstream component of the ER stress- and osmotic stress-induced NRP-mediated cell-death signaling pathway. *BMC Plant Biol* 2011; 11:129; <http://dx.doi.org/10.1186/1471-2229-11-129>; PMID: 21943253

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

12. Hegedus D, Yu M, Baldwin D, Gruber M, Sharpe A, Parkin I, et al. Molecular characterization of Brassica napus NAC domain transcriptional activators induced in

response to biotic and abiotic stress. *Plant Mol Biol* 2003; 53:383 - 97;
<http://dx.doi.org/10.1023/B:PLAN.0000006944.61384.11>; PMID: 14750526

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

23. Alves MS, Reis PAB, Dadalto SP, Faria JAQA, Fontes EPB, Fietto LG. A novel transcription factor, early responsive to dehydration 15, connects ER stress with an osmotic stress-induced cell death signal. *J Biol Chem* 2010; 286:20020 - 30;
<http://dx.doi.org/10.1074/jbc.M111.233494>

[Web of Science ®](#) | [Google Scholar](#)

24. Bertolotti A, Zhang Y, Hendershot LM, Harding HP, Ron D. Dynamic interaction of BiP and ER stress transducers in the unfolded-protein response. *Nat Cell Biol* 2000; 2:326 - 32; <http://dx.doi.org/10.1038/35014014>; PMID: 10854322

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

25. Morris JA, Dorner AJ, Edwards CA, Hendershot LM, Kaufman RJ. Immunoglobulin binding protein (BiP) function is required to protect cells from endoplasmic reticulum stress but is not required for the secretion of selective proteins. *J Biol Chem* 1997; 272:4327 - 34; <http://dx.doi.org/10.1074/jbc.272.7.4327>; PMID: 9020152

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

26. Liu H, Bowes RC 3rd, van de Water B, Sillence C, Nagelkerke JF, Stevens JL. Endoplasmic reticulum chaperones GRP78 and calreticulin prevent oxidative stress, Ca²⁺ disturbances, and cell death in renal epithelial cells. *J Biol Chem* 1997; 272:21751 - 9; <http://dx.doi.org/10.1074/jbc.272.35.21751>; PMID: 9268304

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

27. Liu H, Miller E, van de Water B, Stevens JL. Endoplasmic reticulum stress proteins block oxidant-induced Ca²⁺ increases and cell death. *J Biol Chem* 1998; 273:12858 - 62; <http://dx.doi.org/10.1074/jbc.273.21.12858>; PMID: 9582315

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

8. Gething MJ. Role and regulation of the ER chaperone BiP. *Seminars in Cell and Developmental Biology*.1999; 10: 465-472. PMID:10597629: DOI: 10.1006/scdb.1999.0318.

[Google Scholar](#)

9. Kishi S, Shimoke K, Nakatani Y, Shimada T, Okumura N, Nagai K, et al. Nerve growth factor attenuates 2-deoxy-d-glucose-triggered endoplasmic reticulum stress-mediated apoptosis via enhanced expression of GRP78. *Neurosci Res* 2010; 66:14 - 21; <http://dx.doi.org/10.1016/j.neures.2009.09.003>; PMID: 19766678

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

10. Leborgne-Castel N, Jelitto-Van Dooren EPWM, Crofts AJ, Denecke J. Overexpression of BiP in tobacco alleviates endoplasmic reticulum stress. *Plant Cell* 1999; 11: 459-469. PMID: 10072404: DOI: 10.1105/tpc.11.3.459.

[Google Scholar](#)

11. Reis PAB, Rosado GL, Silva LAC, Oliveira LC, Oliveira LB, Costa MDL, et al. The binding protein BiP attenuates stress-induced cell death in soybean via modulation of the N-rich protein-mediated signaling pathway. *Plant Physiology* 2011; 157:1853-1865. PMID: 22007022: DOI: 10.1104/pp.111.179697.

[Google Scholar](#)

12. Cascardo JCM, Almeida RS, Buzeli RAA, Carolino SMB, Otoni WC, Fontes EPB. The phosphorylation state and expression of soybean BiP isoforms are differentially regulated following abiotic stresses. *J Biol Chem* 2000; 275:14494 - 500; <http://dx.doi.org/10.1074/jbc.275.19.14494>; PMID: 10799532

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

13. Alvim FC, Carolino SMB, Cascardo JCM, Nunes CC, Martinez CA, Otoni WC, et al. Enhanced accumulation of BiP in transgenic plants confers tolerance to water stress. *Plant Physiology* 2001; 126: 1042-1054. PMCID: PMC116461: DOI: 10.1104/pp.126.3.1042.

34. Valente MAS, Faria JAQA, Soares-Ramos JR, Reis PAB, Pinheiro GL, Piovesan ND, et al. The ER luminal binding protein (BiP) mediates an increase in drought tolerance in soybean and delays drought-induced leaf senescence in soybean and tobacco. *J Exp Bot* 2009; 60:533 - 46; <http://dx.doi.org/10.1093/jxb/ern296>; PMID: 19052255

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

35. Ludwig AA, Tenhaken R. A new cell wall located N-rich protein is strongly induced during the hypersensitive response in *Glycine max* L. *Eur J Plant Pathol* 2001; 107:323 - 36; <http://dx.doi.org/10.1023/A:1011202225323>

[Web of Science ®](#)

[Google Scholar](#)

36. Kiyosue T, Yamaguchi-Shinozaki K, Shinozaki K. ERD15, a cDNA for a dehydration-induced gene from *Arabidopsis thaliana*. *Plant Physiol* 1994; 106:1707; <http://dx.doi.org/10.1104/pp.106.4.1707>; PMID: 7846179

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

37. Kariola T, Brader G, Helenius E, Li J, Heino P, Palva ET. EARLY RESPONSIVE TO DEHYDRATION 15, a Negative Regulator of Abscisic Acid Responses in *Arabidopsis*. *Plant Physiology*.2006; 142: 1559–1573. PMID: 17056758: DOI: 10.1105/tpc.108.061002.

[Google Scholar](#)

[Download PDF](#)



Asparagine-rich protein (NRP) mediates stress response by regulating biosynthesis of plant secondary metabolites in Arabidopsis >

Kaikai Zhu et al.

Plant Signaling & Behavior

Published online: 29 Jul 2023



Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG

 Taylor and Francis
Group

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)