



Free access

1,317 24

Views CrossRef citations to date Altmetric

1

Listen

Mini Review

N-rich protein (NRP)-mediated cell death signaling

A new branch of the ER stress response with implications for plant biotechnology

Pedro A.B. Reis & Elizabeth P.B. Fontes

Pages 628-632 | Published online: 14 May 2012

Cite this article <https://doi.org/10.4161/psb.20111>

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

View PDF

Share

We Care About Your Privacy

We and our 909 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting I Accept enables tracking technologies to support the purposes shown under we and our partners process data to provide. Selecting Reject All or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the Show Purposes link on the bottom of the webpage. Your choices will have effect within our Website. For more details, refer to our Privacy Policy. [Here](#)

We and our partners process data to provide:

Use precise geolocation data. Actively scan device

I Accept

Reject All

Show Purposes



attenuates dehydration-induced cell death and promotes a better adaptation of BiP-overexpressing transgenic lines to drought.

Keywords: :

- ERD15
- GmNAC6
- N-rich proteins
- NRP-A
- NRP-B
- cell death response
- endoplasmic reticulum stress
- osmotic stress

The endoplasmic reticulum is a key signaling organelle involved in the activation of cellular stress responses in eukaryotic cells. One such well-characterized signaling event is the unfolded protein response (UPR), which is activated to cope with the disruption of ER homeostasis that results in the accumulation of unfolded or misfolded proteins in the lumen of the organelle. In mammalian cells, the UPR has been associated with other stress response pathways through shared components.¹ In plants, the potential of the ER stress response to accommodate adaptive pathways and its connection with other environmentally induced responses have been the subjects of studies in recent years.^{2,3} One plant-specific, ER stress-shared response is ER and osmotic stress-integrated signaling, which converges on N-rich proteins (NRPs) to transduce signals and activate cell death signaling pathways. For full activation of the UPR, the ER stress response must be integrated with other signaling pathways.

ER str



Any of the misfolded proteins that connect to the protein network through the kinase R, activating transcription factors, not been the ER

transducers. Under normal conditions, the molecular chaperone BiP is bound to the luminal domain of these receptors, rendering them inactive. The accumulation of misfolded proteins in the ER increases the need for the molecular chaperone activity of BiP, and thus, upon ER stress, BiP is released from these receptors. BiP dissociation causes the activation of the three transducers. The activation of PERK suppresses protein synthesis through the phosphorylation of translation initiation factor 2 α (IF2 α). Upon BiP release, Ire1 undergoes dimerization, which sequentially activates its kinase and ribonuclease activity to induce the spliceosome-independent splicing of XBP1 (X-box binding protein-1) mRNA to generate mature mRNA that encode an active transcription factor. The activation of ATF6 promotes its translocation to the Golgi, where it is specifically cleaved by the proteolytic enzymes SP1 and SP2. Upon proteolytic hydrolysis of the transmembrane, the transcriptional activation domain of ATF6 detaches from the membrane and is directed to the nucleus. In the nucleus, XBP1 and ATF6 act in concert to activate the expression of ER-resident molecular chaperones, foldases and components of the ER-associated degradation (ERAD) system to increase the protein folding and processing capacity of the ER under stress conditions. Thus, on a physiological level, the activation of UPR triggers three protective cellular responses: (i) attenuation of protein translation through the PERK-mediated phosphorylation of IF2 α , (ii) upregulation of ER chaperones and (iii) degradation of misfolded proteins by the proteasome (ERAD). However, the failure to restore ER homeostasis under stress

condition
ultimate
module,
transduc

In Arabid
domain-
fusions
sens
complex
activity
mutant
UPR reg
has been
Under no

ways as an
rtite
and ATF6

brane
s. The GFP
the luminal
ene in
nuclease
ouble
E1B as plant
signaling
P60.^{7,9}
microsomes,



Similar to... induced... is synergic... induces... GmNAC6... proteins... represent... 15 different... stress response... consists... NAC domain... which function... regulate

s strongly... sion cells. It... ls, and... toplasts.[20,21](#)... containing... genes... stered into... opment and... e that... called the... nd length,... ately... RPs



expression of NRPs activates the GmNAC6 promoter and induces GmNAC6 expression, suggesting that GmNAC6 is located downstream of NRP in the ER and osmotic stress-induced cell death response.^{[21](#)}

An upstream component of the NRP-mediated cell death response, GmERD15 (Glycine max Early Responsive to Dehydration 15), has been recently identified using one-hybrid screening that targeted the NRP-B promoter in yeast.^{[23](#)} GmERD15 is induced by ER and osmotic stress to activate the expression of NRP genes. GmERD15 binds to and activates the NRP-B promoter in vitro and in vivo, exhibits transcriptional activity, is localized in the nucleus and induces the expression of the NRP-B gene when transiently expressed in soybean protoplasts. GmERD15 belongs to a class of ssDNA binding proteins and specifically recognizes the 12-bp palindromic sequence -511AGCAnnnnTGCT-500 in both SS (single-stranded) and DS (double-stranded) configurations of the NRP-B promoter. As an upstream component of ER stress-induced NRP-mediated signaling, GmERD15 associates stress in the ER with an osmotic stress-induced cell death signal.

The binding protein inhibits NRP-mediated cell death signaling and confers tolerance to drought

Although the mechanism of NRP-mediated cell death signaling has not been totally deciphered, it is believed to play a pivotal role in the ER stress response. Because BiP is the so-called chaperone protein in its capacity to bind to and suppress the aggregation of misfolded proteins, it is thought that BiP has no competitive effect on NRP-mediated cell death signaling. Nevertheless, the activation of NRP-mediated cell death signaling is not the only pathway of UPR.^{[30,4](#)} More recent studies have shown that BiP appears to inhibit NRP-mediated cell death signaling, which might explain



In addition to alleviating ER stress,³⁰ the overexpression of BiP in plants has also been shown to increase their tolerance to water deficits.³²⁻³⁴ The apparent increase in BiP-mediated drought tolerance was not associated with typical short and long-term avoidance responses or with other known tolerance mechanisms.³⁴ The only variations observed in the BiP-overexpressing (OE) lines are a delay in drought-induced leaf senescence and an inhibition of the drought-mediated downregulation of ER molecular chaperone transcripts that occurs under prolonged osmotic stress. Recently, it has been shown that the enhanced expression of BiP in soybean (*Glycine max*) attenuated ER and osmotic-stress-mediated cell death.³¹ In transgenic lines, BiP overexpression attenuates ER and osmotic stress-induced cell death phenotypes, such as foliar necrotic lesions and wilting, percentage of dead cells, induction of senescence-associated gene markers, DNA fragmentation, caspase activity, lipid peroxidation and induction of the cell death marker genes NRP-A, NRP-B and GmNAC6. Accordingly, the prosurvival effect of BiP was associated with the modulation of the ER and osmotic stress-induced NRP-mediated cell death signaling, as determined in transgenic tobacco lines with enhanced and suppressed BiP levels. The enhanced expression of BiP prevented NRP- and NAC6-mediated cell death, whereas the silencing of endogenous BiP accelerated the onset of leaf senescence mediated through the ectopic expression of NRPs and GmNAC6 in tobacco leaves. These results implicate BiP as a negative regulator of the stress-induced NRP-mediated cell death response. Thus, it is not surprising that the overexpression of BiP in soybean plants and tobacco plants under water deprivation

NRP-mediated
response

The
NRP-A, N
stimuli.²
common
evidence
hyperse
expressi

ant

mERD15,
abiotic
represents a
ee lines of
s part of the
first, the
interactions



necrotic lesions, resembling those of the hypersensitive response phenotype, and induces the expression of pathogenesis-related genes.²¹ Finally, ERD15, the transcriptional activator of NRP expression, was first described in Arabidopsis as a dehydration-induced gene³⁶ that functions as a negative regulator of the abscisic acid (ABA)-mediated response and a positive regulator of the salicylic acid (SA)-dependent defense pathway.³⁷

Conclusions

Since the discovery of the stress-induced NRP-mediated cell death response, many aspects of this signaling pathway induced by prolonged ER and osmotic stress have been elucidated. We now know that this pathway integrates the ER and osmotic-stress signals to synergistically increase the expression of N-rich proteins (NRP-A and NRP-B) and the NAC domain-containing protein GmNAC6, which are critical mediators of stress-induced cell death in plants. We also know that NRP-B expression is controlled by the novel ER and osmotic stress-induced transcriptional factor GmERD15 (Glycine max Early Responsive to Dehydration 15). However, several key players of this stress-induced signaling pathway are unknown, and many questions remain unanswered.

What is the ER receptor that molecularly links the ER stress signal with the cell death response? What are the membrane proteins that are the molecular targets of ER and osmotic stress signals? The synergistic action of ER and osmotic stress signals in cell death response? Which upstream signaling molecules are involved in the ER stress response? The identification of NAC6 as a plant ER stress response mediator is a significant step forward in understanding the plant ER signaling pathway.

Recent studies have shown that NAC6 is a positive regulator of the cell death response. NAC6 expression protects plants against ER stress-induced cell death mediated



to modulate the osmotic stress-induced NRP-mediated cell death response. The positive effect of the modulation of the NRP-mediated cell death pathway on plant adaptation to stress might implicate this pathway as an excellent target for engineering superior crops. Because the NRP-mediated cell death signaling pathway represents a shared response to multiple stress signals in plants, it might permit coordinate adaptive cellular responses under a large array of stress conditions.

References

1. Malhotra JD, Kaufman RJ. The endoplasmic reticulum and the unfolded protein response. *Semin Cell Dev Biol* 2007; 18:716 - 31;
<http://dx.doi.org/10.1016/j.semcdb.2007.09.003>; PMID: 18023214
[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)
2. Liu JX, Howell SH. Endoplasmic Reticulum Protein Quality Control and Its Relationship to Environmental Stress Responses in Plants. *Plant Cell* 2010; 22: 1–13. PMID: 20876830; DOI: 10.1105/tpc.110.078154.

Google

3. Moren... response in
plants
97602

4. Co... et al. A
new b... signal
conve... J Biol
Chem... PMID:
18490



- PubMed Web of Science ® Google Scholar

- Google Scholar

- PubMed Web of Science ® Google Scholar

- 

- | | |
|----------|---------------|
| 0. Iwata | ated |
| transc | e. Plant Cell |
| 2008; | 17746 |

1. Hayashi S, Wakasa Y, Takahashi H, Kawakatsu T, Takaiwa F. Signal transduction by IRE1-mediated splicing of bZIP50 and other stress sensors in the endoplasmic reticulum stress response of rice. *Plant J* 2012; 69:946 - 56;
<http://dx.doi.org/10.1111/j.1365-313X.2011.04844.x>; PMID: 22050533

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

2. Lu S-J, Yang Z-T, Sun L, Sun L, Song Z-T, Liu J-X. Conservation of IRE1-Regulated bZIP74 mRNA Unconventional Splicing in Rice (*Oryza sativa* L.) Involved in ER Stress Responses. *Molecular Plant* 2011. PMID: 22199238; DOI:10.1093/mp/ssr115

[Google Scholar](#)

3. Che P, Bussell JD, Zhou W, Estavillo GM, Pogson BJ, Smith SM. Signaling from the endoplasmic reticulum activates brassinosteroid signaling and promotes acclimation to stress in *Arabidopsis*. *Sci Signal* 2010; 3:ra69;
<http://dx.doi.org/10.1126/scisignal.2001140>; PMID: 20876872

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

4. Liu JX, Srivastava R, Che P, Howell SH. An endoplasmic reticulum stress response in *Arabidopsis* is mediated by proteolytic processing and nuclear relocation of a membrane protein. *Proc Natl Acad Sci U S A* 2011; 108:4111-4119. PMID: 21511111

[Google Scholar](#)

5. Gao H, Zhang Y, Zhang Y, Zhang Y, Zhang Y, Zhang Y. The transcription factor bZIP11 defines a novel endoplasmic reticulum stress response in *Arabidopsis*. *Sci U S A* 2011; 108:3105; PMID: 21511111



6. Tajima F, Zhang Y, Zhang Y, Zhang Y, Zhang Y, Zhang Y. The transcription factor bZIP11 defines a novel endoplasmic reticulum stress response in *Arabidopsis*. *Sci U S A* 2011; 108:3105; PMID: 21511111

17. Wang S, Narendra S, Fedoroff N. Heterotrimeric G protein signaling in the Arabidopsis unfolded protein response. *Proc Natl Acad Sci U S A* 2007; 104:3817 - 22; <http://dx.doi.org/10.1073/pnas.0611735104>; PMID: 17360436

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

18. Irsigler AST, Costa MDL, Zhang P, Reis PAB, Dewey RE, Boston RS, et al. Expression profiling on soybean leaves reveals integration of ER- and osmotic-stress pathways. *BMC Genomics* 2007; 8:431; <http://dx.doi.org/10.1186/1471-2164-8-431>; PMID: 18036212

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

19. Tenhaken R, Doerks T, Bork P. DCD - a novel plant specific domain in proteins involved in development and programmed cell death. *BMC Bioinformatics* 2005; 6:169; <http://dx.doi.org/10.1186/1471-2105-6-169>; PMID: 16008837

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

20. Pinheiro GL, Marques CS, Costa MDBL, Reis PAB, Alves MS, Carvalho CM, et al.

Comp xervation and
expres 2009;
444:1 55

21. Faria J The NAC
don ER stress-
an BMC Plant
Biol 20 943253

22. Hegec r
charac induced in



response to biotic and abiotic stress. Plant Mol Biol 2003; 53:383 - 97;
<http://dx.doi.org/10.1023/B:PLAN.00000006944.61384.11>; PMID: 14750526

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

23. Alves MS, Reis PAB, Dadalto SP, Faria JAQA, Fontes EPB, Fietto LG. A novel transcription factor, early responsive to dehydration 15, connects ER stress with an osmotic stress-induced cell death signal. J Biol Chem 2010; 286:20020 - 30;
<http://dx.doi.org/10.1074/jbc.M111.233494>

[Web of Science ®](#)

[Google Scholar](#)

24. Bertolotti A, Zhang Y, Hendershot LM, Harding HP, Ron D. Dynamic interaction of BiP and ER stress transducers in the unfolded-protein response. Nat Cell Biol 2000; 2:326 - 32; <http://dx.doi.org/10.1038/35014014>; PMID: 10854322

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

25. Morris JA, Dorner AJ, Edwards CA, Hendershot LM, Kaufman RJ. Immunoglobulin binding protein (BiP) function is required to protect cells from endoplasmic reticulum stress but is not required for the secretion of selective proteins. J Biol Chem 1997; 272:4327 - 34; <http://dx.doi.org/10.1074/jbc.272.7.4327>; PMID: 9020152

26. Liu H, ... L.
Endop ... ive stress,
Ca²⁺ ... 97;
272:2 ... 04



27. Liu H, ... proteins
block ... 273:12858 -
62; ht

8. Gething MJ. Role and regulation of the ER chaperone BiP. *Seminars in Cell and Developmental Biology*.1999; 10: 465-472. PMID:10597629: DOI: 10.1006/scdb.1999.0318.

[Google Scholar](#)

9. Kishi S, Shimoke K, Nakatani Y, Shimada T, Okumura N, Nagai K, et al. Nerve growth factor attenuates 2-deoxy-d-glucose-triggered endoplasmic reticulum stress-mediated apoptosis via enhanced expression of GRP78. *Neurosci Res* 2010; 66:14 - 21; <http://dx.doi.org/10.1016/j.neures.2009.09.003>; PMID: 19766678

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

10. Leborgne-Castel N, Jelitto-Van Dooren EPWM, Crofts AJ, Denecke J. Overexpression of BiP in tobacco alleviates endoplasmic reticulum stress. *Plant Cell* 1999; 11: 459-469. PMID: 10072404: DOI: 10.1105/tpc.11.3.459.

[Google Scholar](#)

11. Reis PAB, Rosado GL, Silva LAC, Oliveira LC, Oliveira LB, Costa MDL, et al. The binding protein BiP attenuates stress-induced cell death in soybean via modulation of the N-rich protein-mediated signaling pathway. *Plant Physiology* 2011; 157:1853-1865. PMID: 21510000

[Google Scholar](#)

12. Cascaes R, Reis PAB. The binding protein BiP attenuates stress-induced cell death in soybean via modulation of the N-rich protein-mediated signaling pathway. *Plant Physiology* 2011; 157:1853-1865. <http://dx.doi.org/10.1105/tpc.11.3.459>



13. Alvim M, Reis PAB, et al. The binding protein BiP attenuates stress-induced cell death in soybean via modulation of the N-rich protein-mediated signaling pathway. *Plant Physiology* 2011; 157:1853-1865. DOI: 10.1105/tpc.11.3.459

34. Valente MAS, Faria JAQA, Soares-Ramos JR, Reis PAB, Pinheiro GL, Piovesan ND, et al. The ER luminal binding protein (BiP) mediates an increase in drought tolerance in soybean and delays drought-induced leaf senescence in soybean and tobacco. J Exp Bot 2009; 60:533 - 46; <http://dx.doi.org/10.1093/jxb/ern296>; PMID: 19052255

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

35. Ludwig AA, Tenhaken R. A new cell wall located N-rich protein is strongly induced during the hypersensitive response in Glycine max L. Eur J Plant Pathol 2001; 107:323 - 36; <http://dx.doi.org/10.1023/A:1011202225323>

[Web of Science ®](#)

[Google Scholar](#)

36. Kiyosue T, Yamaguchi-Shinozaki K, Shinozaki K. ERD15, a cDNA for a dehydration-induced gene from Arabidopsis thaliana. Plant Physiol 1994; 106:1707; <http://dx.doi.org/10.1104/pp.106.4.1707>; PMID: 7846179

[PubMed](#)

[Web of Science ®](#)

[Google Scholar](#)

37. Kariola T, Brader G, Helenius E, Li J, Heino P, Palva ET. EARLY RESPONSIVE TO DEHYDRATION 15, a Negative Regulator of Absciscic Acid Responses in Arabidopsis. Plant Physiology. 2006; 142: 1559–1573. PMID: 17056758; DOI: 10.1104/pp.106.4.1707

[Google Scholar](#)

[Download](#)

[Related](#)



Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright

Accessib

Registered
5 Howick Pl

or & Francis Group
orma business

